Isolation and characterization of nine microsatellite markers for the red alga *Corallina officinalis*

Ana I. Tavares¹ · Katy R. Nicastro¹ · Regina Kolzenburg² · Federica Ragazzola² · Rita Jacinto¹ · Gerardo I. Zardi³

Abstract

We report the development of nine polymorphic microsatellite markers for *Corallina officinalis* (Linnaeus, 1758), a calcifying intertidal red alga and important ecosystem engineer spread along the North East Atlantic. Characterization and analysis of loci were made using 15 individuals of *C. officinalis* from populations in Iceland and the UK. The average number of alleles per locus was 3.78 (range 2–6) and mean of gene diversity was 0.58 (range 0.38–0.77). The set of microsatellites developed here will provide a useful molecular tool for population genetic and conservation studies.

Keywords *Corallina officinalis* · Genetic diversity · Microsatellites · Red algae · Marginal population

Corresponding author: Ana I. Tavares aintavares@ualg.pt

Introduction

Within the red algae, Corallinales (Rhodophyta) is one of the most species rich orders [1, 2]. Given their role as bioengineers and of their sensitivity to climate change and ocean acidification, species in the Corallinales are of great ecological relevance in various marine, wave-exposed littoral and sublittoral habitats [3, 4]. They are also abundant components in fossil and sub-fossil records, thus representing valuable indicators of paleoenvironmental conditions and providing important information to contextualize ongoing climatic changes [5–8]. *Corallina officinalis* Linnaeus, 1758 (Corallinales, Rhodophyta) is an abundant and widespread calcified geniculate coralline alga on rocky shores in the North Atlantic [9–11]. The distribution of *C. officinalis* along NE Atlantic shores extends from northern Spain and the Azores, in the south, to northern Norway, Iceland and southern Greenland, in the north [10, 12]. The dense turfs created by this species provide shelter in hydrodynamically stressful habitats for many invertebrates [13]. Moreover, *C. officinalis* contributes to carbon dioxide fluxes within the ocean, thus playing a critical role in the carbon cycle of coastal marine ecosystems [14].

Despite its ecological importance, genetic studies for this species are limited [15]. These have used mitochondrial and nuclear sequence data for barcoding, taxonomy, mitogenomics and phylogeny [2, 9, 10, 16–19]. Only very recently have population genetic patterns of *C. officinalis* been investigated [20, 21]. Specifically, novel SNP markers were developed and used to assess population structure based on five locations throughout the NE Atlantic distribution of the species and, at finer spatial scale, along UK shores. Here, with the aim of providing new useful tools using methods previously unutilised for this group, we developed and characterized nine polymorphic microsatellite markers for the red alga *C. officinalis*.

Methods

All samples were hand-collected during low tide from rocky substrate. Tissue was stored in silica drying crystals prior to extraction of DNA. Genomic DNA was extracted from eight individuals, four collected in Iceland (Stafnesviti 63°58'06.5"N 22°45'03.0"W and Stafnes 63°58'28.1"N 22°45'13.8"W) and four in the UK (Margate 51.393200°N 01.383820°E) using the PowerSoil DNA Isolation Kit (MoBio Laboratories, Carlsbad, CA) in accordance with the manufacturer's instructions. Extracted DNA was sent to Ecogenics GmbH molecular marker services (Zurich- Schlieren, Switzerland) to isolate and design microsatellite markers. Genomic DNA size-selected fragments were enriched for simple sequence repeat (SSRs) content by using magnetic streptavidin beads and biotin-labeled CT, GT, AGG, GGT GATA and GTAT repeat oligonucleotides. The SSR-enriched library was analyzed on an Illumina MiSeq platform using the Nano 2 × 250 v2 format. After assembly, 2444 contigs or singlets contained a microsatellite insert with a tetra- or a trinucleotide of at least six repeat units or a dinucleotide of at least ten repeat units. Out of 1129 microsatellite candidates, 20 were designed and tested. Out of these, nine pairs of primers showed polymorphism. The identified, polymorphic primers were further characterized using the eight and seven *C. officinalis* individuals from Icelandic and UK populations respectively (Table 1). The 10 µL volume PCR mix contained ± 10 ng of DNA, 0.3 µM of each primer labelled with a fluorescent marker, 200 µM dNTPs, 1.0 µL of 10× PCR Buffer with 15 mM MgCl₂ and 0.5 U of HotStar Taq. Cycling conditions consisted of an initial denaturing step of 15 min at 95 °C, followed by 35 cycles of 30 s at 95 °C, 45 s at 56 °C annealing temperature, 45 s at 72 °C, and a final elongation step at 72 °C for 30 min. The allele sizes were determined on an ABI3730 (Applied Biosystems), using GeneScan™-500 LIZ Size Standard. Alleles were scored manually using STRand [22]. GENETIX software version 4.05 [23] was used to estimate observed (HO) and expected (HE) heterozygosities. The Hardy–Weinberg expectation (HWE) and linkage disequilibrium were tested using Genepop 4.2 [24] and the resulting p values were Bonferroni-corrected for multiple comparisons ($p < 0.0016$).

Results and discussion

All microsatellites described in this study were polymorphic (Table 1). The number of alleles per locus ranged between 1–6 and 1–5 for the populations from Iceland and the UK, respectively. For the Icelandic populations, expected and observed heterozygosities ranged from 0.37 to 0.69 and from 0.25 to 0.88 respectively (Table 1) while for the UK populations, values ranged from 0.41 to 0.77 and from 0 to 0.86 (Table 1). The highest genetic diversity was found in the UK population, with Coroff_8333 being the most informative locus. The lowest genetic diversity values were given by Coroff_1036 and Coroff_1640 and by Coroff_1258 for Iceland and UK populations respectively. For all loci pairs, we tested for linkage disequilibrium between them, using 1000 replicates and for two pairs (Coroff_2347–Coroff_3261; Coroff_1640–Coroff_7707) there was no significant deviation after Bonferroni correction. Significant deviation from HWE was found for Coroff_11062 in the UK population. Relatively high values of observed heterozygosity recently reported for SNP markers [20] suggest polyploidy in *C. officinalis*. It is important to note that potential polyploidy could significantly affect our ability to reliably utilise codominant scoring and thus interpretation of peak patterns [25]. Most SNP markers recently developed for *C. officinalis* show significant departures from HWE [20]; in our study, significant deviation was

limited to one locus, highlighting the value of the set of microsatellites developed here to investigate ecological and evolutionary dynamics of this intertidal bioengineer associated with habitat contraction and environmental change.

Acknowledgements This research was funded by projects UID/ Multi/04326/2013, IF/01413/2014/CP1217/CT0004 from the Fundação para a Ciência e Tecnologia (FCT-MEC, Portugal) and RDF from University of Portsmouth and further supported by the South African Research Chairs Initiative (SARChI) of the Department of Science and Technology and the National Research Foundation and a scholarship from the South African National Research Foundation (NRF). We thank C.D. McQuaid and C. Yesson for their comments.

References

1. Guiry MD, Guiry GM (2017) AlgaeBase. World-wide electronic publication, National University of Ireland, Galway. <http://www.algae base.org>. Accessed 12 Dec 2017
2. Walker RH, Brodie J, Russell S, Irvine LM, Orfanidis S (2009) Biodiversity of coralline algae in the Northeastern Atlantic. *J Phycol* 297:287–297. <https://doi.org/10.1111/j.1529-8817.2008.0637>
3. Broom JE, Hart DR, Farr TJ, Nelson WA, Neill KF, Harvey AS, Woelkerling WJ (2008) Utility of psbA and nSSU for phylogenetic reconstruction in the Corallinales based on New Zealand taxa. *Mol Phylogenet Evol* 46:958–973. <https://doi.org/10.1016/j.ympev.2007.12.016>
4. McCoy SJ, Kamenos NA (2015) Coralline algae (Rhodophyta) in a changing world: integrating ecological, physiological, and geochemical responses to global change. *J Phycol* 51:6–24. <https://doi.org/10.1111/jpy.12262>
5. Braga JC, Aguirre J (2001) Coralline algal assemblages in upper Neogene reef and temperate carbonates in Southern Spain. *Palaeogeogr Palaeoclimatol Palaeoecol* 175:27–41. [https://doi.org/10.1016/S0031-0182\(01\)00384-4](https://doi.org/10.1016/S0031-0182(01)00384-4)
6. Abbey E, Webster JM, Braga JC, Sugihara K, Wallace C, Iryu Y, Potts D, Done T, Camoin G, Seard C (2011) Variation in deglacial coral algal assemblages and their paleoenvironmental significance: IODP Expedition 310, “Tahiti Sea Level”. *Glob Planet Change* 76:1–15. <https://doi.org/10.1016/j.gloplacha.2010.11.005>
7. Aguirre J, Braga JC, De Reviers B, Woelkerling WJ (2012) Reassessment of Lemoine’s newly discovered types of fossil corallines (Corallinales, Rhodophyta) preserved at the Muséum national d’histoire naturelle, Paris. *Cryptogam Algal* 33:289–326. <https://doi.org/10.7872/crya.v33.iss3.2012.289>
8. Novak V, Santodomingo N, Rösler A, Martino E, Braga JC, Taylor PD, Johnson KG, Renema W (2013) Environmental reconstruction of a late Burdigalian (Miocene) patch reef in deltaic deposits (East Kalimantan, Indonesia). *Palaeogeogr Palaeoclimatol Palaeoecol* 374:110–122. <https://doi.org/10.1016/j.palaeo.2013.01.009>
9. Brodie J, Walker RH, Williamson C, Irvine LM (2013) Epitypification and redescription of *Corallina officinalis* L., the type of the genus, and *C. elongata* Ellis et Solander (Corallinales, Rhodophyta). *Cryptogam Algal* 34:49–56. <https://doi.org/10.7872/crya.v34.iss1.2013.49>
10. Williamson CJ, Walker RH, Robba L, Yesson C, Russell S, Irvine LM, Brodie J (2015) Toward resolution of species diversity and distribution in the calcified red algal genera *Corallina* and *Ellisolandia* (Corallinales, Rhodophyta). *Phycologia* 54:2–11. <https://doi.org/10.2216/14-024.1>
11. Williamson CJ, Perkins R, Voller M, Yallop ML, Brodie J (2017) The regulation of coralline algal physiology, an in situ study of *Corallina officinalis* (Corallinales, Rhodophyta). *Biogeosciences*. <https://doi.org/10.5194/bg-2017-135>
12. Pardo C, Peña V, Barreiro R, Bárbara I (2015) A molecular and morphological study of *Corallina* sensu lato (Corallinales, Rhodophyta) in the Atlantic Iberian Peninsula. *Cryptogam Algal* 36:31–54. <https://doi.org/10.7872/crya.v36.iss1.2015.31>
13. Nelson WA (2009) Calcified macroalgae—critical to coastal ecosystems and vulnerable to change: a review. *Mar Freshw Res* 60:787–801. <https://doi.org/10.1071/MF08335>
14. van der Heijden LH, Kamenos NA (2015) Reviews and syntheses: calculating the global contribution of coralline algae to total carbon burial. *Biogeosciences* 12:6429–6441. <https://doi.org/10.5194/bg-12-6429-2015>
15. Kim KM, Yang EC, Kim JH, Nelson WA, Yoon HS (2015) Complete mitochondrial genome of a rhodolith, *Sporolithon durum*

- (Sporolithales, Rhodophyta). Mitochondrial DNA 26:155–156. <https://doi.org/10.3109/19401736.2013.819500>
16. Robba L, Russell SJ, Barker GL, Brodie J (2006) Assessing the use of the mitochondrial cox1 marker for use in DNA barcoding of red algae (Rhodophyta). Am J Bot 93:1101–1108. <https://doi.org/10.3732/ajb.93.8.1101>
 17. Hind KR, Saunders GW (2013) A molecular phylogenetic study of the tribe Corallineae (Corallinales, Rhodophyta) with an assessment of genus-level taxonomic features and descriptions of novel genera. J Phycol 49:103–114. <https://doi.org/10.1111/jpy.12019>
 18. Hind KR, Gabrielson PW, Lindstrom SC, Martone PT (2014) Misleading morphologies and the importance of sequencing type specimens for resolving coralline taxonomy (Corallinales, Rhodophyta): *Pachyarthon cretaceum* is *Corallina officinalis*. J Phycol 50:760–764. <https://doi.org/10.1111/jpy.12205>
 19. Corallinales, Rhodophyta (2016) Complete mitochondrial genome of the geniculate calcified red alga, *Corallina officinalis*. Mitochondrial DNA Part B 1:326–327. <https://doi.org/10.1080/23802359.2016.1172048>
 20. Yesson C, Jackson A, Russell S et al (2018) SNPs reveal geographical population structure of *Corallina officinalis* (Corallinales, Rhodophyta). Eur J Phycol. <https://doi.org/10.1080/09670262.2017.1402373>
 21. Glynn WH (2018) Population structure in the red calcifying alga *Corallina officinalis* in the North Atlantic: implications in a time of global climate change. Thesis project for the MRes in Biodiversity, Evolution and Conservation at University College London
 22. Toonen RJ, Hughes S (2001) Increased throughput for fragment analysis on an ABI Prism® 377 automated sequencer using a membrane comb and STRand software. Biotechniques 31:1320–1325
 23. Belkhir K, Borsa P, Chikhi L, Raufaste N, Bonhomme F (2013) GENETIX 4.05, logiciel sous Windows TM pour la génétique des populations. Laboratoire Génome, Populations, Interactions, CNRS UMR 5171, Université de Montpellier II, Montpellier
 24. Raymond M, Rousset F (1995) GENEPOP (version 4.2): population genetics software for exact tests and ecumenicism. J Heredity 86:248–249
 25. Dufresne F, Stift M, Vergilino R, Mable BK (2014) Recent progress and challenges in population genetics of polyploid organisms: an overview of current state-of-the-art molecular and statistical tools. Mol Ecol 23:40–69. <https://doi.org/10.1111/mec.12581>

Table 1 Characterization of nine microsatellite loci in two *Corallina officinalis* populations

Locus name	Primer sequences 5'-3'	Repeat motif	Size range (bp)	Ta (°C)	Iceland		UK		H_o	F_{is}	H_e	H_o	F_{is}
Coroff_1036	F-TCGATACGACACAAACAGC R-ATCGGGCATTATGTTCCAC	(AGC) ₇	235-254	56	2	2	3	3	0.001	1*	0.375	0.143	0.778*
Coroff_1258	F-ATAATTTCCGTGGCCAGAACG R-CCGACACCCATACTCTCC	(GA) ₁₃	105-121	56	4	4	2	2	0.375	0.333	0.492	0.001	1*
Coroff_1640	F-GCCCGGGTATTCATAATGGG R-GCTCCAGTATTTGGCGCTAC	(GA) ₁₅	142-145	56	2	2	3	3	0.000	1*	0.375	0.143	0.812**
Coroff_2347	F-CTCAAGTGAAAGCCGCAAG R-GAGTACTTTGGCCACATGCC	(GTGC) ₇	175-646	56	6	6	5	5	0.687	0.333	0.687	0.143	0.8**
Coroff_2797	F-TAAGAAAGCATCCGTCGAG R-CGCTTCACGCTCTCAGTTG	(CA) ₁₅	194-226	56	4	4	4	4	0.648	0.479*	0.648	0.571	0.238
Coroff_3261	F-CCAGCCTAAAGAAGTTCCGG R-GACTGTATAAATCGACGGCGG	(CCG) ₈	224-246	56	5	5	3	3	0.734	-0.126	0.875	0.571	0.059
Coroff_7707	F-CCCCTGTCACGAACGAAAC R-AAGACTTGACCGAGGACAGC	(TGC) ₁₀	77-97	56	4	4	3	3	0.68	-0.225	0.875	0.857	-0.333
Coroff_8333	F-CGTTTCGATAATGTCCTCCAC R-TCTCTGTATGACCAACGGC	(AG) ₂₀	169-197	56	5	5	5	5	0.422	0.461*	0.422	0.429	0.5*
Coroff_11062	F-CGGTTAGGTAGCTTGGGAG R-TTCGCCCTCAGAAAAATGCG	(ACA) ₁₁	211-239	56	4	4	4	4	0.539	0.364	0.539	0.143	0.769**

Locus name, primer sequence, motif, repeat, size range, PCR annealing temperature (Ta)

A number of alleles, H_e expected heterozygosity; H_o observed heterozygosity; F_{is} inbreeding coefficient

*Significant values at $p < 0.05$ and ** at $p < 0.01$