The physiological effects of daily Cold-water immersion on 5-day tournament performance in international standard youth field-hockey players

Authors: Malte Krueger^{1, 2}; Joseph T Costello³; Mirko Stenzel^{2, 4}; Joachim Mester²; Patrick Wahl^{2, 5}

Institutions:

¹ Institute of Cardiology and Sports Medicine - Department of Preventative and Rehabilitative Sports and Performance Medicine, German Sport University Cologne, Cologne, Germany

² The German Research Centre of Elite Sport, German Sport University Cologne, Cologne, Germany

³ Extreme Environments Laboratory, Department of Sport and Exercise Science, University of Portsmouth, Portsmouth, UK

⁴ German Hockey Federation, Mönchengladbach, Germany

⁵ Institute of Cardiology and Sports Medicine – Department of Molecular and Cellular Sports Medicine, German Sport University Cologne, Cologne, Germany

Corresponding author

Malte Krueger

German Sport University Cologne

Institute of Cardiology and Sports Medicine, Depart. of Preventative and Rehabilitative Sports and Performance Medicine

Am Sportpark Müngersdorf 6, 50933 Cologne, Germany

E-mail: malte.krueger@bayer04.de

Phone: +49-(0)221-4982 6062

Abstract

PURPOSE: This study examined the effects of daily post-exercise cold-water immersion (CWI) on match performance, perceptual recovery, and biomarkers of muscle damage and metabolic load during a 5-day international tournament of elite youth field-hockey players.

METHODS: The entire German under-18 national squad (n=18) were randomly assigned to a daily CWI- (5-min at ~6°C; excluding the head; n=9) or passive recovery (CON; n=9) intervention. Training- and match-performance was assessed using a GPS-tracking system and perceived exertion (RPE). Daily ratings of delayed onset muscle soreness (DOMS), perceived stress and recovery, quality of sleep, heart-rate recovery (HRR) and serum creatinekinase (CK), lactate-dehydrogenase, and urea nitrogen were also recorded. Repeated sprint ability (RSA) and counter-movement jump (CMJ) was carried out on day 1 and 5.

RESULTS: There was no significant between intervention differences in time on pitch, total distance, velocity-zones and accelerometer-bases parameters during match performance (all $p > 0.05$). DOMS ($p < 0.01$), RPE ($p < 0.01$), and CK ($p < 0.01$) were significantly elevated over the course of the tournament; however, no between intervention effects were observed (all $p > 0.05$). Both groups were able to maintain RSA and CMJ (all $p > 0.05$).

CONCLUSION: In conclusion, daily post-exercise CWI did not improve match performance, perceptual recovery, or biomarkers of muscle damage and metabolic load in elite youth field-hockey players.

Keywords: Recovery, DOMS, GPS-tracking, Elite athletes, Biomarkers

51 **Abbreviations:**

52	ACC	Accelerations
53	ANOVA	Analysis of variance
54	BUN	Blood urea nitrogen
55	CK	Creatinekinase
56	CMJ	Counter-movement jump
57	CON	Control
58	CWI	Cold-water immersion
59	DEC	Decelerations
60	DOMS	delayed onset muscle soreness
61	ExE	Explosive efforts
62	GPS	Global positioning system
63	HR _{ex}	Exercise heart rate
64	HRR	Heart rate recovery
65	HSD	High-speed distance
66	LDH	Lactate- dehydrogenase
67	RPE	Rating of perceived exertion
68	RSA	Repeated sprint ability
69	TD	Total distance
70	T _{skin}	Skin temperature
71	VZ	Velocity zone

72

73

74

75

76 **Introduction**

77 In elite international field hockey tournaments, players are required to play and achieve
78 their best performance possible almost daily, with the first 3 matches often scheduled
79 within 4 days (Jennings et al. 2012). Repeating high-intensity exercise with limited time
80 for recovery may lead to accumulated fatigue and consequently, reduced performance
81 capacity (Spencer et al. 2005, Rowsell et al. 2011, Clarkson and Hubal 2002). Reductions
82 in exercise duration and intensity for 1-2 days, following high-intensity exercise as
83 recommended (Cheung et al. 2003), are not always possible due to the tournaments
84 schedule. Therefore, protocols to enhancing recovery or reducing fatigue between
85 matches are an important factor for success and the players' ability to maintain
86 performance throughout the tournament.

87 The use of time-motion analysis and GPS-tracking devices is widely spread to quantify
88 physical activity in team sports (Lythe and Kilding 2011, White and MacFarlane 2013,
89 White and MacFarlane 2015, Cummins et al. 2013). It has been well demonstrated that
90 decrements in total running distance (TD) and high-speed running distance (HSD) occur
91 throughout team-sport tournaments (e.g. soccer and rugby), potentially indicating
92 accumulated fatigue (Rowsell et al. 2011, Johnston et al. 2013). In elite field hockey,
93 findings are inconsistent as some studies reported reductions in TD, HSD and number of
94 sprints (Spencer et al. 2005, Ihsan et al. 2017), while another stated no changes
95 (Jennings et al. 2012). However, it is well established that there is a large variability in
96 physical match performance in elite teams, which is likely attributed to technical and
97 tactical requirements (Gregson et al. 2010). Thus, players may not be required to reach
98 their physiological limits and decrements in certain recovery parameters are not
99 necessarily associated with accumulated fatigue.

Therefore, various surrogate outcome measures are used to monitor fatigue throughout consecutive training and competition, such as counter-movement jumps (CMJ), repeated sprint ability (RSA) or heart rate recovery tests (HRR) (Rowell et al. 2009, Buchheit et al. 2013, Thorpe et al. 2015). Moreover, perturbations of various biomarkers related to exercise-induced muscle damage and inflammation are frequently assessed to quantify the athletes load and recovery status during training and competition (Meyer and Meister 2011). Additionally, the players' internal load can be assessed using psychometric questionnaires related to delayed onset muscle soreness (DOMS), freshness, quality of sleep, fatigue and stress (Buchheit et al. 2013, Hitzschke et al. 2015, Williamson and Hoggart 2005). In order to get a better understanding of any potential impact of an recovery-intervention, it has been recommended to use multiple parameters and assessments to quantify performance, fatigue and recovery (Halsen 2014, Kellmann et al. 2018).

Post-exercise cooling is a common recovery modality for prevention and treatment of exercise-induced fatigue (Poppendieck et al. 2013, Hohenauer et al. 2015). Among the most common methods are brief exposures to cold water (cold-water immersion, CWI, ~ 5-13°C for 5-20 min), extremely cold air (whole-body cryotherapy, WBC, ~ -110°C for 2 to 4 min) or extremely cold vaporized liquid nitrogen (partial- body cryotherapy, PBC, - 110-195°C for 1-3 min) (Bleakley et al. 2012, Costello et al. 2015, Bouzigon et al. 2016, Leeder et al. 2012). Despite its increasing popularity and application in sports, evidence that cryotherapy improves recovery is equivocal (Murray and Cardinale 2015, Poppendieck et al. 2013, Leeder et al. 2012). Several reviews of the literature suggest, that by its primary effects of reducing tissue temperature, peripheral blood flow, and nerve

conductance velocity, cryotherapy attenuates the perception of exercise-induced muscle damage and DOMS (Bleakley et al. 2012, Hohenauer et al. 2015, Ihsan et al. 2016). However, concurrent findings are published regarding the effects of cryotherapy on biomarkers of muscle damage and inflammation (Ascensao et al. 2011, Pournot et al. 2011, Ingram et al. 2009, Krueger et al. 2018). These disparate results might be related to differences in methodology, such as cooling temperature, immersion depth, initial exercise modality and time points of recovery parameter assessment. In addition, it has been suggested that CWI is most effective after prolonged whole body endurance and/or intermittent based exercise (Bleakley et al. 2012, Ihsan et al. 2016). However, most studies focused on single exercise bouts and single applications of CWI in recreational trained participants, thus data of daily CWI applications following repetitive high-intensity training and match performance in international level athletes is extremely limited. To our knowledge, no study has investigated the effects of CWI compared to a control condition on 1) match performance and 2) various recovery parameters during a tournament in elite field hockey to date.

Accordingly, the aim of the present study was to investigate the effectiveness of daily post-exercise CWI on physiological and perceptual markers of fatigue and recovery and physical match-performance during a 5-day tournament in international standard youth field-hockey players. We hypothesised that 1) CWI would not improve performance-related parameters or biomarkers of muscle damage or fatigue, and 2) that CWI would improve perceptual markers associated with fatigue and recovery.

Methods

Participants

The under-18 German national Squad in field hockey participated in the present study. The team won the European Championship in the in the year of the present studies conduction. During the investigated tournament the national squad consisted of 18 healthy male field hockey players (mean $\pm$ SD: age 16.6 ± 0.6 yrs.; height 182.1 ± 5.5 ; mass 73.8 ± 7.8 kg). All participants played field hockey for at least five years with three or more training sessions per week and regular league matches in their respective home clubs. Additionally, they were familiar with different recovery methods like message or CWI. Before the tournament, players were informed about the testing procedures and potential risks and familiarized with the questionnaires used during the investigation. After that, all volunteers and their parents or legal guardians gave their written informed consent to participate. The study, including all procedures was conducted in accordance with the Declaration of Helsinki and approved by the ethical committee of the German Sport University Cologne.

Experimental design

An overview of the experimental design is presented in figure 1.

Briefly, the tournament took place immediately after the last week of the regular season with players arriving from their respective home clubs in the morning of day 1. After anthropometric measurements and baseline assessment of questionnaires and biomarkers, the team completed a 15-min warm-up, followed by performance testing of CMJ and 6 x 30m RSA. Afterwards, the players were paired by the coaches according to

their playing position, level of experience and anthropometrics. Randomly (coin toss) one player of each pair was assigned to the intervention (CWI) and control (CON) group, respectively. In the afternoon of day 1 a friendly match against a local club was carried out, consisting of 3 x 20 min. On day 2 tactical coaching and a 2 h – training session was conducted. On days 3 – 5, the team participated in an international tournament with one match per day. In the afternoon of day 5 assessment of CMJ and RSA performance was repeated. Prior to the experiment, the coach had requested that all players would play the same amount of time on the pitch and cover an similar distance. Interchanges were conducted accordingly to achieve this during the matches.

*** Figure 1 near here****

Performance tests (RSA, CMJ)

CMJ-height was measured using the Optojump photoelectric cell system (Microgate, Bolzano, Italy) as previously described (Glatthorn et al. 2011). Players were advised to place their hands at the hips to avoid additional arm swing. The best of three maximal effort jumps was taken for statistical analysis. The RSA-test consisted of 6 maximal 30m efforts, starting every 25 sec as recommended for field hockey players and described elsewhere (Tanner and Gore 2013). Sprint time was assessed using a dual beam sports timing system (Swiftperformance, Wacol, Australia). Gates were placed at the start, 10m, 20m and 30m. Total sprint time was calculated using the sum of all six sprints (total sprint time). The test-retest coefficient of variation for youth male team-sport athletes tested in our laboratory is 2.8% for CMJ and 1.1% for RSA.

Physical Performance Monitoring during match and training

189 The physical activity of each match and training session was monitored using a portable
190 positional tracking device (10hz GPS and GLONASS) with integrated 3D-Accelerometer,
191 3D-Gyroscope and 3D-Magnetometer using a sampling rate of 100hz, respectively
192 (OptimEye S5, Catapult sports, Melbourne, Australia). Each player's individually assigned
193 device was worn at the upper back between the left and right scapulae in a custom-made
194 vest (Catapult sports). All devices were turned on at least 30 min before exercise for
195 satellite connection. During data collection the average number of satellites and horizontal
196 dilution of precision per unit (mean $\pm$ SD) were 12.0 ± 0.0 and 0.61 ± 0.01 , indicating an
197 almost ideal signal quality (Malone et al. 2017). During matches, the players' rotations
198 were precisely documented. After recording, data was downloaded, analysed and
199 reported using the appending software (OpenField v1.17.0). Goalkeepers were excluded
200 from statistical analysis. Bench time of the field players was excluded and values
201 calculated for time on pitch only, as described previously and recommended for time-
202 depended variables in field hockey (White and MacFarlane 2013, Ihsan et al. 2017).
203 Thresholds for six velocity zones (VZ) were matched with previously published
204 investigations in field hockey (VZ1, $< 6 \text{ km}\cdot\text{h}^{-1}$; VZ2, $6 - 11 \text{ km}\cdot\text{h}^{-1}$; VZ3, $11 - 15 \text{ km}\cdot\text{h}^{-1}$;
205 VZ4, $15 - 19 \text{ km}\cdot\text{h}^{-1}$; VZ5, $19 - 23 \text{ km}\cdot\text{h}^{-1}$; VZ6, $> 23 \text{ km}\cdot\text{h}^{-1}$) (White and MacFarlane
206 2015, White and MacFarlane 2013, Lythe and Kilding 2011). The time spend in each VZ
207 was calculated as percentage of time on pitch. HSD was measured for speeds above 19
208 $\text{km}\cdot\text{h}^{-1}$. The software's inertial movement analysis was used to detect accelerations
209 (ACC) and decelerations (DEC) above $2.5 \text{ m}\cdot\text{s}^{-2}$ and explosive efforts (ExE), defined as
210 the sum of accelerations, decelerations, changes of direction to the left and right above
211 $3.5 \text{ m}\cdot\text{s}^{-2}$. TD, HSD, ACC, DEC and ExE were normalised to one minute for comparison.

30 min after the end of each exercise, players were asked to rate their perceived exertion of the previous session (RPE_{session}) on a 1 – 10 Borg scale (Foster et al. 2001).

Cold-water immersion and passive recovery

15 minutes after each match or training on days 1 – 4 the CWI-group fully immersed themselves into an ice-water barrel for 5 minutes, with only the head out of the water. The barrels were prepared with crushed ice to a water temperature between 5 and 8°C ($6.4 \pm 0.8^\circ\text{C}$). During immersion, the temperature was frequently controlled and ice added if necessary. Players were instructed to slowly move their arms and legs to avoid the formation of warmer water layers around the body. Similar protocols have been used extensively elsewhere (Ingram et al. 2009, Rowsell et al. 2009, Costello et al. 2012). During the same time, the CON-group remained passive and seated at room temperature. Afterwards, the whole team continued with the schedule of the tournament together.

Mean Skin temperature

Skin temperature was assessed by means of a thermoimaging camera (Testo 880, Testo GmbH, Vienna, Austria) as described elsewhere (Moreira et al. 2017). Photos were taken of the trunk, thigh and lower arm before and after CWI and analysed using the appending software (IRsoft Version 3.1 SP3, Testo GmbH, Vienna, Austria). Mean skin temperature (T_{skin}) was calculated using the Burton (Burton 1935) equation: $T_{\text{skin}} = 0.5 T_{\text{trunk}} + 0.36 T_{\text{thigh}} + 0.14 T_{\text{arm}}$.

Biomarkers

232 Every morning 300 μL of capillary blood was collected from the earlobe of each athlete,
233 transferred in a heparinized centrifuge tube, centrifuged for 2 minutes and immediately
234 analysed for serum levels in Creatinkinase (CK), Lactate-Dehydrogenase (LDH) and
235 Blood Urea Nitrogen (BUN) using a mobile Point-of-care device (Spotchem EZ SP-4430,
236 Axonlab, Baden-Daettwil, Germany). Serum levels were determined by optical
237 measurement of reflection intensity (at 550 or 610 nm) for a reagent colour reaction on
238 three single reagent strips. The reagent strip is composed of a multi-layered test field
239 containing reagents necessary to generate a colour that is quantified by reflectance
240 spectrophotometry. The precision (CV %) of consecutive replicate measurements was
241 3.8% for CK ($94.7 \text{ U}\cdot\text{L}^{-1}$), 3.1 % for BUN ($6.1 \text{ mmol}\cdot\text{L}^{-1}$) and 3.7% for LDH ($322.9 \text{ U}\cdot\text{L}^{-1}$).

242 ***Psychological questionnaires, quality of sleep and Pain scale***

243 Every morning before breakfast, the players answered a questionnaire assessing their
244 acute state of recovery and stress. To avoid misunderstanding and translational problems
245 a questionnaire in German language was chosen, that has already been validated in field
246 hockey (Hitzschke et al. 2015). The items of the “Short Recovery and Stress Scale for
247 Sport” were rated on a scale from 0 (“not at all”) to 6 (“absolutely”). Furthermore, players
248 were asked to rate the quality of last nights’ sleep on a scale from 1 (“very good”) to 6
249 (“very bad”).

250 Moreover, DOMS was measured every morning and evening, using a 100 mm visual
251 analog scale, ranging from 0 (“no soreness”) to 100 (“severe soreness”) (Williamson and
252 Hoggart 2005). The players rated their individual DOMS after slowly sitting down and
253 standing up from a chair (knee angle approximately 90°).

Heart rate recovery test

A heart rate recovery test was included into the warm-up session before each match or training. On an indoor pitch (length: 50m) the entire team ran for 5 min at a constant speed of 9 km/h, controlled by a digital metronome, followed by sitting in a relaxed position for another 5 min. Heart rate was recorded continuously using a mobile team system (Acentas GmbH, Hörgertshausen, Germany). Exercise heart rate (HR_{ex}), defined as the average heart rate during the last 30 seconds and heart rate recovery (HRR), defined as the difference between HR_{ex} and heart rate after one minute of recovery were calculated as described elsewhere (Buchheit 2014, Borresen and Lambert 2007). HR_{ex} was expressed as percentage of the players' individual maximal HR, HRR was expressed as percentage of HR_{ex} .

Statistical analysis

All statistical tests were carried out using the Statistica software package for Windows® (version 13.0, StatSoft Inc., Tulsa, OK, U.S.A). The distribution of data was assessed using descriptive methods (skewness, outliers, and distribution plots) and inferential statistics (Shapiro–Wilk test). Normal distribution was observed for all parameters except CK. Normally distributed data is presented as mean values $\pm$ standard deviations (SD), and mean differences (MD) with 95% confidence intervals for performance tests and distribution of velocity zones. Data deviating from normal distribution is presented as median and inner-quartile range (Q_{25} and Q_{75}). A two way (treatment [CWI, CON] * time [Days 1-5]) repeated-measures analysis of variance (ANOVA) was applied to compare all biomarkers (CK, LDH, BUN), Questionnaires (perceived recovery, perceived stress,

quality of sleep, DOMS), heart rate recovery tests (HR_{ex} , HRR), performance tests (RSA, CMJ) and tracking data (time on pitch, total distance, normalised distance, HSD, ACC, DEC and ExE). If main effects for time, intervention or interaction were identified, Bonferroni post-hoc analysis was applied where appropriate. Student paired t-tests were used for analysis of percentage time spend in each velocity zone during tournament matches (days 3-5) as well as the effect of CWI on T_{skin} (pre vs. post). Statistical significance was accepted at $p < 0.05$.

Results

Tracking data

All GPS and accelerometer-based parameters are presented in Table 1 and 2 (means $\pm$ SD). All parameters (time on pitch, TD, normalised distance, normalised HSD, normalised DEC) except normalised ACC ($p = 0.21$) and normalised ExE ($p = 0.62$) significantly increased from day 1 to day 5 ($p < 0.01$). However, this is most likely related to the extended training duration on day 2. There were no significant intervention or interaction effects (all $p > 0.05$) (Table 1). During tournament matches there were also no differences (all $p > 0.05$) between CWI and CON in time spent in each velocity zone (Table 2).

*** Table 1 and 2 near here****

Performance tests

There were no time, interaction or intervention effects (all $p > 0.05$) in RSA total sprint time (mean $\pm$ SD; mean difference [95% CI]) from day 1 (CWI: 26.23 ± 1.06 s, CON: 26.05

297 ± 0.69 s; MD: 0.18 [-0.78 – 1.14]) to day 5 (CWI: 26.37 $\pm$ 1.06 s, CON: 26.34 $\pm$ 0.73 s;
298 MD: 0.04 [-0.86 – 0.92]). The same results (all $p > 0.05$) were observed in CMJ with
299 similarly no differences from day 1 (CON: 42.06 $\pm$ 3.48 cm, CWI: 41.37 $\pm$ 5.31 cm; MD:
300 0.69 [-3.71 – 5.09]) to day 5 (CON: 41.82 $\pm$ 3.64 cm, CWI: 41.59 $\pm$ 4.61 cm; MD: 0.23 [-
301 3.84 – 4.30]), respectively.

302 ***Heart rate recovery / Questionnaires***

303 The results for the daily measurements of the heart rate recovery and questionnaires are
304 detailed in Table 3. Throughout the course of the tournament RPE_{session} and perceived
305 stress increased, while perceived recovery, HR_{ex} and HRR decreased (all $p < 0.05$).
306 However, no significant intervention or interaction effects were observed in heart rate
307 recovery- or in questionnaire data (all $p > 0.05$). DOMS (Figure 2) increased over time (p
308 < 0.01), but no intervention ($p = 0.09$) and interaction ($p = 0.41$) effect was evident.

309 *** Table 3 and Figure 2 near here****

310 ***Biomarkers***

311 Serum biomarkers data for CK, LDH and BUN is presented in Figure 3. CK increased
312 over time ($p < 0.01$), but no other significant differences were observed (all $p > 0.05$).

313 *** Figure 3 near here****

314 ***CWI - Mean skin temperature***

315 T_{skin} was significantly reduced ($p < 0.01$; $d = 11.5$) from 29.8 $\pm$ 1.3 °C to 14.9 $\pm$ 1.3 °C
316 immediately after each application of CWI.

317

318 **Discussion**

319 The present study is the first to delineate the effects of daily post exercise CWI on
320 performance capacity, biomarkers of muscle damage and metabolic load, fatigue, and
321 recovery during a 5-day junior elite field hockey tournament. The main findings of the
322 present study are: (1) in support of our hypothesis, daily CWI did not alter performance
323 parameters during consecutive matches, (2) regardless of recovery intervention, elite
324 youth field hockey players were able to maintain match-to-match performance, (3)
325 exercise-induced alterations in biomarkers were not attenuated by CWI, and (4) contrary
326 to our second hypothesis, CWI did not improve perceptual recovery. Collectively, these
327 data suggest that daily post exercise CWI has no beneficial effects during a 5-day
328 tournament and therefore its usage is not recommended in international standard male
329 junior field hockey players

330 It has previously been reported that CWI might be able to reduce fatigue associated
331 decrements in physical performance (i.e. TD and HSD) during consecutive tournament
332 soccer matches in junior males (Rowse et al. 2011). The findings of the current study
333 suggest that these benefits may not be extended to field hockey (Table 1 and 2), as elite
334 youth athletes were able to maintain their physical match performance even without the
335 use of CWI. These contrasting findings might be explained by the large differences in
336 running characteristics observed between both sports. Total duration is less in field
337 hockey (~ 30 min) and unlimited interchanges allow individual recovery breaks and higher
338 intensities on the pitch (Spencer et al. 2005). However, the present data suggests that

the repeated high intensity exercise throughout the tournament did not lead to reduced match running performance or exercise intensity. Distribution of percentage durations per velocity zones was constant over the course of the tournament and significant time effects in time on pitch, TD and normalised distance, HSD and DEC were caused by day 2 training session. These results corroborate the findings of Jennings et al. (Jennings et al. 2012) and Ihsan et al (Ihsan et al. 2017), who also reported that elite hockey players were able to maintain exercise intensity when playing six matches in nine days. Contrary, Spencer and colleagues (Spencer et al. 2005) described a decrease in jogging and increase in walking duration in three subsequent field-hockey matches within four days. It must be acknowledged that match performance variables may not reflect the true status of recovery, as tactical requirements may result in high between match variability (Gregson et al. 2010). Regardless, it appears likely that young elite field hockey players are able to maintain their physical performance throughout consecutive matches.

This is further supported by the results of the jump and repeated sprint performance outcomes (CMJ and RSA). These tests have previously been used to monitor fatigue and recovery state in team sports, as alterations are associated with training and match performance (Thorpe et al. 2015, Buchheit et al. 2013, Claudino et al. 2017, Ingram et al. 2009). In contrast to observations in other team sport tournaments (Rowse et al. 2009, Montgomery et al. 2008), performance in CMJ remained consistent from day 1 to day 5 and RSA was only reduced by ~1%, with no differences between CWI and CON-group. These results indicate that the 5-day tournament did not induce accumulated neuromuscular fatigue and CWI had no impact on athletic performance.

Significant reductions in HR_{ex} and HRR were observed in the present study in the CWI- and CON-groups, with the lowest values recorded on day 3 and 4. This is counterintuitive, as lower HR_{ex} and HRR (when expressed as %HR_{ex}) is typically associated with improved fitness and reduced fatigue (Daanen et al. 2012, Buchheit et al. 2013). Comparable data have been explained as demonstrating symptoms of overreaching (Borresen and Lambert 2007); however, this is very unlikely in the present study as physical performance was maintained throughout the tournament. Further, these international standard players were highly trained prior to the start of the tournament. CWI has been proposed to increase parasympathetic activity along with cardiac output (Ihsan et al. 2016), consequently lower HR_{ex} and HRR could have been expected in the CWI-group. However, CWI did not induce changes in exercise-induced alterations of heart-rate derived variables compared to CON.

Post-exercise ratings of perceived exertion (RPE_{session}) increased from day 1 to 4 in both groups and slightly dropped on day 5 (Table 3). As performance values and exercise intensity were on the same level throughout tournament matches, this increase might be an indicator for increasing perceptual fatigue as shown during a 5-day soccer tournament as well (Rowell et al. 2009). DOMS also accumulated throughout the course of the tournament. Days 3-5 showed significantly elevated levels compared to baseline, however, there were no intervention or interaction effects in this field based analysis (Figure 2). Even though some studies (Pournot et al. 2011, Corbett et al. 2012) reported no effects of CWI on DOMS as well, several empirical laboratory and field based studies (Ascensao et al. 2011, Bailey et al. 2007, Rowell et al. 2009, Ingram et al. 2009) and reviews (Ihsan et al. 2016, Hohenauer et al. 2015, Bleakley et al. 2012, Leeder et al.

2012) suggest that post exercise CWI is effective in reducing DOMS. These findings may be explained by the training status of the international level athletes, which one would expect to result in relatively limited DOMS observed in the current study. Furthermore, DOMS typically peaks ~48 hrs after exercise (Goodall and Howatson 2008), however the team was required to daily, thus the time course for peak DOMS of a single exercise is difficult to identify. Moreover, as exercise itself has been shown to reduce DOMS (Corbett et al. 2012), it is plausible that the exercise may have led to an attenuation of DOMS in the CON- group. Perceived stress and recovery remained similar throughout the tournament and only increased from day 4 to day 5. As the final match decided about winning the tournament, these time effects are most likely related to this. The exercise itself and the recovery modality had no influence on the players' ratings of perceived stress, recovery, or quality of sleep (Table 3).

Findings regarding the effects of CWI on exercise- induced alterations of biomarkers associated with muscle damage and inflammation are contradictory in the literature [for review see (Leeder et al. 2012, Bleakley et al. 2012, Hohenauer et al. 2015, Ihsan et al. 2016)]. For example, several studies have demonstrated decreased levels of CK, Myoglobin or C-reactive Protein compared to thermoneutral or contrast water immersion and passive control (Ascensao et al. 2011, Pournot et al. 2011, Bailey et al. 2007) while others reported no change (Ingram et al. 2009, Pointon et al. 2012, Corbett et al. 2012). In line with the current investigation Rowsell et al. (Rowsell et al. 2009) found no effect on CK-clearance following the use of CWI after 5-day soccer tournament. These finding corroborates the current results as CK increased from day 1 to 2 and remained elevated throughout the course of the tournament, indicating high muscular load in both groups;

yet, CWI did not ameliorate the elevation in CK. However, no changes in LDH or BUN were observed (Figure 3). Similar differences in CK and LDH responses to exercise have been reported before and are potentially attributed to the different structural areas where they are sequestered within the muscle sarcomere (Pournot et al. 2011). Nevertheless, it could be argued that exercise intensity over the course of the five days was not severe enough to induce a higher protein turnover rate, which would have been indicated by elevated levels in BUN (Meyer and Meister 2011).

The present study has limitations that warrant mention. Firstly, we acknowledge the small sample size of n=18. However, the entire national squad participated in the study; therefore, this sample represent the largest number of international level athletes we were able to recruit. Secondly, as indicated by elevated levels in CK, DOMS and stress at baseline, the athletes did not commence the tournament in a completely 'rested' state. This is not surprising and reflects the intense schedule of elite junior team sport athletes that participate in tournaments or international competitions alongside, or immediately after, their regular season. Thirdly, performance data (i.e. CMJ and RSA) was assessed after the athletes arrived on site (circa 10 am) on day 1 and on day 5 three hours following the final match of the tournament (circa 4 pm). We acknowledge that the data may be reflective of this; however, again this represents the challenge of conducting applied research on international athletes and working within their time schedules. However, both the CWI and the control groups were randomised and we do not believe this invalidates our findings regarding the effectiveness, or lack thereof, of the CWI intervention. Finally, we acknowledge that additional follow up data after the tournament (i.e. day 6 and 7)

would have added additional insights, but this was not feasible as the athletes returned home, all across the country, after the tournament.

Conclusion

Despite elevated levels of CK, RPE and DOMS, several parameters (i.e. LDH, BUN, perceived recovery and stress, quality of sleep) were not altered over the course of the 5-day tournament. This suggests that there was an absence of severe cumulative fatigue and/or the international standard athletes were sufficiently capable of completing and recovering from the intensive training load, as indicated by maintained physical performance. The results of the present study also indicate that CWI had no beneficial effects on physical performance, perceptual wellbeing, or biomarkers associated with muscle damage compared to passive recovery. In conclusion, the application of daily CWI appears to be superfluous and we do not recommend its use in elite junior field-hockey players during a 5-day tournament.

Acknowledgements

The authors would like to thank all field-hockey players of the under- 18 German national Squad who participated in this study, and all trainers and staff members for their help and support.

Disclosure statement

The authors confirm there are no conflicts of interest.

Funding

This study received no funding from any grant or organization.

FIGURE CAPTIONS

Fig. 1 Schematic presentation of the study design. CWI: Cold-water immersion; RSA: Repeated sprint ability; CMJ: Counter movement jump; DOMS: rating of delayed onset muscle soreness

Fig. 2 Perceived ratings (mean $\pm$ SD) of delayed onset muscle soreness (DOMS) in the morning (pre) and evening (post) of each day (Days 1-5). * $P < 0.05$ time effects compared to baseline (day 1 pre), for both interventions (cold-water immersion [CWI] and control [CON]) combined

Fig. 3 Serum concentrations of Creatinkinase (CK) (median $\pm$ inner quartile range), Lactate Dehydrogenase (LDH) and Blood Urea Nitrogen (BUN) (mean $\pm$ SD) in the

morning of each day (Days 1-5). * $P < 0.05$ time effects compared to baseline (day 1), for both interventions (cold-water immersion [CWI] and control [CON]) combined

References

- Ascensao A, Leite M, Rebelo AN, Magalhaes S, Magalhaes J (2011) Effects of cold water immersion on the recovery of physical performance and muscle damage following a one-off soccer match. *Journal of sports sciences* 29(3):217–225. doi: 10.1080/02640414.2010.526132
- Bailey DM, Erith SJ, Griffin PJ, Dowson A, Brewer DS, Gant N, Williams C (2007) Influence of cold-water immersion on indices of muscle damage following prolonged intermittent shuttle running. *Journal of sports sciences* 25(11):1163–1170. doi: 10.1080/02640410600982659
- Bleakley C, McDonough S, Gardner E, Baxter GD, Hopkins JT, Davison GW (2012) Cold-water immersion (cryotherapy) for preventing and treating muscle soreness after exercise. *The Cochrane database of systematic reviews*(2):CD008262. doi: 10.1002/14651858.CD008262.pub2
- Borresen J, Lambert MI (2007) Changes in heart rate recovery in response to acute changes in training load. *European journal of applied physiology* 101(4):503–511. doi: 10.1007/s00421-007-0516-6
- Bouzigon R, Grappe F, Ravier G, Dugue B (2016) Whole- and partial-body cryostimulation/cryotherapy. Current technologies and practical applications. *Journal of thermal biology* 61:67–81. doi: 10.1016/j.jtherbio.2016.08.009
- Buchheit M (2014) Monitoring training status with HR measures: do all roads lead to Rome? *Frontiers in physiology* 5:73. doi: 10.3389/fphys.2014.00073
- Buchheit M, Racinais S, Bilsborough JC, Bourdon PC, Voss SC, Hocking J, Cordy J, Mendez-Villanueva A, Coutts AJ (2013) Monitoring fitness, fatigue and running performance during a pre-season training camp in elite football players. *Journal of science and medicine in sport* 16(6):550–555. doi: 10.1016/j.jsams.2012.12.003

507 Burton AC (1935) Human calorimetry II: the average temperature of the tissues of the body. *Journal of*
 508 *Nutrition* 9(3):261–280
 509 Cheung K, Hume P, Maxwell L (2003) Delayed onset muscle soreness. Treatment strategies and
 510 performance factors. *Sports medicine (Auckland, N.Z.)* 33(2):145–164
 511 Clarkson PM, Hubal MJ (2002) Exercise-induced muscle damage in humans. *American journal of physical*
 512 *medicine & rehabilitation* 81(11 Suppl):S52-69. doi: 10.1097/01.PHM.0000029772.45258.43
 513 Claudino JG, Cronin J, Mezêncio B, McMaster DT, McGuigan M, Tricoli V, Amadio AC, Serrão JC (2017)
 514 The countermovement jump to monitor neuromuscular status. A meta-analysis. *Journal of science*
 515 *and medicine in sport* 20(4):397–402. doi: 10.1016/j.jsams.2016.08.011
 516 Corbett J, Barwood MJ, Lunt HC, Milner A, Tipton MJ (2012) Water immersion as a recovery aid from
 517 intermittent shuttle running exercise. *European journal of sport science* 12(6):509–514. doi:
 518 10.1080/17461391.2011.570380
 519 Costello JT, Baker PRA, Minett GM, Bieuzen F, Stewart IB, Bleakley C (2015) Whole-body cryotherapy
 520 (extreme cold air exposure) for preventing and treating muscle soreness after exercise in adults.
 521 *The Cochrane database of systematic reviews*(9):CD010789. doi:
 522 10.1002/14651858.CD010789.pub2
 523 Costello JT, Culligan K, Selfe J, Donnelly AE (2012) Muscle, skin and core temperature after -110°C cold
 524 air and 8°C water treatment. *PloS one* 7(11):e48190. doi: 10.1371/journal.pone.0048190
 525 Cummins C, Orr R, O'Connor H, West C (2013) Global positioning systems (GPS) and microtechnology
 526 sensors in team sports. A systematic review. *Sports medicine (Auckland, N.Z.)* 43(10):1025–1042.
 527 doi: 10.1007/s40279-013-0069-2
 528 Daanen HAM, Lamberts RP, Kallen VL, Jin A, van Meeteren NLU (2012) A systematic review on heart-rate
 529 recovery to monitor changes in training status in athletes. *International journal of sports physiology*
 530 *and performance* 7(3):251–260
 531 Foster C, FLORHAUG JA, FRANKLIN J, GOTTSCHALL L, HROVATIN LA, PARKER S, DOLESHAL P, DODGE C
 532 (2001) A New Approach to Monitoring Exercise Training. *Journal of strength and conditioning*
 533 *research* 15(1):109–115. doi: 10.1519/00124278-200102000-00019
 534 Glatthorn JF, Gouge S, Nussbaumer S, Stauffacher S, Impellizzeri FM, Maffiuletti NA (2011) Validity and
 535 reliability of Optojump photoelectric cells for estimating vertical jump height. *Journal of strength*
 536 *and conditioning research* 25(2):556–560. doi: 10.1519/JSC.0b013e3181ccb18d
 537 Goodall S, Howatson G (2008) The effects of multiple cold water immersions on indices of muscle
 538 damage. *Journal of sports science & medicine* 7(2):235–241
 539 Gregson W, Drust B, Atkinson G, Salvo VD (2010) Match-to-match variability of high-speed activities in
 540 premier league soccer. *International journal of sports medicine* 31(4):237–242. doi: 10.1055/s-
 541 0030-1247546
 542 Halson SL (2014) Monitoring training load to understand fatigue in athletes. *Sports medicine (Auckland,*
 543 *N.Z.)* 44 Suppl 2:S139-47. doi: 10.1007/s40279-014-0253-z
 544 Hitzschke B, Kölling S, Ferrauti A, Meyer T, Pfeiffer M, Kellmann M (2015) Entwicklung der Kurzskaala zur
 545 Erfassung von Erholung und Beanspruchung im Sport (KEB). *Zeitschrift für Sportpsychologie*
 546 22(4):146–162. doi: 10.1026/1612-5010/a000150

547 Hohenauer E, Taeymans J, Baeyens J-P, Clarys P, Clijsen R (2015) The Effect of Post-Exercise Cryotherapy
 548 on Recovery Characteristics. A Systematic Review and Meta-Analysis. *PloS one* 10(9):e0139028. doi:
 549 10.1371/journal.pone.0139028
 550 Ihsan M, Tan F, Sahrom S, Choo HC, Chia M, Aziz AR (2017) Pre-game perceived wellness highly
 551 associates with match running performances during an international field hockey tournament.
 552 *European journal of sport science* 17(5):593–602. doi: 10.1080/17461391.2017.1301559
 553 Ihsan M, Watson G, Abbiss CR (2016) What are the Physiological Mechanisms for Post-Exercise Cold
 554 Water Immersion in the Recovery from Prolonged Endurance and Intermittent Exercise? *Sports*
 555 *medicine (Auckland, N.Z.)* 46(8):1095–1109. doi: 10.1007/s40279-016-0483-3
 556 Impellizzeri FM, Rampinini E, Castagna C, Bishop D, Ferrari Bravo D, Tibaudi A, Wisloff U (2008) Validity
 557 of a repeated-sprint test for football. *International journal of sports medicine* 29(11):899–905. doi:
 558 10.1055/s-2008-1038491
 559 Ingram J, Dawson B, Goodman C, Wallman K, Beilby J (2009) Effect of water immersion methods on
 560 post-exercise recovery from simulated team sport exercise. *Journal of science and medicine in sport*
 561 12(3):417–421. doi: 10.1016/j.jsams.2007.12.011
 562 Jennings D, Cormack SJ, Coutts AJ, Aughey RJ (2012) GPS analysis of an international field hockey
 563 tournament. *International journal of sports physiology and performance* 7(3):224–231
 564 Johnston RD, Gabbett TJ, Jenkins DG (2013) Influence of an intensified competition on fatigue and match
 565 performance in junior rugby league players. *Journal of science and medicine in sport* 16(5):460–465.
 566 doi: 10.1016/j.jsams.2012.10.009
 567 Kellmann M, Bertollo M, Bosquet L, Brink M, Coutts AJ, Duffield R, Erlacher D, Halson SL, Hecksteden A,
 568 Heidari J, Kallus KW, Meeusen R, Mujika I, Robazza C, Skorski S, Venter R, Beckmann J (2018)
 569 Recovery and Performance in Sport. Consensus Statement. *International journal of sports*
 570 *physiology and performance* 13(2):240–245. doi: 10.1123/ijsp.2017-0759
 571 Krueger M, Costello JT, Achtzehn S, Dittmar K-H, Mester J (2018) Whole-body cryotherapy (-110 °C)
 572 following high-intensity intermittent exercise does not alter hormonal, inflammatory or muscle
 573 damage biomarkers in trained males. *Cytokine*. doi: 10.1016/j.cyto.2018.07.018
 574 Leeder J, Gissane C, van Someren K, Gregson W, Howatson G (2012) Cold water immersion and recovery
 575 from strenuous exercise. A meta-analysis. *British journal of sports medicine* 46(4):233–240. doi:
 576 10.1136/bjsports-2011-090061
 577 Lythe J, Kilding AE (2011) Physical demands and physiological responses during elite field hockey.
 578 *International journal of sports medicine* 32(7):523–528. doi: 10.1055/s-0031-1273710
 579 Malone JJ, Lovell R, Varley MC, Coutts AJ (2017) Unpacking the Black Box. Applications and
 580 Considerations for Using GPS Devices in Sport. *International journal of sports physiology and*
 581 *performance* 12(Suppl 2):S218–S226. doi: 10.1123/ijsp.2016-0236
 582 Meyer T, Meister S (2011) Routine blood parameters in elite soccer players. *International journal of*
 583 *sports medicine* 32(11):875–881. doi: 10.1055/s-0031-1280776
 584 Montgomery PG, Pyne DB, Hopkins WG, Dorman JC, Cook K, Minahan CL (2008) The effect of recovery
 585 strategies on physical performance and cumulative fatigue in competitive basketball. *Journal of*
 586 *sports sciences* 26(11):1135–1145. doi: 10.1080/02640410802104912
 587 Moreira DG, Costello JT, Brito CJ, Adamczyk JG, Ammer K, Bach AJE, Costa CMA, Eglin C, Fernandes AA,
 588 Fernández-Cuevas I, Ferreira JJA, Formenti D, Fournet D, Havenith G, Howell K, Jung A, Kenny GP,

- Kolosovas-Machuca ES, Maley MJ, Merla A, Pascoe DD, Priego Quesada JI, Schwartz RG, Seixas ARD, Selfe J, Vainer BG, Sillero-Quintana M (2017) Thermographic imaging in sports and exercise medicine. A Delphi study and consensus statement on the measurement of human skin temperature. *Journal of thermal biology* 69:155–162. doi: 10.1016/j.jtherbio.2017.07.006
- Murray A, Cardinale M (2015) Cold applications for recovery in adolescent athletes. A systematic review and meta analysis. *Extreme physiology & medicine* 4:17. doi: 10.1186/s13728-015-0035-8
- Pointon M, Duffield R, Cannon J, Marino FE (2012) Cold water immersion recovery following intermittent-sprint exercise in the heat. *European journal of applied physiology* 112(7):2483–2494. doi: 10.1007/s00421-011-2218-3
- Poppendieck W, Faude O, Wegmann M, Meyer T (2013) Cooling and performance recovery of trained athletes: a meta-analytical review. *International journal of sports physiology and performance* 8(3):227–242
- Pournot H, Bieuzen F, Duffield R, Lepretre P-M, Cozzolino C, Hausswirth C (2011) Short term effects of various water immersions on recovery from exhaustive intermittent exercise. *European journal of applied physiology* 111(7):1287–1295. doi: 10.1007/s00421-010-1754-6
- Rowell GJ, Coutts AJ, Reaburn P, Hill-Haas S (2009) Effects of cold-water immersion on physical performance between successive matches in high-performance junior male soccer players. *Journal of sports sciences* 27(6):565–573. doi: 10.1080/02640410802603855
- Rowell GJ, Coutts AJ, Reaburn P, Hill-Haas S (2011) Effect of post-match cold-water immersion on subsequent match running performance in junior soccer players during tournament play. *Journal of sports sciences* 29(1):1–6. doi: 10.1080/02640414.2010.512640
- Spencer M, Rechichi C, Lawrence S, Dawson B, Bishop D, Goodman C (2005) Time-motion analysis of elite field hockey during several games in succession. A tournament scenario. *Journal of science and medicine in sport* 8(4):382–391
- Tanner RK, Gore CJ (2013) *Physiological tests for elite athletes*, 2. ed. Human Kinetics, Champaign, Ill.
- Thorpe RT, Strudwick AJ, Buchheit M, Atkinson G, Drust B, Gregson W (2015) Monitoring Fatigue During the In-Season Competitive Phase in Elite Soccer Players. *International journal of sports physiology and performance* 10(8):958–964. doi: 10.1123/ijsspp.2015-0004
- White AD, MacFarlane N (2013) Time-on-pitch or full-game GPS analysis procedures for elite field hockey? *International journal of sports physiology and performance* 8(5):549–555
- White AD, MacFarlane NG (2015) Analysis of international competition and training in men's field hockey by global positioning system and inertial sensor technology. *Journal of strength and conditioning research* 29(1):137–143. doi: 10.1519/JSC.0000000000000600
- Williamson A, Hoggart B (2005) Pain. A review of three commonly used pain rating scales. *Journal of clinical nursing* 14(7):798–804. doi: 10.1111/j.1365-2702.2005.01121.x

Table 1 Comparison of external load parameters (mean $\pm$ SD) between recovery modalities cold-water immersion (CWI) and control (CON)

Variable	Intervention	Day 1 – FM	Day 2 – TR	Day 3 – TM 1	Day 4 – TM 2	Day 5 – TM 3	p-value		
							[time]	[int.]	[time*int.]
Time on pitch [min]	CWI	46.6 $\pm$ 10.4	120.0 $\pm$ 0.0	53.5 $\pm$ 6.0	46.5 $\pm$ 11.1	51.0 $\pm$ 5.4	< 0.01	0.87	0.61
	CON	42.3 $\pm$ 5.4	120.0 $\pm$ 0.0	51.5 $\pm$ 6.4	47.6 $\pm$ 8.7	54.2 $\pm$ 8.9			
Total distance [m]	CWI	5487 $\pm$ 1013	6883 $\pm$ 690	6193 $\pm$ 800	5614 $\pm$ 1408	5765 $\pm$ 614	< 0.01	0.81	0.70
	CON	5146 $\pm$ 697	6990 $\pm$ 394	5940 $\pm$ 956	5337 $\pm$ 976	6185 $\pm$ 1082			
Normalised distance [m·min ⁻¹]	CWI	119.3 $\pm$ 11.0	57.4 $\pm$ 5.7	115.8 $\pm$ 7.2	120.5 $\pm$ 6.2	113.3 $\pm$ 7.0	< 0.01	0.82	0.24
	CON	121.9 $\pm$ 7.6	58.2 $\pm$ 3.3	115.3 $\pm$ 12.8	112.8 $\pm$ 11.8	114.0 $\pm$ 10.6			
Normalised HSD [n·min ⁻¹]	CWI	8.86 $\pm$ 4.63	2.16 $\pm$ 1.02	7.61 $\pm$ 3.81	10.02 $\pm$ 4.63	8.40 $\pm$ 2.60	< 0.01	0.65	0.83
	CON	10.32 $\pm$ 2.14	2.38 $\pm$ 0.74	8.56 $\pm$ 2.27	10.43 $\pm$ 3.14	7.60 $\pm$ 1.79			
Normalised ACC [n·min ⁻¹]	CWI	0.29 $\pm$ 0.09	0.22 $\pm$ 0.06	0.25 $\pm$ 0.07	0.25 $\pm$ 0.13	0.18 $\pm$ 0.05	0.21	0.10	0.54
	CON	0.19 $\pm$ 0.06	0.17 $\pm$ 0.07	0.13 $\pm$ 0.07	0.13 $\pm$ 0.07	0.23 $\pm$ 0.08			
Normalised DEC [n·min ⁻¹]	CWI	0.35 $\pm$ 0.17	0.14 $\pm$ 0.08	0.33 $\pm$ 0.16	0.38 $\pm$ 0.10	0.30 $\pm$ 0.12	< 0.01	0.63	0.42
	CON	0.28 $\pm$ 0.12	0.18 $\pm$ 0.08	0.31 $\pm$ 0.11	0.31 $\pm$ 0.10	0.32 $\pm$ 0.09			
Normalised ExE [n·min ⁻¹]	CWI	0.42 $\pm$ 0.12	0.35 $\pm$ 0.08	0.46 $\pm$ 0.15	0.37 $\pm$ 0.09	0.32 $\pm$ 0.08	0.62	0.19	0.52
	CON	0.29 $\pm$ 0.08	0.33 $\pm$ 0.08	0.32 $\pm$ 0.12	0.41 $\pm$ 0.17	0.39 $\pm$ 0.10			

624 FM: friendly match; TR: training; TM: tournament match; HSD: high speed distance; ACC: accelerations; DEC: decelerations; ExE: explosive efforts;

625 Int.: intervention

Table 2 Comparison of percentage duration [mean $\pm$ SD; MD and 95% CI] spend in each velocity zone during tournament matches 1-3 combined

Velocity zone	CWI	CON	MD [95% CI]	p- value
---------------	-----	-----	-------------	----------

< 6 km·h ⁻¹	54.8 ± 3.7	56.4 ± 6.0	1.6 [-6.5-3.3]	0.34
6 – 11 km·h ⁻¹	24.4 ± 2.1	23.3 ± 2.8	1.1 [-1.3-3.5]	0.18
11 – 15 km·h ⁻¹	12.9 ± 1.8	12.2 ± 2.6	0.7 [-1.5-2.9]	0.30
15 – 19 km·h ⁻¹	5.4 ± 1.2	5.5 ± 1.5	0.1 [-1.4-1.2]	0.77
19 – 23 km·h ⁻¹	1.9 ± 0.7	1.9 ± 0.5	0.0 [-0.6-0.6]	0.82
> 23 km·h ⁻¹	0.5 ± 0.4	0.5 ± 0.3	0.0 [-0.3-0.3]	0.49

626 CWI: cold water immersion; CON: control

627

Table 3 Comparison of perceptual and HR-based parameters (mean $\pm$ SD) between recovery modalities cold-water immersion (CWI) and control (CON)

Variable	Intervention	Day 1 – FM	Day 2 – TR	Day 3 – TM 1	Day 4 – TM 2	Day 5 – TM 3	[time]	p-value [int.]	[time*int.]
RPE _{session} [AU]	CWI	4.9 $\pm$ 0.6	5.1 $\pm$ 0.7	5.7 $\pm$ 1.3	6.8 $\pm$ 1.7 *	6.0 $\pm$ 1.3 *	< 0.01	0.58	0.81
	CON	5.2 $\pm$ 1.0	5.7 $\pm$ 0.7	5.7 $\pm$ 0.7	6.7 $\pm$ 1.2	6.3 $\pm$ 1.2			
Quality of sleep [AU]	CWI	2.6 $\pm$ 1.2	2.6 $\pm$ 1.0	2.7 $\pm$ 0.8	2.3 $\pm$ 0.9	2.7 $\pm$ 1.1	0.38	0.67	0.81
	CON	2.8 $\pm$ 1.2	2.2 $\pm$ 0.6	3.1 $\pm$ 1.3	2.3 $\pm$ 0.8	2.9 $\pm$ 0.7			
Perceived stress [AU]	CWI	2.6 $\pm$ 0.7	2.1 $\pm$ 0.3	2.1 $\pm$ 0.4	2.2 $\pm$ 1.0	2.8 $\pm$ 1.2 *	0.02	0.87	0.38
	CON	2.2 $\pm$ 0.7	2.5 $\pm$ 0.6	2.3 $\pm$ 1.0	2.0 $\pm$ 0.9	2.8 $\pm$ 0.7			
Perceived recovery [AU]	CWI	3.6 $\pm$ 0.5	3.6 $\pm$ 0.5	4.0 $\pm$ 0.6	3.9 $\pm$ 0.9	3.1 $\pm$ 1.0 *	< 0.01	0.82	0.85
	CON	3.7 $\pm$ 0.8	3.6 $\pm$ 0.7	3.7 $\pm$ 0.8	3.9 $\pm$ 0.8	3.1 $\pm$ 0.6			
HR _{ex} [%HR _{max}]	CWI	77.5 $\pm$ 5.0	79.5 $\pm$ 4.0	73.6 $\pm$ 4.4 *	73.2 $\pm$ 4.9 *	76.0 $\pm$ 3.2	< 0.01	0.67	0.69
	CON	77.2 $\pm$ 3.9	80.6 $\pm$ 4.3	75.8 $\pm$ 4.1	73.9 $\pm$ 4.8	76.2 $\pm$ 3.3			
HRR [%HR _{ex}]	CWI	67.4 $\pm$ 3.2	65.4 $\pm$ 4.0	62.8 $\pm$ 6.3 *	60.3 $\pm$ 6.1 *	62.8 $\pm$ 5.0	< 0.01	0.36	0.26
	CON	70.1 $\pm$ 3.7	64.7 $\pm$ 3.8	62.4 $\pm$ 6.0	63.8 $\pm$ 6.5	66.4 $\pm$ 5.3			

628 FM: friendly match; TR: training; TM: tournament match; RPE: rating perceived exertion; HR_{ex}: exercise heart rate; HR_{max}: maximum heart rate;

629 HRR: heart rate recovery; Int.: intervention; * indicates significant time effects ($p < 0.05$) compared to Day 1

630

631

632

KEY: Questionnaire Biomarkers Heart rate recovery GPS tracking

633

634

635

636