ORIGINAL PAPER

The role of negative cognition, intolerance of uncertainty, mindfulness, and self-compassion in weight regulation among male army recruits.

Michail Mantzios • Janet Clare Wilson • Maggie Linnell • Paul Morris

M. Mantzios • J. C. Wilson • M. Linnell • P. Morris

Department of Psychology, University of Portsmouth, Hampshire, Portsmouth PO1 2DY
e-mail: michael.mantzios@port.ac.uk

Abstract

The present study examined weight gain in a stressful environment (i.e., following compulsory enrollment in the Greek Army). It was predicted that higher levels of mindfulness and self-compassion would relate negatively to weight gain, whilst negative automatic thoughts and intolerance of uncertainty would positively relate to weight gain. This research also explored the strength of independent variables when predicting weight gain, plus the additional contribution of self-compassion when controlling for the effect of mindfulness on weight gain. Ninety-seven military recruits completed the psychological scales on the first day of enrollment. Their height and weight were measured at baseline and after five weeks to record weight gain. Results revealed that 43 participants gained weight, while 54 lost weight. Those who lost weight reported significantly higher scores in mindfulness and self-compassion, whereas those who gained weight reported significantly higher scores in negative automatic thoughts and intolerance of uncertainty. Furthermore, negative automatic thoughts and intolerance of uncertainty did not significantly predict weight gain, after mindfulness and self-compassion were taken into account. Also, self-compassion uniquely contributed to the negative prediction of weight gain, once mindfulness was taken into account. This research concluded that negative cognition may play a role in weight gain; however, mindfulness and self-compassion may be more useful traits in predicting weight gain, given that once they are taken into account, negative cognition stop being significant predictors.

KEYWORDS: Self-compassion; Mindfulness; Weight Regulation; Intolerance of Uncertainty; Negative Automatic Thoughts

Introduction

The increasing number of people gaining weight is of international concern (Finucane, et al., 2011). Weight gain is the result of an imbalance between energy intake and expenditure and is explained by genetic, environmental, and psychological factors (e.g., Cohen & Farley, 2008; Elfhag & Morey, 2008; Maes, Neale, & Eaves, 1997; Wansink, 2004). To overcome the problem, behavioral scientists have attempted to support people with behavioral and cognitive practices, which adjust eating and assists weight loss (e.g., Goodrick, Poston, Kimball, Reeves, & Foreyt, 1998; Telch, Agras, Rossiter, Wilfley, & Kenardy, 1990). However, when dieting, many people lose less weight than required to improve their health, and quickly regain the weight that was lost (Stroebe, Mensink, Aarts, Schut, & Kruglanski, 2008; Wadden & Butryn, 2003).

Indeed, most research on weight management explored regulation in dieters. However, the present study did not require dieting. Instead, this research explored weight gain, which often occurs when one is not actively dieting, but when one is between diets (and unintentionally fails to maintain weight loss after dieting), or, when one is exposed to stressful and uncertain environments (e.g., a competitive work environment or unemployment). In fact, gaining weight is often observed in stressful environments (Greeno & Wing, 1994), especially when those environments project an uncertain future (e.g., Byrne, Cooper & Fairburn, 2003; Elfhag & Morey, 2008; Ganley, 1989), such as joining the armed forces.

Joining the armed forces can be stressful (e.g., Clemons, 1996; Dawson, 2000). However, as all men are automatically obliged to serve one year in the military in Greece, this adds two additional stressors. First, many are ill prepared for a stringent training regime. Second, some

men are obese, making the training particularly challenging. Normally, when one has chosen to join the military as a recruit, one loses a little fat and gains a little muscle mass due to the rigorous physical training regime. On the other hand, food is the only distraction and old habit that recruits are exposed to during their initial training. Preliminary findings showed that about 50% of the recruits unintentionally gained weight on entering the Greek military (Mantzios, 2012). This presented an excellent opportunity to study those who unintentionally gain weight in a stressful environment, despite rigorously exercising. Furthermore, these new recruits are immediately exposed to uncertainties (such as handling weapons, loss of independence and freedom, intimidating physical and psychological demands, involuntary deployment), as well as a restricted diet; all of which can give rise to negative automatic thinking.

At first glance, negative automatic thoughts and, in particular, intolerance of uncertainty may not appear related to weight gain. However, negative automatic thoughts are internal events that disrupt psychological well-being and require management to enable the self-regulation of behavior. In particular, negative automatic thoughts (e.g., 'I'm a failure, I don't know the point of even trying') often sabotage weight regulation efforts (e.g., Kuehnel & Wadden, 1994; Waller, Ohanian, Meyer, & Osman, 2000). Additionally, as they are automatic, they are more difficult to manage (e.g., Stewart & Payne, 2008). When people attempt to control such thoughts, they often find themselves immersed in an on-going battle, amplifying them as they do so (Wegner, 1989; Wegner, Schneider, Carter & White, 1987). If one's main coping strategy for dealing with negative thoughts (e.g., threats to personal feelings of safety) is distracting oneself with food, this will certainly result in weight gain if the thoughts continue. This is not uncommon when joining the military, as this is one of a few distractions available for soldiers who are often deprived

access to other distractions, e.g., television or alcohol (see Johnsen, Laberg & Eid, 1998 for avoidance-coping in military personnel).

There is also a subset of negative automatic thoughts that may cause further problems – specifically, negative thoughts around uncertainty. Intolerance of uncertainty is defined as the inclination to respond adversely to uncertain situations and events (Dugas, Buhr, & Ladouceur, 2004). Intolerance of uncertainty often leads to anxiety and depression (e.g., Asmundson & Carleton, 2005; Berenbaum, Bredemeier, & Thompson, 2008; Holaway, Heimberg, & Coles, 2006; McEvoy & Mahoney, 2011), as well as cognitive avoidance and perfectionism (Buhr & Dugas, 2006; Laugesen, Dugas, & Bukowski, 2003). Although everyone is likely to experience uncertainty, in cases where uncertainty is too frequent or too difficult to manage, it becomes problematic (e.g., keeping a job whilst others experience job loss or being automatically recruited into the military). When unable to tolerate such uncertainties, intolerance may lead to worry, anxiety and depression; all of which have been linked to overeating and weight gain (e.g., Elfhag & Morey, 2008). Furthermore, Heatherton and Baumeister (1991) proposed that people often overeat to escape self-critical thoughts. These self-critical thoughts are frequently associated with uncertainty (e.g., ‘I may screw up; I don’t know how to deal with this situation’). Therefore, this research envisaged that higher levels of negative automatic thoughts and intolerance of uncertainty would relate to weight gain.

Once avoiding negative thoughts in an uncertain and stressful environment may not be a viable option, how one relates to such thoughts may be more promising. Mindfulness allows people to face experiences rather than avoid them, and thus, prevents the escalation of rumination, worries, or negative automatic thoughts (Segal, Williams, & Teasdale, 2002). For example, Verplanken, Friborg, Wang, Trafimow, and Woolf (2007) found a significant negative

correlation between mindfulness and habitual negative thinking. Diminishing the adverse effects of negative thinking through mindfulness is achieved by (a) becoming aware of when and where the automatic thoughts arise, and (b) by reducing the importance of the negative thoughts through acceptance (Verplanken & Tangelder, 2011). In recent years, some weight loss strategies have included mindfulness, and this has been shown to be useful for people who attempt to lose weight (Forman, Butryn, Hoffman, & Herbert, 2009; Lillis, Hayes, Bunting, & Masuda, 2009; Mantzios & Wilson, 2013a; Tapper et al., 2009). However, this research attempted to identify factors that would promote or hinder weight gain, whilst such previous research looks at developing mindfulness as a skill, and not at the trait of mindfulness.

The trait of mindfulness has been described as the experience of centering attention and awareness on what is taking place in the present moment (Brown & Ryan, 2003). Research demonstrates that high scores on the trait of mindfulness predicts lower depression, anxiety, and unpleasant affect (Brown & Ryan, 2003; Shapiro, Oman, Thoresen, Plante, & Flinders, 2008; Weinstein, Brown & Ryan, 2009). Furthermore, higher trait mindfulness also relates to successful self-regulation, including lower reactivity to emotional stimuli (Creswell, Way, Eisenberger, & Lieberman, 2007) and lower automatic and impulsive reactions (Brown & Ryan, 2003; Lattimore, Fisher, & Malinowski, 2011; Levesque & Brown, 2007; Wenk-Sormaz, 2005). However, if one is low on mindfulness, the reverse is more likely to happen; that is, negative automatic thoughts may pull the person away from the present moment and into negative rumination (Verplanken et al., 2007). Thus, those high in mindfulness or who have practiced mindfulness may be able to accept negative automatic thoughts and intolerance of uncertainty and weaken their impact (Verplanken & Tangelder, 2011). Consequently, mindfulness emerges as a factor that could potentially hinder weight gain. However, mindfulness does not target the

suffering that comes with negative automatic thoughts, as mindfulness acknowledges all present moments (positive, negative, neutral) and the experience rather than the individual that has the experience (see Neff & Germer, 2013 for review). Perhaps reframing suffering and reinforcing acceptance as seen in self-compassion research may also be helpful.

Neff (2003a) described self-compassion as experiencing one's own suffering through feelings of kindness toward oneself, with a mindful awareness and the recognition that one's suffering is part of the common human experience (see Neff, 2003a; 2003b for review). Self-compassion is negatively associated with self-criticism, depression, anxiety, rumination, thought suppression, and neurotic perfectionism and positively with life satisfaction and social connectedness (Neff, 2003a; 2003b). Gilbert (2005, 2009) suggested that compassion uniquely stimulates the neural self-soothing system that assists affect regulation and redirects attention with feelings of gentleness and kindness.

There are two ways where self-compassion can be helpful to people who are trying not to gain any weight. First, self-compassion can assist by supporting mindfulness. Indeed, recent research showed that higher scores of self-compassion increased the effectiveness of mindfulness training (Birnie, Speca, & Carlson, 2010); and self-compassion partly mediated the relationship between mindfulness and well-being (Hollis-Walker & Colosimo, 2011). In other words, mindfulness may support efforts to self-regulate, since successful self-regulation depends on paying close attention to present behavior (Baumeister, Heatherton, & Tice, 1994; Carver & Scheier, 1981; Duval & Wicklund, 1972; Mischel, Cantor, & Feldman, 1996). Self-compassion, however, may assist in managing affect in overwhelming situations further, since a sense of common humanity and self-kindness are strengthening acceptance, and may make such occurrences (i.e., adjusting to the military and weight regulation) more tolerable (see Allen &

Leary, 2010). Second, self-compassion can be of assistance directly to eating behaviors.

Consider for instance restrictive eaters, who often display the paradoxical tendency to eat more after breaking their diet, simply as a way of coping with failure to restrict oneself in the first place (Heatherton, Polivy, & Herman, 1990). Adams and Leary (2007) demonstrated that self-compassion can put an end to such paradox, by assisting people in feeling more compassionate about their failings.

Furthermore, the common humanity component of self-compassion has been shown to help military recruits. The belief that ‘we are all in this together’ assisted recruits in reducing stress (Gold & Friedman, 2000), whilst reducing stress may aid people who are attempting to regulate their weight (see Elfhag & Morey, 2008). Therefore, this research expected higher levels of self-compassion and mindfulness to prevent weight gain in stressful situations.

The main aim examined in the current study was to investigate psychological factors relating to weight gain in a stressful environment. Negative automatic thoughts and intolerance of uncertainty were expected to positively predict weight gain, as these would serve to undermine weight regulation efforts. Mindfulness and self-compassion, on the other hand, were expected to negatively predict weight gain, as they should theoretically allow for the acceptance of stressful situations, and thus, reduce the desire to overeat as a coping mechanism. Also, this research tried to identify which independent variables prevailed when tested together, as well as the contribution of self-compassion when accounting for the effect of mindfulness in weight gain.

Method

Participants

A sample of 102 military recruits from an army base in Greece participated in the study. A general assessment of medical and psychological well-being was performed for all recruits (as a military requirement) prior to recruitment. This assessment served as a basis for excluding participants with eating or clinical disorders. Five participants were released from their duty after the first week for medical reasons, and were thus excluded from the analyses. The remaining sample of 97 men had a Body Mass Index of $M = 25.9$ (Age $M = 21.03$ years, $SD = 2.4$ years). None were underweight, 47 were of normal weight ($BMI_{\text{range}} = 18.5\text{-}24.9$), and 50 were overweight; that is, 33 were overweight according to their BMI ($BMI_{\text{range}} = 25\text{-}29.9$), and 17 were obese ($BMI_{\text{range}} = 30\text{-}39.9$) (see WHO, 2012 for BMI particulars).

During the study, no participant maintained the same weight during the five week training period. Therefore, two groups were created - those who gained weight and those who lost weight. Of the 97 participants, 43 gained weight over the five weeks of training ($M = 2.40\text{kg}$, $SD = 1.05$; range 1-7 kg; $BMI = 26.33$) and 54 lost weight ($M = 3.83\text{kg}$, $SD = 1.24$; range = 2-8 kg; $BMI = 24.67$). Note that the BMI of the two groups was not significantly different [$t(95) = -0.30$, $p = .76$], thus any differences between these groups is not due to proportional weight differences. Of those who gained weight, 21 were overweight/obese and 22 were of normal weight; and of those who lost weight, 29 were overweight/ obese and 25 of normal weight.

Instruments

Preceding the description of the scales that were used, it should be noted that all scales were translated into Greek. The translation process that was used was forward-backward translation. Furthermore, all scales were accepted as valid and reliable measures before continuing into any further analyses (see Mantzios, Wilson, & Giannou, 2013 for review).

Self-compassion Scale (Neff, 2003a). The scale calculates the qualities of self-compassion with a high score indicating increased feelings of compassion towards oneself. Responses were given on a 5-point scale from 1 (*almost never*) to 5 (*almost always*) with items such as “When times are really difficult, I tend to be tough on myself”. It is a 26-item scale which is composed of six subscales: self-kindness, self-judgment, common humanity, isolation, mindfulness and over-identification. The original scale demonstrated excellent reliability ($\alpha = .93$; e.g., Neff, 2003a), as did the Greek version ($\alpha = .87$; Mantzios et al., 2013). The Cronbach’s alpha for the present study was $\alpha = .73$.

Mindful Attention and Awareness Scale (Brown & Ryan, 2003). The scale is a 15-item, single factor instrument that measures one’s tendency to function on automatic pilot without attention to present experience, with a high score indicating increased mindfulness. Responses were given on a 6-point scale from 1 (*almost always*) to 6 (*almost never*) and include, for example, “I forget a person’s name almost as soon as I’ve been told it for the first time” and “I rush through activities without being really attentive to them”. The original version of the scale reported a Cronbach’s alpha of $\alpha = .82$ (Brown & Ryan, 2003) and for the Greek scale, researchers reported

a Cronbach's alpha of $\alpha = .86$ (Mantzios et al., 2013). The Cronbach's alpha for the present study was $\alpha = .89$.

Intolerance of Uncertainty Scale (Buhr & Dugas, 2002). The Intolerance of Uncertainty scale is composed of 27 items such as "Uncertainty makes me uneasy, anxious, or stressed" and "When it's time to act, uncertainty paralyses me". Respondents rated on a 5-point scale how strongly each of 27 items best described intolerance of uncertainty with higher scores yielding increased intolerance. The authors of the English scale reported a Cronbach's $\alpha = .94$ and the Greek version was $\alpha = .89$ (Mantzios, 2012). The Cronbach's alpha for the present study was $\alpha = .90$.

Automatic Thoughts Questionnaire (Hollon & Kendall, 1980). The questionnaire is a 30-item questionnaire used to assess negative cognition by measuring the cognitive self-statements of an individual. Responses were given on a 5-point scale from 1 (*not at all*) to 5 (*all the time*) and include items such as "I feel so helpless" and "I wish I were somewhere else". Individuals respond on a 5-point scale with higher scores indicating increased occurrences of negative automatic thoughts. Developers of the scale reported a Cronbach's alpha of $\alpha = .98$ and the Greek version has a Cronbach's alpha of $\alpha = .91$ (Mantzios, 2012). The Cronbach's alpha for the present study was $\alpha = .93$.

Procedure and Design

Potential participants gathered in a theatre and they were told about the study (that is, that it related to weight maintenance in the military). Participants who wished to take part were given a questionnaire pack. As part of the Army procedures, their weight and height were measured at baseline and their weight was measured again five weeks later. The consent form gave

permission to access the weight measurements for the present study. Furthermore those who chose to participate were informed that many soldiers appear to gain weight, thus, they should watch their weight, but not diet or skip meals. Overall, the goal of new recruits was not to gain any weight and maintain their current weight status.

Results

Data overview

The descriptive statistics, including inter-correlations between mindfulness, self-compassion, negative automatic thoughts, intolerance of uncertainty and weight change are presented in Table 1. As shown in Table 1, mindfulness and self-compassion are positively correlated and both relate negatively to negative automatic thoughts and intolerance of uncertainty (all $p < .01$). *Weight gain* (i.e., $\text{weight}_{\text{Follow-up}} - \text{weight}_{\text{Baseline}}$) related negatively to mindfulness and self-compassion and positively to negative automatic thoughts and intolerance of uncertainty.

INSERT TABLE 1 HERE

Main Analyses

Initially, since no other research explored weight gain with negative automatic thoughts and intolerance of uncertainty, this research investigated whether weight gain would be predicted by higher levels of intolerance of uncertainty and negative automatic thoughts. As shown in Table 2, those who gained weight were significantly higher in negative automatic thoughts, including intolerance of uncertainty, than those who lost weight. Those who lost weight were significantly higher in mindfulness and self-compassion, but lower in negative automatic thoughts, including intolerance of uncertainty, than those who gained weight. These findings

were further explored through regressions. Since intolerance of uncertainty is a sub-group of negative automatic thoughts, the chance of multicollinearity among these predictors creating instability in the regression coefficients was high, making it difficult to assess the separate effects of the independent variables. Therefore, two separate linear regressions were conducted. Weight gain was explained by intolerance of uncertainty [adjusted $R^2 = .098$, $F(1, 95) = 11.45$, $p < .001$] and negative automatic thoughts [adjusted $R^2 = .191$, $F(1, 95) = 23.7$, $p < .001$]. The relationship between Weight gain and intolerance of uncertainty ($\beta = .33$, $p < .001$), as well as between weight gain and negative automatic thoughts were positive, ($\beta = -.45$, $p < .001$, see Table 2). Therefore, increases in negative automatic thoughts, including intolerance of uncertainty were associated with higher weight gain.

INSERT TABLE 2 HERE

To determine the effect of all independent variables on weight gain, a multiple regression was performed with self-compassion, mindfulness, negative automatic thoughts, and intolerance of uncertainty. Results indicated that only mindfulness ($\beta = -.40$, $p < .001$) and self-compassion ($\beta = -.45$, $p < .001$) scores significantly predicted Weight Gain [adjusted $R^2 = -.48$, $F(4, 92) = 23.19$, $p < .001$] (see Table 3). Therefore, the results indicated that the effect of negative automatic thoughts and intolerance of uncertainty was non-significant, once the effects of mindfulness and self-compassion were taken into account.

INSERT TABLE 3 HERE

Last, this research aimed to explore if self-compassion accounted for additional variance when predicting weight gain and controlling for mindfulness. A hierarchical regression revealed that self-compassion accounted for additional variance in weight gain when controlling for

mindfulness [$\Delta R^2 = .17$; $\Delta F(2, 94) = 31.4$, $p < .001$]. Therefore, self-compassion appears to uniquely contribute to the prediction of weight gain.

Discussion

The main aim of the current study was to explore psychological factors relating to weight gain in a stressful environment. As expected, it was found that positive predictors of weight gain were negative automatic thoughts and intolerance of uncertainty, while self-compassion and mindfulness related negatively to weight gain. However, results were not that straightforward.

To begin with, results revealed that 44% of the sample gained weight, while 56% lost weight and none of the participants maintained their weight, thus, replicating previous findings in new military recruits (Mantzios, 2012). Exploring those groups further revealed that those who gained weight were significantly higher in negative automatic thoughts and intolerance of uncertainty, than those who lost weight. Furthermore, those who lost weight were significantly higher in mindfulness and self-compassion than those who gained weight.

Further analyses revealed that negative automatic thoughts and intolerance of uncertainty were predictors of weight gain when examined independently; although they did not significantly predict weight gain once the effect of mindfulness and self-compassion were taken into account. People usually dwell over experiences or try to avoid them. Either way, the effects are usually negative (e.g., Cribb, Moulds, & Carter, 2006), explaining the relationship between negative cognitions and weight gain (see also Mantzios & Wilson, 2013b). On the other hand, instead of ruminating or avoiding an experience, there is the option of relating differently to the experience,

and such difference is best described through acceptance. Mindfulness enables acceptance and assists in acknowledging all experiences, as well as in seeing thoughts as thoughts (instead of rigid expressions that require rumination, avoidance, or, suppression). This way, negative cognitions may exist, but not affect the way people react to them when they are around food.

Furthermore, self-compassion and mindfulness work differently and add something unlike to the negative prediction of weight gain, as self-compassion appeared to explain an additional amount of variance when controlling for mindfulness. Apart from the reasons given in the introduction, another way that could explain the additional variance is the emphasis on suffering instead of whatever arises in the present moment (see Neff & Germer, 2013). This emphasis on suffering may be another feature that assists weight regulation behavior. In reality, managing weight translates into two main decisions for weight watchers. First, people restrict themselves and regulate their food intake, which adds some form of deprivation and suffering to the present moment. The second choice is for people to give into the temptation of their favorite foods, but such decisions are usually followed by feelings of distress and shame afterwards. Regardless of the choice one makes when managing weight, there is always some form of suffering that needs to be handled with gentleness, a kind attitude and more compassion. Future research should explore eating behavior and the role of suffering and self-compassion more thoroughly.

The new military recruits that were used as a sample added some benefits and some limitations to the current study. The benefits included having a controlled environment to examine the psychological traits. All participants were eating the same food (although they had some control over portions) and were exercising at the same level, thus, living in an environment where energy intake and expenditure was relatively controlled. Furthermore, the participants

followed a routine day that was highly planned and controlled (e.g., waking up at 6 a.m., in bed by 10 p.m., eating at specific times and places, and only specific food being available, exercising at certain times), making everyday uncertainty negligible. However, the uncertainty of whether they could cope effectively with such a regime was still very much present (e.g., Hogg, 2007). Furthermore, the automatic recruitment of civilians allowed for a relatively randomized sample from the general population and included those who fell within a wide range on the Body Mass Index, allowing therefore for an exploration of weight gain that is apparent in both average, as well as overweight populations.

On the other hand, limitations were evident also. First, the absence of female participants and the unusual military environment (i.e., being constantly under pressure for five weeks) suggest that caution should be used when generalizing and interpreting the present results. People who serve in the military usually need to adjust and tolerate the anxiety and uncertainty of using weapons (and threats to personal safety), communal living (and loss of personal privacy, as well as sleep deprivation), leaving behind civilian rights and belongings (i.e., isolation from former social networks, loss of independence and freedom, loss of control over one's life). All of these factors may have strengthened the relationships observed in this study, especially with regards to negative cognitions. Second, future studies may benefit by examining specific uncertainties associated to weight regulation versus more global life-based uncertainties. For example, future studies could explore whether people who successfully manage their weight do so because they are fine with uncertainty (both in weight management and/or universally). Third, almost half of the sample was of an average weight (according to their BMI) and participants were not asked to diet. In fact, if participants were asked to diet (or if the sample was of a higher BMI), results could have been more significant, but such assumptions remain to be explored in

future research. Fourth, weight gain could represent an increase in muscle mass (instead of adipose). Considering the daily exercise routine, the foot patrols, and the frequent engagement in physical activities (such as domestic chores), there is a chance that the weight increase resulted from the labor-intensive environment. Future research should address this limitation by including measures that will measure changes in adipose (and in effect, determine changes in muscle mass).

In conclusion, this research complements the potential and significance of mindfulness and self-compassion in weight regulation research. The findings of the present research also provide preliminary evidence that mindfulness and self-compassion may together be a useful strategy to aid weight maintenance, despite of any negative cognition that may be present. Experimental research could clarify the relationship between mindfulness and self-compassion with negative cognitions and uncertainty further. This research suggests that both mindfulness and self-compassion may be of assistance; consequently, future research should preferably use both components, instead of one or the other.

References

- Adams, C. E., & Leary, M. R. (2007). Promoting self-compassionate attitudes toward eating among restrictive and guilty eaters. *Journal of Social and Clinical Psychology, 26*(10), 1120-1144.
- Allen, A. B., & Leary, M. R. (2010). Self-compassion, stress and coping. *Social and Personality Psychology Compass, 4*(2), 107-118.
- Asmundson, G. J. G., & Carleton, R. N. (2005). Fear of pain is elevated in adults with co-occurring trauma-related stress and social anxiety symptoms. *Cognitive Behavior Therapy, 34*(4), 248–255.
- Baumeister, R. F., Heatherton, T. F., Tice, D. M. (1994). *Losing Control: How and Why People Fail at Self-Regulation*. San Diego: Academic Press.
- Berenbaum, H., Bredemeier, K., & Thompson, R.J. (2008). Intolerance of uncertainty: Exploring its dimensionality and associations with need for cognitive closure, psychopathology, and personality. *Journal of Anxiety Disorders, 22*(1), 117–125.
- Birnie, K., Speca, M., & Carlson, L. E. (2010). Exploring self-compassion and empathy in the context of mindfulness-based stress reduction (MBSR). *Stress and Health, 26*(5), 359-371.
- Brown, K. W., & Ryan, R. M. (2003). The benefits of being present: Mindfulness and its role in psychological well-being. *Journal of Personality and Social Psychology, 84*(4), 822-848.

- Buhr, K., & Dugas, M. J. (2002). The intolerance of uncertainty scale: Psychometric properties of the English version. *Behavior Research and Therapy*, 40(8), 931-945.
- Buhr, K., & Dugas, M. J. (2006). Investigating the construct validity of intolerance of uncertainty and its unique relationship with worry. *Journal of Anxiety Disorders*, 20(2), 222-236.
- Byrne, S., Cooper, Z., & Fairburn, C. (2003). Weight maintenance and relapse in obesity: a qualitative study. *International Journal of Obesity*, 27(8), 955-962.
- Carver, C. S., & Scheier, M. (1981). *Attention and self-regulation: A control-theory approach to human behavior*. New York: Springer.
- Clemons, E. P. (1996). Monitoring anxiety levels and coping skills among military recruits. *Military Medicine*, 161(1), 18-21.
- Cohen, D., & Farley, T. (2008). Eating as an automatic behavior. *Preventing Chronic Disease*, 5(1), A23.
- Creswell, J. D., Way, B. M., Eisenberger, N. I., & Lieberman, M. D. (2007). Neural correlates of dispositional Mindfulness during affect labelling. *Psychosomatic Medicine*, 69(6), 560-565.
- Cribb, G., Moulds, M. L., & Carter, S. (2006). Rumination and experiential avoidance in depression. *Behavior Change*, 23(3), 165-176.
- Dawson, K. J. (2000). Psychological adjustment to basic military training: application of a stress and coping model (Unpublished thesis). University of Queensland, Australia.

- Dugas, M. J., Buhr, K., & Ladouceur, R. (2004). The role of intolerance of uncertainty in the etiology and maintenance of generalized anxiety disorder. In: R. G. Heimberg, C. L. Turk, & D. S. Mennin (Eds.), *Generalized anxiety disorder: advances in research and practice* (pp. 143–163). New York: Guilford Press.
- Dugas, M. J., Hedayati, M., Karavidas, A., Buhr, K., Francis, K., & Phillips, N. A. (2005). Intolerance of uncertainty and information processing: Evidence of biased recall and interpretations. *Cognitive Therapy and Research*, 29(1), 57-70.
- Duval, S., & Wicklund, R. A. (1972). *A theory of objective self-awareness*. New York: Academic Press.
- Elfhag, K., & Morey, L. C. (2008). Personality traits and eating behavior in the obese: poor self-control in emotional and external eating but personality assets in restrained eating. *Eating Behaviors*, 9(3), 285-293.
- Finucane, M. M., Stevens, G. A., Cowan, M. J., Danaei, G., Lin, J. K., Paciorek, C. J., . . . Bahalim, A. N. (2011). National, regional, and global trends in body-mass index since 1980: systematic analysis of health examination surveys and epidemiological studies with 960 country-years and 9· 1 million participants. *The Lancet*, 377(9765), 557-567.
- Forman, E. M., Butryn, M. L., Hoffman, K. L., & Herbert, J. D. (2009). An open trial of an acceptance-based behavioral intervention for weight loss. *Cognitive and Behavioral Practice*, 16(2), 223-235.
- Fujita, K., & Roberts, J. C. (2010). Promoting prospective self-control through abstraction. *Journal of Experimental Social Psychology*, 46(6), 1049-1054.

- Ganley, R. M. (1989). Emotion and eating in obesity: a review of the literature. *International Journal of Eating Disorders*, 8(3), 343-61.
- Gilbert, P. (2005). *Compassion: Conceptualisations, research and use in psychotherapy*. New York: Routledge.
- Gilbert, P. (2009). *The compassionate mind*. London: Constable.
- Gold, M. A., & Friedman, S. B. (2000). Cadet basic training: An ethnographic study of stress and coping. *Military Medicine*, 165 (2), 147-152.
- Goodrick, G. K., Poston II, W. S. C., Kimball, K. T., Reeves, R. S., & Foreyt, J. P. (1998). Non-dieting versus dieting treatment for overweight binge-eating women. *Journal of Consulting and Clinical Psychology*, 66(2), 363-368.
- Greeno, C. G., & Wing, R. R. (1994). Stress-induced eating. *Psychological Bulletin*, 115(3), 444-464.
- Heatherton, T. F., & Baumeister, R. F. (1991). Binge eating as escape from self-awareness. *Psychological Bulletin*, 110(1), 86-108.
- Herman, C. P., & Mack, D. (1975). Restrained and unrestrained eating. *Journal of Personality*, 43(4), 647-660.
- Hogg, M. A. (2007). Uncertainty–identity theory. *Advances in Experimental Social Psychology*, 39, 69-126.

- Holaway, R. M., Heimberg, R. G., & Coles, M. E. (2006). A comparison of intolerance of uncertainty in analogue obsessive-compulsive disorder and generalized anxiety disorder. *Journal of Anxiety Disorders, 20*, 157–174.
- Hollis-Walker, L., & Colosimo, K. (2011). Mindfulness, self-compassion, and happiness in non-meditators: A theoretical and empirical examination. *Personality and Individual Differences, 50*(2), 222-227.
- Hollon, S. D., & Kendall, P. C. (1980). Cognitive self-statements in depression: Development of an automatic thoughts questionnaire. *Cognitive Therapy and Research, 4*(4), 383-395.
- Johnsen, B. H., Laberg, J. C., & Eid, J. (1998). Coping strategies and mental health problems in a military unit. *Military Medicine, 163* (9), 599-602.
- Kuehnel, R. H., & Wadden, T. A. (1994). Binge eating disorder, weight cycling, and psychopathology. *International Journal of Eating Disorders, 15*(4), 321-329.
- Lattimore, P., Fisher, N., & Malinowski, P. (2011). A cross-sectional investigation of trait disinhibition and its association with mindfulness and impulsivity. *Appetite, 56*(2), 241-248.
- Laugesen, N., Dugas, M. J., & Bukowski, W. M. (2003). Understanding adolescent worry: the application of a cognitive model. *Journal of Abnormal Child Psychology, 31*(1), 55-64.
- Levesque, C., & Brown, K. W. (2007). Mindfulness as a moderator of the effect of implicit motivational self-concept on day-to-day behavioural motivation. *Motivation and Emotion, 31*(4), 284-299.

- Lillis, J., Hayes, S. C., Bunting, K., & Masuda, A. (2009). Teaching acceptance and Mindfulness to improve the lives of the obese: A preliminary test of a theoretical model. *Annals of Behavioral Medicine*, 37(1), 58-69.
- Maes, H., H., M., Neale, M., C., & Eaves, L., J. (1997). Genetic and environmental factors in relative body weight and human adiposity. *Behavior Genetics*, 27(4), 325-351.
- Mantzios, M. (2012). *The development of self-compassion and mindfulness as a possible intervention for weight loss* (Unpublished doctoral dissertation). University of Portsmouth, United Kingdom.
- Mantzios, M. & Wilson, J. C. (2013a). Exploring Mindfulness and Mindfulness with Self-Compassion centred interventions to assist weight loss – Theoretical considerations and preliminary results of a longitudinal study (Submitted for publication)
- Mantzios, M. & Wilson, J. C. (2013b). Making concrete construals mindful: A novel approach of developing mindfulness and self-compassion to assist weight loss, *Psychology & Health*, 1-41.
- Mantzios, M., & Wilson, J. C. (2013). Psychometric properties of the Greek versions of the Self-Compassion and Mindful Attention and Awareness scales, *Mindfulness*, 1-10.
- McEvoy, P. M., & Mahoney, A. E. J. (2011). Achieving certainty about the structure of intolerance of uncertainty in a treatment seeking sample with anxiety and depression. *Journal of Anxiety Disorders*, 25(1), 112–122.
- Mischel, W., Cantor, N., & Feldman, S. (1996). Principles of self-regulation: The nature of willpower and self-control. In: E. T., Higgins, A. W. Kruglanski, (Eds). *Social Psychology: Handbook of Basic Principles* (pp. 329–360). New York: Guilford Press.

- Neff, K., D. (2003a). The development and validation of a scale to measure Self-compassion. *Self and Identity*, 2(3), 223-250.
- Neff, K., D. (2003b). Self-compassion: An alternative conceptualization of a healthy attitude toward oneself. *Self and Identity*, 2(2), 85-101.
- Neff, K. D., & Germer, C. K. (2013). A pilot study and randomized controlled trial of the mindful self-compassion program. *Journal of Clinical Psychology*, 69(1), 28-44.
- Segal, Z. V., Williams, J. M. G., & Teasdale, J. D. (2002). *Mindfulness-based cognitive therapy for depression: A new approach to preventing relapse*. New York: Guilford Press.
- Shapiro, S. L., Oman, D., Thoresen, C. E., Plante, T. G., & Flinders, T. (2008). Cultivating Mindfulness: effects on well-being. *Journal of Clinical Psychology*, 64(7), 840-862.
- Stewart, B. D., & Payne, B. K. (2008). Bringing automatic stereotyping under control: Implementation intentions as efficient means of thought control. *Personality and Social Psychology Bulletin*, 34(10), 1332-1345.
- Stroebe, W., Mensink, W., Aarts, H., Schut, H., & Kruglanski, A. W. (2008). Why dieters fail: Testing the goal conflict model of eating. *Journal of Experimental Social Psychology*, 44(1), 26-36.
- Tapper, K., Shaw, C., Ilsley, J., Hill, A. J., Bond, F. W., & Moore, L. (2009). Exploratory randomised controlled trial of a Mindfulness-based weight loss intervention for women. *Appetite*, 52(2), 396-404.

- Telch, C. F., Agras, W. S., Rossiter, E. M., Wilfley, D., & Kenardy, J. (1990). Group cognitive-behavioural treatment for the non-purging bulimic: An initial evaluation. *Journal of Consulting and Clinical Psychology*, 58(5), 629-635.
- Verplanken, B., & Tangelder, Y. (2011). No body is perfect: The significance of habitual negative thinking about appearance for body dissatisfaction, eating disorder propensity, self-esteem, and snacking. *Psychology and Health*, 26(6), 685–701.
- Verplanken, B., Friborg, O., Wang, C. E., Trafimow, D., & Woolf, K. (2007). Mental habits: Metacognitive reflection on negative self-thinking. *Journal of Personality and Social Psychology*, 92(3), 526–541.
- Wadden, T. A., & Butryn, M. L. (2003). Behavioral treatment of obesity. *Endocrinology and Metabolism Clinics of North America*, 32(4), 981-1003.
- Waller, G., Ohanian, V., Meyer, C., & Osman, S. (2000). Cognitive content among bulimic women: The role of core beliefs. *International Journal of Eating Disorders*, 28(2), 235-241.
- Wansink, B. (2004). Environmental factors that increase the food intake and consumption volume of unknowing consumers. *Annual Review of Nutrition*, 24, 455-479.
- Wegner, D. M. (1989). *White Bears and Other Unwanted Thoughts*. New York: Penguin.
- Wegner, D. M., Schneider, D. J., Carter, S. R., White, T. L. (1987). Paradoxical effects of thought suppression. *Journal of Personality and Social Psychology*, 53(1), 5-13.

Weinstein, N., Brown, K. W., & Ryan, R. M. (2009). A multi-method examination of the effects of Mindfulness on stress attribution, coping, and emotional well-being. *Journal of Research in Personality*, 43(3), 374-385.

Wenk-Sormaz, H. (2005). Meditation can reduce habitual responding. *Advances in Mind-Body Medicine*, 21(3-4), 33-49.

World Health Organization (2012). Fact Sheet No. 311. Available at:

<http://www.who.int/mediacentre/factsheets/fs311/en/>. Accessed: June 17, 2012.

Table 1

Correlation coefficients between Mindfulness, Self-compassion, Negative Automatic Thoughts, Intolerance of Uncertainty, and Weight Maintenance ($n=97$). Also, a summary of Mean scores (M) and Standard Deviations (SD) are presented.

Measures	1	2	3	4	M	SD
(1) Mindful Attention Awareness Scale	-				4.47	.94
(2) Self Compassion Scale	.393**	-			3.38	.44
(3) Automatic Thoughts Questionnaire	-.431**	-.548**	-		2.54	.66
(4) Intolerance of Uncertainty Scale	-.349**	-.518**	.610**	-	1.94	.49
(5) Weight Gain	-.571**	-.602**	.447**	.322**	1.07	3.32

Note: * $p < .05$; ** $p < .01$

Table 2

Means and Standard Deviations of Mindfulness, Self-Compassion, Automatic Thoughts and Intolerance of Uncertainty according to weight change, i.e., weight loss (n=54) compared to weight gain (n=43).

Measures	Weight		<i>SD</i>	<i>t</i>	Significance
	Change	Mean			
Mindfulness Attention and Awareness Scale	Loss	4.99	0.71	-7.87	< .001
	Gain	3.81	0.77		
Self Compassion Scale	Loss	3.63	0.37	-9.13	< .001
	Gain	3.05	0.26		
Automatic Thought Questionnaire	Loss	52.04	9.72	5.25	< .001
	Gain	66.12	16.42		
Intolerance of Uncertainty Scale	Loss	63.17	16.34	3.60	.001
	Gain	75.47	17.22		

Table 3

Summary of multiple regression for Self-compassion, Mindfulness, Negative Automatic Thoughts, Intolerance of Uncertainty in predicting Weight Maintenance.

Variables	B	SE B	Beta	<i>t</i>	Significance	Confidence Intervals	
						Lower	Upper
SCS	-3.27	.68	.45	-4.82	<.001	1.92	4.61
MAAS	-1.37	.29	.39	-4.66	<.001	.77	1.95
NATQ	.02	.02	-.09	.91	.368	-.07	.03
IUS	-.02	.02	.10	-.99	.324	-.02	.05

Note: NATQ=Negative Automatic Thought Questionnaire, IUS= Intolerance of Uncertainty Scale, MAAS= Mindful Attention and Awareness Scale, SCS= Self-compassion Scale.