

# A black hole mass threshold from non-singular quantum gravitational collapse

Martin Bojowald<sup>1</sup>, Rituparno Goswami<sup>2</sup>, Roy Maartens<sup>3</sup>, Parampreet Singh<sup>4</sup>

<sup>1</sup>*Max-Planck-Institut für Gravitationsphysik, Albert-Einstein-Institut, D-14476 Potsdam, Germany*

<sup>2</sup>*Tata Institute for Fundamental Research, Colaba, Mumbai, India*

<sup>3</sup>*Institute of Cosmology and Gravitation, University of Portsmouth, Portsmouth PO1 2EG, UK and*

<sup>4</sup>*Institute for Gravitational Physics and Geometry, Pennsylvania State University, PA 16802, USA*

(Dated: February 7, 2008)

Quantum gravity is expected to remove the classical singularity that arises as the end-state of gravitational collapse. To investigate this, we work with a toy model of a collapsing homogeneous scalar field. We show that non-perturbative semi-classical effects of Loop Quantum Gravity cause a bounce and remove the black hole singularity. Furthermore, we find a critical threshold scale, below which no horizon forms – quantum gravity may exclude very small astrophysical black holes.

PACS numbers: 04.60.Pp, 04.70.Dy, 97.60.Lf

Singularity formation during gravitational collapse signals the breakdown of classical general relativity. In a more complete theory of quantum gravity the singularity should be removed. However, a satisfactory quantum gravity theory has yet to be developed. In addition, the dynamics of general collapse is very complicated. Thus we can only expect to make partial progress in tackling the problem, using a candidate for quantum gravity and a collapse model that is simple enough to be tractable.

A non-perturbative approach to quantizing gravity is loop quantum gravity or quantum geometry [1], which gives rise to a discrete spatial structure [2] and whose successes include prediction of black hole entropy [3]. Applied to the early universe, loop quantum effects can remove the big bang singularity [4]. A natural question is: do these effects also remove the black hole singularity as the end-state of collapse? Techniques to handle inhomogeneous systems are under development and give promising indications [5], but they do not easily reveal the physical picture. We thus consider a simple toy model of a collapsing *homogeneous* scalar field. Classically, this model always produces a black hole, but we show that loop quantum effects change this situation dramatically.

Since we do not yet know semi-classical non-perturbative effects in inhomogeneous cases, we are unable to perform our analysis in the general case. However, when we split the system into a homogeneous star interior and an inhomogeneous outside region, known quantum effects in the interior can be carried into the exterior indirectly through matching conditions. The collapsing homogeneous scalar field cannot be matched to a Schwarzschild exterior because the pressure does not vanish at the boundary. But in any case, we expect that quantum effects will include small non-stationary corrections and thus use a non-stationary spherically symmetric exterior. The generalized Vaidya metric provides a reasonable starting point. It is sufficiently general to allow for a broad range of behavior, including radiative effects.

Our analysis is based on effective equations for the interior which have been established in the cosmological

setting. Fundamentally, the evolution is described by a wave function subject to a difference equation, and effective equations describe the motion of semiclassical wave packets [6]. As long as one stays in semiclassical regimes, which can e.g. be checked using the size of curvature, one gets reliable expectations for the quantum situation.

We first review the classical collapse and the inevitability of a black hole singularity covered by a horizon, for any initial mass. The isotropic interior metric is [7]

$$ds^2 = -dt^2 + a(t)^2(1 + r^2/4)^{-2} [dr^2 + r^2 d\Omega^2], \quad (1)$$

and the massless scalar field  $\phi(t)$  has pressure and energy density  $p = \rho = \frac{1}{2}\dot{\phi}^2$ . The Friedmann equation is

$$\dot{a}^2/a^2 = 4\pi\ell_p^2 \dot{\phi}^2/3 - 1/a^2. \quad (2)$$

The Klein-Gordon equation,  $a\ddot{\phi} + 3\dot{a}\dot{\phi} = 0$ , has solution

$$\dot{\phi} = L/a^3, \quad (3)$$

where  $L$  is a length scale associated with the maximal size of the collapse region, since (2) implies

$$a \leq a_m \equiv (4\pi/3)^{1/4} \sqrt{\ell_p L}. \quad (4)$$

At the singularity  $a \rightarrow 0$ , we have  $\dot{\phi}, \rho \rightarrow \infty$ . The solution of the Friedmann equation is

$$t - t_0 = a_m \int_{a/a_m}^{a_0/a_m} \frac{b^2 db}{\sqrt{1 - b^4}}, \quad (5)$$

where  $a_0 (\leq a_m)$  gives the initial size of the collapse region at time  $t_0$ . The singularity  $a = 0$  is covered by a horizon (see below), and reached in finite proper time for any  $a_0$ :

$$\frac{1}{a_m} (t_s - t_0) < \int_0^1 \frac{db}{\sqrt{1 - b^4}} = \frac{1}{\sqrt{2}} F\left(\frac{\pi}{2}, \frac{1}{\sqrt{2}}\right), \quad (6)$$

where  $F$  is an elliptic integral of the first kind.

We now add non-perturbative modifications to the dynamics, motivated by loop quantum gravity [8]. The quantization introduces a fundamental length scale

$$\ell_* = 0.28\sqrt{j}\ell_p, \quad (7)$$

where  $j(> 1)$  is a half-integer that is freely specifiable. For  $a < \ell_*$ , the dynamics is increasingly different from general relativity. For  $a \lesssim \ell_p$ , the continuum approximation to the spacetime geometry begins to break down, and the fully quantum gravity regime is reached. In the intermediate regime  $\ell_p \lesssim a \lesssim \ell_*$ , loop quantum effects may be treated semi-classically, i.e., the spacetime metric behaves classically, while the dynamics acquires non-perturbative modifications to general relativity [6]. The non-perturbative semi-classical regime exists provided  $\ell_* \gg \ell_p$ , i.e., for  $j \gg 1$ .

The key feature of the loop quantization scheme is the prediction that the geometrical density,  $1/a^3$ , does not diverge as  $a \rightarrow 0$ , but remains finite. The expectation values of the density operator are approximated by  $d_j(a) = D(a)a^{-3}$ , where the loop quantum correction factor is [9]

$$D(a) = (8/77)^6 q^{3/2} \left\{ 7 \left[ (q+1)^{11/4} - |q-1|^{11/4} \right] - 11q \left[ (q+1)^{7/4} - \text{sgn}(q-1)|q-1|^{7/4} \right] \right\}^6, \quad (8)$$

with  $q \equiv a^2/\ell_*^2$ . In the classical limit we recover the expected behavior of the density, while the quantum regime shows a radical departure from classical behavior:

$$a \gg \ell_* : D \approx 1 \quad , \quad a \ll \ell_* : D \approx (12/7)^6 (a/\ell_*)^{15}. \quad (9)$$

Then  $d_j$  remains finite as  $a \rightarrow 0$ , unlike in conventional quantum cosmology, thus evading the problem of the big-bang singularity in a closed model [10]. Intuitively, one can think of the modified behavior as meaning that gravity, which is classically always attractive, becomes repulsive at small scales when quantized. This effect can produce a bounce where classically there would be a singularity, and can also provide a new mechanism for high-energy inflationary acceleration [11]. In the semi-classical regime (where the spectrum can be treated as continuous),  $d_j$  has a smooth transition from classical to quantum behavior, varying from  $a^{-3}$  to  $a^{12}$ . We emphasize that this is but one possibility for a bounce which we use for concreteness, while bounces in general appear more generically in loop cosmology [12].

In loop cosmology the Hamiltonian of a scalar field in a closed universe is

$$\mathcal{H} = a^3 V(\phi) + d_j P_\phi^2/2, \quad P_\phi = d_j^{-1} \dot{\phi}, \quad (10)$$

where  $P_\phi$  is the momentum canonically conjugate to  $\phi$ . This leads to a modified Friedmann equation [11, 13],

$$\frac{\dot{a}^2}{a^2} = \frac{8\pi\ell_p^2}{3} \left[ V(\phi) + \frac{1}{2D} \dot{\phi}^2 \right] - \frac{1}{a^2}, \quad (11)$$




FIG. 1: The scale factor  $a(t)$  of the collapsing interior, for classical (dashed) and semi-classical quantum dynamics (solid).

and a modified Klein-Gordon equation [14]

$$\ddot{\phi} + 3a^{-1}\dot{a}(1-\alpha)\dot{\phi} + DV(\phi) = 0, \quad \alpha \equiv a\dot{D}/(3\dot{a}D). \quad (12)$$

For  $a \ll \ell_*$ , we have  $\alpha \rightarrow 5$ , whereas classically  $D = 1$  and hence  $\alpha = 0$ . Thus in the semi-classical regime,  $0 < \alpha \leq 5$ .

For a massless scalar field,  $V = 0$ , the solution of Eq. (12), generalizing Eq. (3), is

$$\dot{\phi} = Ld_j(a), \quad (13)$$

so that  $P_\phi = L = \text{const}$ . Then the Friedmann equation becomes

$$\dot{a}^2 + 1 = D(a)(a_m/a)^4. \quad (14)$$

The energy density and pressure are modified as  $\rho = \dot{\phi}^2/2D$ ,  $p = \dot{\phi}^2(1-\alpha)/2D$ , so that

$$w \equiv p/\rho = 1 - \alpha. \quad (15)$$

(The modified  $\rho$  and  $p$  satisfy the usual conservation equation if  $\phi$  satisfies the modified Klein-Gordon equation.) Since  $\alpha$  varies from 0 to 5 as  $a$  decreases, the  $\dot{\phi}$  term in Eq. (12), which classically behaves as *anti-frictional* during collapse, starts to behave as *frictional* when  $\alpha > 1$ . Thus, contrary to classical behavior, where  $\dot{\phi}$  increases as  $a$  decreases, in the semi-classical regime the scalar field starts slowing down with collapse. In fact at  $\alpha = 2$  the magnitude of the frictional term becomes exactly equal to the classical anti-frictional term. Thereafter at smaller values of the scale factor the term becomes increasingly frictional and the collapse further slows down, and may turn around.

The point where  $\alpha = 2$  is also the point beyond which the null energy condition is violated:  $w < -1$ , by Eq. (15). Violations of the null energy condition by

quantum gravity effects are to be expected, and in loop quantum gravity this occurs for  $\alpha > 2$ , when the scalar field effectively behaves as a “phantom” field.

In order to see qualitatively how the non-perturbative frictional quantum effects remove the classical singularity, we assume that, over a small interval of scale factor, we can take  $\alpha \approx \text{constant}$ , so that  $D \approx D_* (a/\ell_*)^{3\alpha}$ , where  $D_*$  is a dimensionless constant. By Eq. (13),

$$\dot{\phi} \approx LD_* \ell_*^{-3\alpha} a^{3(\alpha-1)}, \quad (16)$$

which shows how the kinetic energy decreases with decreasing  $a$  when  $\alpha > 1$ , contrary to the classical case. The modified Friedmann equation (14) gives

$$\dot{a}^2 \approx (a_m^4 \ell_*^{-3\alpha} D_*) a^{3\alpha-4} - 1. \quad (17)$$

In general relativity, where  $\alpha = 0$  and  $D_* = 1$ , this shows that for  $a < a_m$ , there is no turning point in  $a$ , i.e.,  $\dot{a} \neq 0$ . With loop quantum effects, for  $\alpha > \frac{4}{3}$ , the equation  $\dot{a}(t_c) = 0$  has a solution,  $a_c \approx (\ell_*^{3\alpha}/D_* a_m^4)^{1/(3\alpha-4)} \ll a_m$ . Thus the collapse leads to a bounce and singularity avoidance. The numerical integration of the modified Friedmann and Klein-Gordon equations confirms the qualitative analysis, and the results are illustrated in Fig. 1. As is clear from the figure, the classical curve (dashed line) hits the singularity in finite time, whereas the quantum-corrected curve bounces and avoids the singularity. The key question is whether a horizon forms in the quantum-corrected collapse.

The formation or avoidance of the singularity  $a = 0$  is independent of the matching to the exterior. But in order to understand horizon formation in the semi-classical quantum case, we need to impose the matching conditions. Since the pressure is nonzero at the boundary, given in comoving coordinates by  $r = R = \text{constant}$ , the interior cannot be directly matched to a static Schwarzschild exterior. However we can match to an intermediate non-stationary region – for example, a generalized Vaidya region [15],

$$ds^2 = -[1 - 2M(v, \chi)/\chi] dv^2 + 2dv d\chi + \chi^2 d\Omega^2. \quad (18)$$

The usual Vaidya mass  $M/\ell_p^2$  is generalized so that  $\partial M/\partial \chi$  may be nonzero. The total mass measured by an asymptotic observer is  $m = m_M + m_\phi$ , where  $m_M$  is the total mass in the generalized Vaidya region, and  $m_\phi = \int \rho dV$  the interior mass. By Eqs. (1), (4) and (13),

$$\frac{m_\phi}{m_p} = \frac{3a}{2\ell_p} \left(\frac{a_m}{a}\right)^4 D(a) \left[ \tan^{-1} \frac{R}{2} - \frac{R(1-R^2/4)}{2(1+R^2/4)^2} \right]. \quad (19)$$

Since we do not specify the matter content in the exterior, and do not know the modified field equations there, we cannot determine  $M(v, \chi)$  and thus  $m_M$ . However, as we discuss below, we can still draw qualitative conclusions about the behavior of horizons close to the matter shells.




FIG. 2: The speed of collapse,  $|\dot{a}|$ , against the scale factor  $a$ , for the evolution shown in Fig. 1, up to the bounce. The dashed curve is for classical dynamics and semi-classical quantum dynamics gives the solid curve. The horizontal dotted lines correspond to different values of  $R$  in Eq. (24): for the upper line there is no horizon in the quantum-corrected case, the middle line corresponds to the threshold for a horizon, the lower line to the case of an inner and outer horizon.

Matching the 1st and 2nd fundamental forms we obtain

$$\chi(v) = Ra(t)/(1 + R^2/4), \quad (20)$$

$$dv/dt = (1 + R^2/4)/(1 - R^2/4 - R\dot{a}), \quad (21)$$

$$2M = aR^3(\dot{a}^2 + 1)/(1 + R^2/4)^3, \quad (22)$$

$$-M_{,v} = \chi_{,vv} + (1 - 2M/\chi - 3\chi_{,v})(M/\chi - M_{,\chi}). \quad (23)$$

The exterior region can contain trapped surfaces when the condition  $2M(v, \chi) = \chi$  is satisfied. Evaluating this at the matching surface, using Eqs. (20) and (22), gives

$$|\dot{a}| = R^{-1}(1 - R^2/4). \quad (24)$$

When this value is reached, a dynamical horizon [16] intersects the matching surface. This always occurs classically since during the collapse  $|\dot{a}|$  varies from zero to infinity. With the modified dynamics, however,  $|\dot{a}|$  is *bounded* throughout the evolution, so that *it depends on initial values whether or not a horizon forms* (Fig. 2). Moreover, after the bounce,  $\dot{a}$  grows again, so that the condition can be satisfied a second time. This results in a picture where the bounce, replacing the classical singularity, may be shrouded by an evaporating dynamical horizon outside, as shown in Fig. 3. There will be a second point where the horizon condition is satisfied since  $|\dot{a}|$  decreases between the peak of  $d_j(a)$  and the bounce.

When it intersects the matching surface, the horizon is always null, as follows from Eq. (23). Its later behavior depends on the details of the outer region, which can not be determined here. Nevertheless, one can expect that both horizons will become timelike and evaporate. Horizon evaporation in this model does not only come from




FIG. 3: Eddington-Finkelstein diagram of the collapse, with boundary  $\Sigma$ . Dotted lines show constant  $v$  (outside) and constant  $t$  (inside). Quantum modifications imply a bounce of the collapsing field, which for large enough mass is covered by an inner and outer evaporating horizon (dashed). A single matching suffices only until the inner horizon disappears. The dot-dash curves correspond to the subsequent evolution which is not determined in our model.

Hawking radiation, which may be included effectively in the outside matter content, but also from violations of energy conditions around the bounce, which may lead to effective outgoing negative energy.

The model is not able to specify the future of the system after it re-emerges out of the horizon. Equation (21) shows that  $dv/dt$  diverges if and only if  $\dot{a} > 0$  and the matching surface becomes trapped. Thus, we can describe the collapse with a single matching until a horizon disappears, at which point the interior  $t$  ceases to be a good coordinate. One has to continue with a second matched region to analyze the future of the system, but this is beyond the scope of our model.

The qualitative picture that emerges from our toy model is thus the following:

- We do obtain black holes, i.e., dynamical horizons, for large masses, but they contain a bounce of the infalling matter rather than a singularity. For large mass, violations of energy conditions are initially small and the evaporation takes a long time, so that there are only small deviations from classical results.
- For small enough mass however, black holes do not form; horizons do not develop during collapse and the bounce is uncovered. The critical threshold scale for horizon formation is given by the turning point in the  $|\dot{a}|$  curve. By Eqs. (8) and (14), the critical scale is

$$a_{\text{crit}} = 0.987\ell_* = 0.276\sqrt{j}\ell_p. \quad (25)$$

The corresponding threshold mass is  $m_{\text{crit}} = m_M + m_\phi(a_{\text{crit}})$ , but we are unable to compute this mass because the exterior dynamics remains undetermined.

Our estimates may be strongly influenced by the simplifications, in particular a homogeneous interior, we are forced to impose on the problem. However, the qualitative features should be robust, and can provide guidance for further more general analysis. In particular, they mean that there could be *lower bounds on the masses of black holes that form by gravitational collapse*. This could rule out primordial black holes below the threshold mass, and thus modify estimates of Hawking radiation effects from very small black holes. More speculative is an extension to highly non-spherical situations such as particle collisions. If LQG effects can in future be shown to encode some of the non-perturbative aspects of string theory, then our results may have implications for the production of black holes in colliders, as predicted in brane-world gravity [17]. The black hole horizon threshold would be a multiple not of  $\ell_p$ , but of the higher-dimensional Planck scale, which could be as low as  $O(\text{TeV})$ . This would mean that higher collision energies are needed for horizon formation, so that black hole production could be significantly reduced.

**Acknowledgments:** RM is supported by PPARC. PS is supported by Eberly research funds of Penn State and NSF grant PHY-00-90091.

- 
- [1] See, e.g., A. Ashtekar and J. Lewandowski, *Class. Quantum Grav.* **21**, R53 (2004).
  - [2] C. Rovelli and L. Smolin, *Nucl. Phys.* **B442**, 593 (1995); A. Ashtekar and J. Lewandowski, *Class. Quantum Grav.* **14**, A55 (1997).
  - [3] A. Ashtekar et al., *Phys. Rev. Lett.* **80**, 904 (1998).
  - [4] M. Bojowald, *Phys. Rev. Lett.* **86**, 5227 (2001).
  - [5] M. Bojowald, *Class. Quantum Grav.* **21**, 3733 (2004); V. Husain and O. Winkler, gr-qc/0410125; M. Bojowald, *Phys. Rev. Lett.* (to appear), gr-qc/0506128.
  - [6] M. Bojowald, P. Singh, and A. Skirzewski, *Phys. Rev. D* **70**, 124022 (2004).
  - [7] S. W. Hawking and G. F. R. Ellis, *The Large Scale Structure of Space-Time* (Cambridge University Press, 1973), p. 135.
  - [8] T. Thiemann, *Class. Quantum Grav.* **15**, 1281 (1998); M. Bojowald, *Class. Quantum Grav.* **19**, 2717 (2002).
  - [9] M. Bojowald, *Class. Quantum Grav.* **19**, 5113 (2002).
  - [10] P. Singh and A. Toporensky, *Phys. Rev. D* **69**, 104008.
  - [11] M. Bojowald, *Phys. Rev. Lett.* **89**, 261301 (2002).
  - [12] G. Date and G. Hossain, *Phys. Rev. Lett.* **94**, 011302 (2005).
  - [13] M. Bojowald, *Class. Quant. Grav.* **18**, L109 (2001).
  - [14] M. Bojowald and K. Vandersloot, *Phys. Rev. D* **67**, 124023 (2003).
  - [15] P. S. Joshi and I. H. Dwivedi, *Class. Quant. Grav.* **16**, 41 (1999); A. Wang and Y. Wu, *Gen. Rel. Grav.* **31**, 107 (1999).
  - [16] S. Hayward, *Phys. Rev. D* **49**, 6467 (1994); A. Ashtekar

- and B. Krishnan, Phys. Rev. Lett. **89**, 261101 (2002).  
[17] D. M. Eardley and S. B. Giddings, Phys. Rev. **D66**, 044011 (2002).