

**GL Brown Lecture**

**Environmental extremes: origins, consequences and amelioration in humans.**

M. J. Tipton

Extreme Environments Laboratory  
Department of Sport & Exercise Science  
University of Portsmouth  
Portsmouth PO1 2ER  
UK

[michael.tipton@port.ac.uk](mailto:michael.tipton@port.ac.uk)



This is the peer reviewed version of the following article: doi: 10.1113/EP085362, w

## Abstract

Professor Sir George Lindor Brown (1903–1971) is known for his pioneering research into cholinergic neuromuscular transmission. However, during WWII he worked in hyperbaric physiology and his research into underwater physiology greatly improved the safety of divers. It is perhaps fitting therefore that this review, which accompanies the Physiological Society's GL Brown lecture for 2015<sup>1</sup>, explores the impact and mitigation of the environmental stresses which, to varying extents, have shaped our past, threaten our present and inform our future. From a whole-body, integrative perspective, this review examines our current understanding of micro-gravity, hypo- and hyperbaria, heat, cold air and cold water as both individual as well as combined stresses. Consideration is given to ways of mitigating the threat posed by environmental extremes, including the differing extents to which humans can demonstrate adaptation to them. Finally, recommendations for further study are suggested that might result in both direct and indirect insights.

---

Since the iconic picture, "Earthrise", was taken from Apollo 8 as it orbited the moon in 1968 we have considered our world a "blue planet", an oasis amongst otherwise inhospitable and indifferent spheres. However, the idea of an Eden-like home that we have evolved to be in harmony with, that nurtures us as we care for it, is some way from the truth. On one side of the ecological coin we have the damage we do to our world by our existence. On the other side we have the fact that 70 % of our planet is covered in water, with an average ocean depth of 4,000 m. Light penetrates up to about 80 m into water, so most of our planet is in permanent darkness; the rest of it is in intermittent darkness with day and night. Only 15 % of the surface of the Earth is not water, desert, ice or mountain. For humans, a sub-tropical, 1-G, low altitude, air-breathing homeothermic animal, most of our home represents an "extreme environment", defined as a place where it is difficult to survive.

Whilst we can, to differing extents, demonstrate physiological adaptation to heat, cold and altitude; if constrained by these capacities we would still inhabit a very small portion of our planet, close to our equatorial origins. The 3–4 million-year evolution of humans has been influenced and determined by the environment. The migration from our equatorial origins was primarily determined by the ability to adapt behaviourally; it was this, underpinned by intellect and invention, that allowed us to protect ourselves, recreate our desired macro and microclimates, and thereby colonise most of the rest of the planet, and even play one-handed golf on the Moon (Alan Shepard, NASA astronaut, 1971). So it is that, wherever in the world you take the measurement, if resting humans report being "comfortable" they will have a deep body temperature of 37 °C and mean skin temperature of 33 °C, just as they would have had living naked in their ancestral home with its air temperature of 28 °C.

---

<sup>1</sup> The GL Brown lecture is given in memory of the distinguished physiologist Sir George Lindor Brown (1903-1971). It is aimed at a younger audience with a view to stimulating an interest in physiology. M. Tipton gave the GL Brown lecture in 2015 at (chronological order) the Universities of Keele, Cambridge, Westminster, Bath, Portsmouth, Trinity College Dublin, Brighton, Sheffield and at the Physiological Society, London.

The technical abilities that have enabled us to inhabit the rest of the planet have also made us dependent on that technology, and thereby limited the extent of our physiological adaptation. Indeed, there is an argument that those stressors that humans have historically been most able to counter, are also the ones to which it is most difficult to identify an adaptation. The oldest building made by homonids was a windbreak constructed by *Australopithecus*  $\approx$  3.25 million years ago. Over a million years ago, *Homo erectus* was building huts from stones, branches and furs. The earliest evidence for the use of fire, a source of light and radiant heat, comes from China (600,000 BC), whilst the oldest clothed body yet discovered was that of a man who died about 35,000 years ago (Tipton et al, 2008). Conversely, those environmental stressors which were more difficult for our ancestors to mitigate (e.g. heat, altitude) can produce significant physiological adaptation with repeated exposure. Our technical abilities also explain why humans do not possess the morphological and anatomical adaptations observed in other species.

The problem comes when things go wrong and we have to fall back on our relatively primitive, unevolved, physiological defences. The consequences of their limitations include: drowning, hypothermia, cold injury, hyperthermia and barotrauma. These pathophysiological threats can arise in natural as well as man-made environments: from deserts to space craft. A simple hierarchy of survival is presented in Table 1, although the threats differ, the ultimate problem is usually hypoxia.

**Table 1.** The hierarchy of survival

| <b>Threat</b>                                       | <b>Survival Time</b> |
|---|----------------------|
| Lack of Oxygen (respiratory/circulatory impairment) | Minutes              |
| Thermal imbalance (hyper/hypothermia)               | Hours                |
| Fluid imbalance (dehydration)                       | Days                 |
| Energy imbalance (starvation)                       | Weeks                |

The lack of precision in the “survival times” in Table 1 is indicative of the large variation in the responses of humans to environmental threats. This variability is caused by a combination of environmental factors (water/air temperature, humidity, pressure *etc.*) and individual phenotypical and genotypical factors (sex, race, health, fitness, acclimatisation status *etc.*). For example, survival time from dehydration differs in a marine ( $\approx$  6 days) compared to a desert environment ( $\approx$  3 days, Golden & Tipton, 2002). The variation is not only apparent between environments and individuals, but also within individuals on different occasions. Thus, an experienced mountaineer, with no previous history of acute mountain sickness, may succumb to it on a subsequent visit to altitude. The reasons for this are not clearly understood; the role of subtle predisposing changes in the physiology of the climber, including asymptomatic infection, have not been elucidated. Additionally, the co-existence of additional stresses may result in a problem that does not exist otherwise; those that are most likely to become hypothermic in a life-raft at sea are also those that are seasick and hungry; motion illness impairs the heat loss defence mechanisms (decreased vasoconstriction [Mekjavic et al, 2001]) and

hypoglycaemia impairs shivering (Gale et al, 1981). Motion illness also decreases vasoconstriction and arterial pressure and this, in turn, decreases G-tolerance (Eiken et al, 2005).

The investigation of the impact of the environment on human physiology has been undertaken for a variety of reasons, including to: i. gain an understanding of the physiology and pathophysiology surrounding a specific threat. ii. prepare and protect those entering extreme environments for occupational or leisure reasons. This applies to groups as diverse as the chemical, baking, canning and energy industries, as well as the military and sports enthusiasts ranging from outdoor pursuits to fencing. iii. obtain insights into other medical conditions.

Starting from a high point, the remainder of this article briefly examines different extreme environments with a view to highlighting these aspects.

### Micro-gravity

One has to travel 6.37 million km into space to reach a point where the gravitational pull from the Earth is one millionth of that at the surface. As spacecraft orbit 200-450 km from the surface astronauts experience apparent weightlessness or “micro-gravity” ( $\mu\text{G}$ ) rather than zero gravity, and orbit is equivalent to free fall. The resulting reduced hydrostatic gradients and loading acting upon the body produce a range of cardiovascular, pulmonary, musculoskeletal, hormonal, vestibular, neurosensory, haematological, thermoregulatory, fluid and electrolyte changes (Lujan & White, 1994; Convertino, 2000; Mekjavic et al, 2005). The time-courses of these alterations vary, with the fluid shifts and cardiac deconditioning happening earlier (days to weeks) and muscle and bone mass changes later (months). These changes are less significant when astronauts are in  $\mu\text{G}$  than when they return to 1G, where the physiological deconditioning they have experienced can become pathological. This has led to many studies examining possible interventions to mitigate the reduction in the anatomical and physiological status caused by exposure to  $\mu\text{G}$ . Because the changes seen mimic those seen with detraining, inactivity and immobilization, most interventions involve careful crew pre-selection and the completion of ingenious forms of exercise in space to try and reduce the deleterious effects of  $\mu\text{G}$  (Nicogossian et al, 1994).

The expense and logistical demands of spaceflight mean other methodologies have been employed to simulate exposure to  $\mu\text{G}$ , these include parabolic flight, immersion, immobilisation and bed rest. It is the last of these methodologies that is most widely used, it also provides an alternative rationale for the study of  $\mu\text{G}$ ; the investigation of physiological aspects of space travel is seldom justified in terms of the future of our species, the need to leave the solar system within the next 4.5 billion years is a powerful argument, but is a little too far outside five year spending plans. Instead, the relevance of such investigations for other conditions such as ageing, bone changes (osteoporosis) in post-menopausal women and extended bed rest in critical care patients is emphasised.


Vico et al (2000) reported a 25% loss of bone density after 6 months of space travel in 11 cosmonauts in the Russian MIR space station. This is somewhat more than the 5-10% seen in post-menopausal women, and occurred in the legs rather than arms, with large inter-individual variation. The time to fully recover was longer than the mission. The cause of the reduction in bone mineral density has been a matter of some debate, with mechano-adaptation following the loss of muscle contractions/strength now being preferred to fluid shifts and changed perfusion pressure gradients as the primary mechanism (Rittweger et al, 2010). This “muscle-bone” hypothesis is supported by the finding that exercise that maintains muscle strength is an effective countermeasure to bone loss (Shackelford et al, 2004).

Bed rest studies modelling the impact of  $\mu$ G can also provide valuable information for acute muscle wasting in critical illness (Puthuchery et al, 2013), with some of the countermeasures being applicable to both  $\mu$ G and the care, maintenance and rehabilitation of the long-term critically ill (Greenleaf, 1997; Parry et al, 2012).

### Hypobaria

Barometric pressure decreases in an approximately exponential function with distance above the earth. One atmosphere is the pressure exerted by the 38 km of air above us and is equal to about 760 mmHg (101.33 kPa; 14.7 psi); because air is compressible half an atmosphere is at an altitude of about 5,500 m.

The primary problem associated with altitude is the fall in barometric pressure and the consequent fall in the partial pressure of oxygen and oxygen availability at the tissues; this problem is compounded when oxygen demand is increased by exercise. The problem is characterised by the S-shape of the oxygen-dissociation curve which means that normally, at sea level, small falls in the partial pressure of oxygen have relatively little impact on the saturation of haemoglobin, however, below a  $PO_2$  of around 50 mmHg (equivalent to an altitude of about 5,486 m) such falls have a more significant impact on %Hb saturation, this represents the highest elevation with permanent residents (just below half an atmosphere, Figure 1).



**Figure 1.** The physical, physiological and pathophysiological consequences of going to altitude (from: McArdle, Katch & Katch [2001] Exercise Physiology 5<sup>th</sup> Edition Lippincott, Williams & Wilkins).

Above 1,500 m maximum aerobic capacity decreases by about 7.7-11 % for every 1000 m increase in altitude up to 6,300 m at which point it declines at a greater rate (Fulco et al, 1998). The symptoms of hypoxia progress with increasing altitude with the major dangers being high altitude cerebral oedema [HACE] and high altitude pulmonary oedema [HAPE]). The pathogenesis of these conditions has been something of a mystery, however there is a mounting body of evidence of a relationship between genotype and mountain sickness; several authors have reported a relationship between the angiotensin-converting enzyme (ACE) genotype and high altitude pulmonary hypertension, and endothelial nitric oxide synthase gene polymorphisms and high altitude pulmonary oedema (Morrell et al, 1999; Aldashev et al, 2002; Droma et al, 2002).

Two further areas of high altitude research with the possibility of illuminating pathological states include studies into the regulation of cerebral blood flow (CBF) at high altitude, and the individual and temporal response to hypoxia. The initial increase in CBF on exposure to high altitude (Severinghaus et al, 1966) and the change in the cerebrovascular response to  $PCO_2$  at high altitude can influence breathing responses (Ainslie & Duffin, 2009). The altered cerebrovascular reactivity to  $CO_2$  and breathing instability seen at high altitude may have implications for unstable breathing patterns in patients with congestive heart failure (Xie et al, 2005) and obstructive sleep apnoea (Burgess et al, 2010). The mechanisms underlying the changes in CBF at high altitude remain to be fully elucidated; 40% of the increase in CBF at 5050 m appears to be related to the balance of  $PaO_2 / PaCO_2$ , and be determined by ventilatory sensitivity. Whilst the other 60% is thought to be due to neuronal and local (e.g. endothelium-derived nitric oxide) factors (Lucas et al, 2011). 7-9 days of acclimatisation leads to normalisation of CBF due to hypoxic-induced increases in ventilation, with a consequently higher  $PaO_2 / PaCO_2$  ratio resulting in less hypoxia-induced dilatation and more hypercapnia-induced constriction in the cerebral circulation (Lucas, 2011).

Another justification for high altitude research has been its relevance for intensive care medicine, this is part of a paradigm that suggests that the physiological and pathophysiological responses to extreme environments may be similar to those seen in critical illness (Grocott et al, 2007). The cellular hypoxia associated with critical illness and the responses to high altitude may have parallels in terms of the progression from acute responses that attempt to overcome tissue hypoxia by increasing oxygen delivery, to longer-term responses that reduce oxygen utilisation by increasing efficiency (Hochachka et al, 1996). Furthermore, a better understanding of the variability observed in the response of individuals to altitude may help explain the variability in the outcome from critical illness founded on hypoxia. For example, the genes identified as assisting high altitude performance might also improve outcome in critical illness: the ACE gene insertion allele improves performance at altitude and lowers mortality from various conditions including acute respiratory distress syndrome (Marshall et al, 2002).

Finally, there remains some debate about the validity of using normobaric hypoxia as a model for hypobaric hypoxia (Beidleman et al, 2014); a recent systematic review and evaluation of 13 studies (Coppel et al, 2015) suggested there were differences between the two in a number of variables including minute ventilation and nitric oxide levels, suggesting the presence of a physiological difference. However, the authors were unable to make a definitive conclusion due to a lack of methodological standardisation between the studies.

### Hyperbaria

Diving has a long history; Greek sponge divers of 2500 years BC are mentioned in Homer's Iliad and Odyssey. In 1500 Leonardo da Vinci designed the first snorkel with fins for hands and feet. In 1650 Von Guericke developed the first air pump which was used by Robert Boyle and led to his first description of decompression sickness with the appearance of bubbles in the eyes of a viper following compression and decompression. Sir Edmund Haley patented the first diving bell in 1690 and the first confirmed submarine battle (US Turtle vs. HMS Eagle) took place in 1776. The first prototype self-contained breathing apparatus (SCUBA) was invented by James in 1825 and developed by Fleus in 1878. In 1873 Smith reported decompression sickness (DCS) in workers leaving their pressurized caisson working on the Brooklyn Bridge, the workers bent forward from sudden joint pain; this resembled the fashionable "Grecian bend" of the tightly corseted New York society women. The term "bends" remains to this day. In 1878 Paul Bert published "*La Pression Barometrique*", suggesting that nitrogen bubbles cause decompression sickness (DCS) and that gradual ascent and recompression help alleviate the condition. This was confirmed by Boycott et al (1908) with the suggestion of staged decompression.

The physiology and pathophysiology associated with the hyperbaric environment are dominated by the physical Laws of Boyle (barotrauma), Dalton (oxygen toxicity, nitrogen narcosis) and Henry (decompression sickness) and, more than any other environment in this review, are inextricably linked to the associated technology. Indeed, the development of new techniques and equipment in diving

usually preceded an understanding of the possible physiological consequences of their use (Brubak & Neuman, 2003).

Perhaps the most important diving-related response to mention within the confines of this review is pulmonary barotrauma resulting in air embolism. This is the second leading cause of death after drowning for those entering the water. The greatest change in lung volume occurs near the surface (Boyle's Law: between 10 m and the surface the pressure is halved so the volume doubles). Having taken a breath from a source of compressed air, breath holding during ascent can lead to rupture of the alveolar membrane and air emboli entering the pulmonary vasculature and systemic circulation where blockage of the pulmonary, coronary or cerebral circulation results in symptoms ranging from confusion and weakness to death. The shallowest depth from which this has occurred is thought to be 1 m (Tipton et al, 1995), and the most unfortunate death occurred in a diver at a fixed shallow water depth who happened to hold his breath whilst concentrating on his camera as the peak and then trough of a wave passed above him, effectively reducing his depth of immersion and causing pulmonary barotrauma (Shilling et al, 1984).


In most regions, the thermal consequences of diving have to be considered. At depths of greater than 180 m a diver requires 1.2-1.3 kW of external heating to maintain thermal balance. At depths greater than 100 m heating the inspired gas is mandatory as heat loss from the respiratory passages changes from primarily evaporative to convective (due to increased gas density) and becomes a major route of heat loss. At 30 atmospheres, divers breathing gas at a temperature of 4°C will lose all of their metabolic heat via the respiratory tract, independent of the insulation provided at the skin surface. Respiratory heat loss can be reduced by breathing less dense gas mixes, but even when using helium/oxygen mixes ("heliox") at depths of 250 m hypothermia can occur in less than 20 minutes when breathing un-heated gas, despite normal skin temperatures (Hoke et al, 1976).

Hyperbaric oxygen therapy has been recommended for a wide range of conditions. However, in some cases the definitive experimentation underpinning these treatments is either missing, or lacking the necessary control of normobaric oxygen (Kindwall & Whelan, 2008).

### Heat

Despite being a "tropical animal", designed to live naked in air temperatures around 28 °C and possessing powerful effector mechanisms for heat loss, humans can suffer heat illness including heat stroke and death (Figure 2). This usually occurs when individuals choose, or are required, to exercise in hot and/or humid environments, or when they exercise when wearing thermally inappropriate clothing (e.g. chemical, biological, radiological or nuclear protective clothing). If the body were prevented from losing any of the heat it produced, a fatal level of heat storage would be reached in about four hours at rest or 25 minutes with moderate exercise. Some cases of exertional heat illness occur as a result of malignant hyperthermia, an inherited pharmacogenetic disorder of skeletal muscle characterised by an elevated calcium release from the skeletal muscle sarcoplasmic reticulum. This

condition has been associated with pathogenic variants in the RYR1 and CACNA1S genes (Fischer et al, 2015).



**Figure 2.** The physiological and pathophysiological signs and symptoms of increasing heat illness.

The problems caused by heat result from decreased circulating blood volume, alterations in regional blood flow, increased blood viscosity and the direct effect of temperature on the respiratory centres and proteins. Lower levels of heat stress, particularly when compounded by dehydration, can cause physiological changes, including cardiovascular insufficiency, that impair exercise performance (Tipton, 2015, Table 2).


**Table 2.** Physiological changes when hot compared to cool (Rowell, 1986; Tipton, 2015).

| Variable                            | Consequences   |
|-------------------------------------|--|
| Central venous pressure – reduced   | Submaximal exercise performance can be affected e.g. marathon performance declines by about 1 minute for each 1°C increase in air temperature above 15 °C. |
| Central blood volume - reduced      |  |
| Heart rate – higher                 |  |
| Stroke volume – reduced/unchanged   |  |
| Cardiac output - reduced            |  |
| Arterial blood pressure - reduced   | Compromised muscle and liver blood flow results in an earlier onset of anaerobic metabolism and blood lactate accumulation                                 |
| Oxygen consumption – same           |  |
| a-vO <sub>2</sub> difference - same |  |
| Splanchnic blood flow – reduced     |  |
| Splanchnic blood volume – reduced   | Muscle glycogen utilisation is increased, and fatigue occurs earlier during prolonged moderate exercise in the heat  |
| Renal blood flow – reduced          |  |
| Muscle blood flow - reduced         |  |
| Skin blood flow - increased         |  |
| Skin blood volume - increased       |  |

Various options for the active cooling of hyperthermic individuals exist. Of these, whole body fanning and hand immersion in cold water, which use the physiological responses of the body (sweating and peripheral blood flow) are quick and efficient (Barwood et al, 2009). If the victim is suffering heat stroke, with an absence of sweating and compromised peripheral blood flow, conductive cooling by immersion in cool water is required.

### Cold Air

Provided adequate and appropriate (wind/waterproof) thermal protection is worn and metabolism is maintained through exercise, hypothermia is rarely a problem in even very cold environments. The problem comes when injury or exhaustion intervene and limit the metabolic production of heat. The signs and symptoms of hypothermia are presented in Figure 3, hypothermia decreases cellular metabolism, blood flow and neural activation.



**Figure 3.** The physiological and pathophysiological signs and symptoms of hypothermia. \* Time varies due to individual and environmental factors.


A more likely risk in very cold environments is cold injury of the peripheral tissue. “Frostbite” is a well-known and understood condition, with the pathology resulting from intracellular freezing during rapid cooling and direct mechanical disruption. During slower cooling, extracellular water crystallisation causes osmotic outflow leading to intracellular dehydration, damage to capillary wall endothelium and cell death. As a consequence there is a local release of inflammatory mediators (such as prostaglandin F<sub>2</sub> [PGF<sub>2</sub>], thromboxane A<sub>2</sub> [TA<sub>2</sub>], and oxygen radicals), a reduction in plasma volume, oedema, reduced local blood flow, capillary sludging, platelet aggregation, vessel thrombosis, local hypoxia and gangrene (Reamy, 1998; Handford et al, 2014).

A much less well known and understood condition is non-freezing cold injury (NFCI) in which the tissues are damaged by long-term exposure to cold and often wet environments, without tissue freezing. The long-term sequelae of NFCI include cold sensitivity (exaggerated cold-induced vasoconstriction), hyperhidrosis and intractable pain in the injured area. The symptomatology of NFCI suggests neuropathy and vascular endothelial damage associated with impaired nitric oxide (NO) dependent endothelial function, tipping the balance in favour of Rho kinase activity leading to increased vasoconstriction (Wettsschureck & Offermanns, 2002). Some support for this mechanism comes from the finding that the impairment in the vasodilator response seen in individuals with cold-sensitivity can be overcome by the use of exogenous sublingual glyceryl trinitrate, an endothelial-

independent NO donor (Hope et al, 2014). A better understanding of the pathological mechanism associated with NFCI may shed light on the microvascular disorders associated with other conditions such as diabetes and Raynaud's.

### Cold Water

Despite an historic pre-occupation with hypothermia as the primary cause of death on immersion in cold water, a significant amount of anecdotal, statistical and experimental evidence points to the initial minutes of immersion being the most hazardous. It takes at least 30 minutes for an adult to become hypothermic in cold water, yet about 55% of the annual open-water deaths in the UK occur within a few minutes and within 3 m of a safe refuge. The initial inspiratory “gasp” and uncontrollable hyperventilation on immersion in cold water (“Cold shock”, Tipton, 1989), evoked by the dynamic response of the peripheral cold receptors with rapid skin cooling, can be the pre-cursors to drowning. Cold shock is followed by cooling of the superficial nerves and muscle, particularly in the limbs resulting in swim failure and, in the absence of airway protection, drowning (Tipton et al, 2000). Cold shock and neuromuscular incapacitation can both occur without deep body temperature falling to hypothermic levels (<35 °C). When hypothermia does eventually occur, and unconsciousness is lost (Figure 3), drowning again intervenes unless the airway is supported by a lifejacket (Figure 4).




**Figure 4.** The “Physiological Pathways to Drowning” following immersion / submersion in cold water, with possible interventions for partial mitigation (dashed boxes). IS = Immersion suit; LJ = lifejacket; EBA = emergency breathing aid.

Drowning remains the third most common cause of unintentional injury death worldwide (7% of all injury-related deaths), with a conservatively estimated 372,000 (1020 per day) drowning deaths

worldwide in 2012 (WHO, 2014). It is a “disease of youth”; 64 % of those that drown are under 30 years of age, 25 % are under 5 years old.

Cardiac problems on immersion have probably been under-estimated due to the fact that electrical disturbances of the heart which result in agonal gasps and aspiration of water (Figure 4) are likely to be considered as drowning deaths. Also, young, fit and healthy volunteers are tested in the laboratory. Even then head out-immersion results in arrhythmia and dysrhythmia in about 1-3 % of immersions (Datta & Tipton, 2006). This figure increases to over 80 % if maximum breath holding and submersion are undertaken; coincidental activation of the cold shock response (sympathetic activation) and “diving response” with face immersions (vagal activation) causes “Autonomic conflict” (Shattock & Tipton, 2012; Tipton, 2013). It is thought that the arrhythmias caused by autonomic conflict can, in the presence of predisposing factors, descend into fatal arrhythmias (Figure 5); it is suggested that the same mechanism may initiate sudden cardiac death in other situations (Shattock & Tipton, 2012).




**Figure 5.** Autonomic Conflict – a proposed mechanism for sudden cardiac death (from: Shattock & Tipton [2012]. “Autonomic conflict”: a different way to die on immersion in cold water? *Journal of Physiology*. 590[14], 3219-30).

On rare occasions individuals, particularly children, survive prolonged submersion and make a full recovery. The current “record” is 66 minutes in a 2.5 year old female (Bolte et al, 1988). It has been

suggested that cold water entering the lungs during the two minutes it takes to drown, selectively cools the heart and the brain via the carotid artery and, via this mechanism, the hypometabolic consequence of hypothermia extends the hypoxic survival time of the brain (Tipton & Golden, 2012). The critical water temperature for this mechanism to operate appears to be around 6 °C, and the larger surface area to mass ratio of smaller individuals, which promotes faster supplementary surface cooling, helps explain why those that have survived prolonged submersion have tended to be children or small adults. This information has been communicated to the search and rescue community in the form of a decision making guide for the rescue and resuscitation of submerged individuals (Tipton & Golden, 2012). Thus, an understanding of the physiological and pathophysiological processes associated with immersion helps with the prevention and treatment of these fatalities, sheds light on sudden death in other situations and helps inform search and rescue.

### Adaptation

As mentioned, to a greater or lesser extent humans can, following repeated exposure, adapt physiologically to the environmental stressors described above. These adaptations may take the form of inherited (genotypic) or acquired (phenotypic) variations (Tipton et al, 2008). Adaptation theory (Figure 6) addresses most of the applied questions surrounding adaptation.



**Figure 6.** General Adaptation Theory: showing adaptation threshold, physiological response and decay with time (from: Taylor & Cotter [2006]).

The capacity of any stress to induce an adaptation is a function of its application frequency, duration, intensity and variability. The time-course of adaptation has six general characteristics (Adolph, 1964):

1. The stimulus (forcing function) has to be greater than an “adaptation threshold” to induce adaptation.
2. Variable phase delay (“adaptation latency”) exists before adaptation
3. For each effector organ there is a physiological and possibly a genetically determined maximum.

4. The speed of adaptation varies between stressors and individuals.
5. Optimal adaptation occurs when the cumulative adaptation impulse is maximised, whilst ensuring the physiological strain is tolerable.
6. Cessation of adaptation stimulus results in variable decay.

The most common applied questions concerning adaptation are: “how long does it take?” and “how long does it last?” The traditional answers with regard to heat and altitude are 10-14 days with major benefits retained for about a week, but 75% lost within three weeks (Armstrong and Maresh, 1991; Tipton et al, 2008). More recent evidence suggests that short-term heat acclimation (plasma volume expansion, reduced exercise heart rate, increased forearm perfusion, lower deep body temperature, improved rowing performance in the heat) can be achieved in five days (Garrett et al, 2014); the value of concurrent permissive dehydration in this response is a matter for debate (Neal et al, 2015). There is also recent evidence that heat acclimation can be used as an ergogenic aid for performance in cool conditions (Lorenzo et al, 2010; Corbett et al, 2014).

Some of the adaptations to altitude, heat and cold (water and air) are listed in Table 3. For the reasons suggested above, the picture for adaptation to cold is somewhat more confused than that for heat or altitude.

**Table 3.** Changes in, and consequences of, the responses to different environmental stresses with adaptation (Milledge, 1998; Armstrong and Maresh, 1991; Tipton et al, 2008. HR = heart rate; Q = cardiac output; SV = stroke volume; PV = plasma volume)

| Environmental Stressor | Alteration  | Consequence  |
|------------------------|---|--|
| Altitude >2300 m       |   |  |
| Immediate              | Hyperventilation; Body fluid alkalosis; Increased submax HR & Q; SV & max Q same or slight fall.  | Increased supply of oxygen to tissues; improved efficiency of oxygen utilisation; Improved work capacity |
| Longer-term            | Hyperventilation; Right-shift in O <sub>2</sub> -Hb dissociation curve; Excretion of base (HCO <sub>3</sub> <sup>-</sup> ) by kidneys – decreased alkaline reserve; Increased sympathetic neurohumoral activity; Submax HR remains increased; Submax and max Q fall; SV & PV decrease; Mass and lean body mass fall. Increased: haematocrit, haemoglobin concentration, red blood cell count & 2,3-DPG concentration, skeletal muscle capillarisation, mitochondrial density & aerobic enzyme concentrations. |  |

|                   |  |   |
|-------------------|--|---|
| Heat              | Increased secretion and sensitivity to aldosterone increased ▶ sodium & chloride reabsorption in sweat ducts and renal tubules ▶ sweat sodium reduced ▶ osmotic expansion of plasma volume; Sweating: earlier onset, greater rate and gland recruitment.   | Improved salt balance; Greater cardiovascular stability, less CV strain; Better distribution of cardiac output; Improved cutaneous blood flow; Lower resting and exercising deep body temperature; Lower skin temperature for a given level of exercise; Improved work capacity; Decreased perceived exertion, Carbohydrate sparing.                    |
| Cold – Peripheral | Increased skin temperature; cold-induced vasodilatation; thermal comfort. Decreased pain; cold pressor response; sympathetic activation.   | Warmer extremities ▶ decreased chance of cold injury; Greater thermal comfort in heat.  |
| Cold – Whole body | <u>Water</u> : Cold shock response reduced (decreased ventilatory, cardiac, blood pressure and catecholamine response to cold immersion); Shivering: delayed onset (same sensitivity).<br><u>Air</u> : Delayed onset of shivering; Increased sympathetic activation, plasma noradrenaline, catecholamine sensitivity; Decreased skin temperature and blood pressure response to cold; Greater amount/activation of brown adipose tissue (non-shivering thermogenesis). | Decreased risk of drowning on initial immersion; Faster fall of deep body temperature on exposure to cold (hypothermic adaptation); Decreased fall of deep body temperature on exposure to cold (Insulative adaptation); Increased metabolism in cold (Metabolic adaptation)? Increased thermal comfort in cold; Increased cold-induced vasodilatation. |

There is increasing evidence of a genetic contribution to the adaptation of those living at high altitude in terms of variations in the function of factors such as hypoxic-inducible factor 1 $\alpha$  (Suzuki et al, 2003) and the endothelial nitric oxide synthase gene (Droma et al, 2006; Simonson et al, 2010). Climate also appears to have been an important selective factor acting on genes responsible for common metabolic disorders (Hancock et al, 2008).

The methodologies used to induce adaptation have been varied and thereby a source of some confusion. In general two techniques prevail, repeated exposure to a constant stimulus or repeated achievement of a constant physiological strain (e.g. deep body temperature, heart rate, level of exertion). The second approach maintains the stimulus to adapt and provides more effective

physiological adaptation (Taylor, 2014) but, at least in terms of the heat, provided the method employed provides a threshold stimulus, it does not appear to make a difference to the improvement in thermo-tolerance as represented by leukocyte hsp72 mRNA (Gibson et al, 2015). The relationship between physiological adaptation and thermo-tolerance requires further investigation.

### Multi-stressor studies

A weakness in the field is the relative paucity of studies investigating combined environmental stresses, despite their prevalence in nature. Excluding heat and humidity, which are usually regarded as a single entity, between 1948 and 2012 there were only 43 published human studies involving combined stresses (Tipton, 2012). Whilst recent work has begun to investigate the combination of bed rest and hypoxia, and cold and hypoxia (e.g. McDonnell et al, 2014; Keramidis et al, 2014), it is an area that is worthy of more attention, as is the associated field of “cross-adaptation”. Lunt et al (2012) reported that cold habituation reduced the catecholamine response and number and severity of symptoms on exposure to hypoxia. These data suggest a generic autonomic cross-adaptive effect between cold habituation and exposure to acute hypoxia in humans.

### Conclusion

The investigation of the impact of extreme environments on human physiology has historically been justified in terms of the pathological consequences of such exposures. As areas like the Arctic and other hazardous and remote areas are considered for occupational and leisure use, this justification remains extant. In addition, and increasingly, the responses of fit and healthy individuals to extreme environmental stress are being used to gain insights into other pathological conditions. This appears to be a helpful approach, but more time is required to evaluate its overall utility. In the future, a better understanding of the metabolomic and biomolecular basis of responses and their variation in extreme environments, and the influence of multiple concurrent environmental stressors and their cross-adaptation, should help demonstrate the benefit of extreme environmental research for other conditions and processes.

### References

Adolph EF (1964). Perspectives of adaptation: some general properties. In: Dill DB, Adolph EF (Eds) Handbook of physiology. Section 4. Adaptation to the environment. American Physiological Society, Washington, DC, 27-35.

Aldashev AA, Sarybaev AS, Sydykov AS, Kalmyrzaev BB, Kim EV, Mamanova LB, Maripov R, Kojonazarov BK, Mirrakhimov MM, Wilkins MR & Morrell NW (2002). Characterization of high-altitude pulmonary hypertension in the Kyrgyz: association with angiotensin-converting enzyme genotype. *Am J Respir Crit Care Med* 166, 1396-1402.

Ainslie PN & Duffin J (2009). Integration of cerebrovascular CO<sub>2</sub> reactivity and chemoreflex control of breathing: mechanisms of regulation, measurement and interpretation. *Am J Physiol Regul Integr Comp Physiol* 296, R1473-R1495.

Armstrong LE & Maresh CM (1991). The induction and decay of heat acclimatisation in trained athletes. *Sports Med* 12(5), 302-12.

Barwood MJ, Davey S, House JR & Tipton MJ (2009). Post-exercise cooling techniques in hot, humid conditions. *Eur J Appl Physiol* 107(4), 385-96.

Beidleman BA, Fulco CS, Staab JE, Andrew SP & Muza SR (2014). Cycling performance decrement is greater in hypobaric versus normobaric hypoxia. *Extreme Physiology & Medicine* 3, 8.

Bert P (1878). *La pression barometrique*. Paris, Masson.

Bolte RG, Black PG, Bowers RS, Thorne JK & Corneli HM (1998). The use of extracorporeal rewarming in a child submerged for 66 minutes. *JAMA* 260, 377-9.

Boycott AE, Damant GCC & Haldane J (1908). The prevention of compressed air illness. *J Hyg Camb* 8, 342-443.

Brubak AO & Neuman TS (2003). *Bennett & Elliott's Physiology and Medicine of diving* 5<sup>th</sup> Edition. Saunders, Elsevier Science, London.

Burgess KR, Fan J-L, Peebles KC, Thomas KN, Lucas SJE, Lucas RAI, Dawson A, Swart M, Shepherd K & Ainslie PN (2010). Exacerbation of obstructive sleep apnea by oral indomethacin. *Chest* 137, 707-710.

Convertino VA (2000). Effects of microgravity on exercise performance. In: Garrett WE & Kirkendall DT (Eds) *Exercise and sports science*. Philadelphia: Lippincott Williams & Wilkins.

Coppel J, Hennis P, Gilbert-Kawai E & Grocott MPW (2015). The physiological effects of hypobaric hypoxia versus normobaric hypoxia: a systematic review of crossover trials. *Extreme Physiology & Medicine* 4, 2.

Corbett J, Neal RA, Lunt HC & Tipton MJ (2014). Adaptation to heat and exercise performance under cooler conditions: a new hot topic. *Sports Med* 44(10), 1323-1331.

Droma Y, Hanaoka M, Ota M, Katsuyama Y, Koizumi T, Fujimoto K, Kobayashi T, Kubo K (2002). Positive association of the endothelial nitric oxide synthase gene polymorphisms with high-altitude pulmonary edema. *Circulation* 106, 826-830.

Droma Y, Hanaoka M, Basnyat B, Arjyal A, Neupane P, Pandit A, Sharma D, Miwa N, Ito M, Katsuyama Y, Ota M & Kubo K (2006). Genetic contribution of the endothelial nitric oxide synthase gene to high altitude adaptation in sherpas. *High Alt Med Biol.* 7(3), 209–220.

Datta A & Tipton MJ (2006). Respiratory responses to cold water immersion: neural pathways, interactions and clinical consequences. *J Appl Physiol* 100(6), 2057-2064.

Eiken O, Tipton MJ, Kölegård R, Lindborg B & Mekjavic IB (2005). Motion sickness decreases arterial pressure and therefore acceleration tolerance. *Aviat Space Environ Med* 76(6), 541-546.

Fischer D, Shaw MA, Fisher NA, Carr IM, Gupta PK, Watkins EJ, Roiz de Sa D, Kim JH & Hopkins PM (2015). Next-generation sequencing of RYR1 and CACNA1S in malignant hyperthermia and exertional heat illness. *Anesthesiology* 122(5): 1033-1046.

Fulco CS, Rock PB & Cymerman A (1998) Maximal and submaximal exercise performance at altitude. *Aviat Space Environ Med* 69(80): 793-801.

Gale EAM, Bennett T, Green HJ & MacDonald IA (1981). Hypoglycaemia, hypothermia and shivering in man. *Clinical Science* 61(4), 463-469.

Garrett AT, Goosens NG, Rehrer NJ, Patterson MJ, Harrison J, Sammut I, Cotter JD. Short-term heat acclimation is effective and may be enhanced rather impaired by dehydration. *Am J Hum Biol* 26(3), 311-320.

Gibson OR, Mee JA, Taylor L, Tuttle JA, Watt PW & Maxwell NS (2015). Isothermic and fixed-intensity heat acclimation method elicit equal increases in Hsp72 mRNA. *Scand J Med Sci Sports Suppl* 1, 259-268.

Golden FStC & Tipton MJ (2002). *Essentials of sea survival*. Human Kinetics, Illinois.

Greenleaf JE (1997). Intensive exercise training during bed rest attenuates deconditioning. *Med Sci Sports Exerc* 29(2), 207-215.

Grocott M, Montgomery H & Vercueil A (2007). High-altitude physiology and pathophysiology: implications and relevance for intensive care medicine. *Critical Care* 11, 203.

Hancock AM, Witonsky DB, Gordon AS, Eshel G, Pritchard JK, Coop G, Di Rienzo A (2008). Adaptations to climate in candidate genes for common metabolic disorders. *PLoS Genetics* 4(2), e32.

Handford C, Buxton P, Russell K, Imray CEA, McIntosh SE, Freer AC, Imray CHE (2014) Frostbite: a practical approach to hospital management. *Extreme Physiology & Medicine* 3, 7.

Hochachka PW, Buck LT, Doll CJ & Land SC (1996). Unifying theory of hypoxia tolerance: molecular/metabolic defense and rescue mechanisms for surviving oxygen lack. *Proc Natl Acad Sci USA* 93, 9493-9498.

Hoke B, Jackson DL, Alexander JM & Flynn ET (1976). Respiratory heat loss and pulmonary function during cold gas breathing at high pressures. In: *Underwater Physiology V. Proceedings of the fifth symposium on underwater physiology* edited by Lambertsen CJ. Fed Am Soc Exp Biol, Bethesda, USA.

Hope K, Eglin C, Golden F & Tipton MJ (2014). Sublingual glyceryl trinitrate and the peripheral thermal responses in normal and cold-sensitive individuals. *Microvascular Res* 91, 84-89.

Keramidas ME, Kolegard R, Mekjavic IB & Eiken O (2014). Acute effects of normobaric hypoxia on hand-temperature responses during and after local cold stress. *Eur J Appl Physiol* 15(2), 183-191.

Kindwall EP & Whelan HT (2008). *Hyperbaric medicine and practice – 3<sup>rd</sup> Edition*. Best Publishing Company, Flagstaff, USA.

Lorenzo S, Halliwill JR, Sawka MN, Minson CT (2010). Heat acclimation improves exercise performance. *J Appl Physiol* 109, 1140-1147.

Lucas SJE, Burgess KR, Thomas KN, Donnelly J, Peebles KC, Lucas RAI, Fan J-L, Basnyat R, Cotter JD & Ainslie PN (2011). Alterations in cerebral blood flow and cerebrovascular reactivity during 14 days at 5050 m. *J Physiol* 589, 741-753.

Lujan BF & White RJ (1994). *Human physiology in space*. NASA.

Lunt H, Barwood MJ, Corbett J & Tipton MJ (2010). "Cross-adaptation": habituation to short repeated cold-water immersions affects the response to acute hypoxia in humans. *Journal of Physiology* 588(18), 3605-3613.

Marshall RP, Webb S, Bellingham GJ, Montgomery HE, Chaudhari B, McAnulty RJ, Humphries SE, Hill MR & Laurent GJ (2002). Angiotensin converting enzyme insertion/deletion polymorphism is associated with susceptibility and outcome in acute respiratory distress syndrome. *Am J Resp Crit Care Med* 166(5), 646-650.

McArdle WD, Katch FI & Katch VL (2001). *Exercise physiology* 5<sup>th</sup> edition. Lippincott, Williams & Wilkins, USA.

McDonnell AC, Eiken O, Mekjavic PJ & Mekjavic IB (2014). Circadian rhythm of peripheral perfusion during 10-day hypoxic confinement and bed rest. *Eur J Appl Physiol* 114(10), 2093-2104.

Mekjavic IB, Golja P, Tipton MJ & Eiken O (2005). Human thermoregulatory function during exercise and immersion after 35 days horizontal bed rest. *Eur J Appl Physiol* 95, 163-171

Mekjavic IB, Tipton MJ, Gennser M & Eiken O (2001). Motion sickness potentiates core cooling during immersion in humans. *Journal of Physiology* 1:535.2, 619-623.

Milledge JS (1998). Altitude. In: *Oxford Textbook of Sports Medicine* 2<sup>nd</sup> Edition. Edited by: Harries M, Williams C, Stanish WD & Micheli LJ. Oxford University Press, London.

Morrell NW, Sarybaev AS, Alikhan A, Mirrakhimov MM & Aldashev AA (1999). ACE genotype and risk of high altitude pulmonary hypertension in Kyrgyz highlanders. *Lancet* 353, 814.

Neal RA, Corbett J, Massey HC & Tipton MJ (2015). Effect of short-term heat acclimation with permissive dehydration on thermoregulation and temperate exercise performance. *Scand J Med Sci Sports* doi: 10.1111/sms12526.

Nicogossian AE, Leach Huntoon C & Pool SL (1994). *Space physiology and medicine*. Philadelphia: Lea Febiger.

Parry SM, Berney S, Koopman R, Bryant A, El-Ansary D, Puthuchery Z, Hart N, Warrillow S & Denehy L (2012). Early rehabilitation in critical care (eRiCC): functional electrical stimulation with cycling protocol for a randomised control trial. *BMJ Open* 13:2 (5), e001891

Puthuchery ZA, Rawal J, McPhail M, Connolly B, Ratnayake G, Chan P, Hopkinson NS, Phadke R, Dew T, Sidhu PS, Velloso C, Seymour J, Agle CC, Selby A, Limb M, Edwards LM, Smith K, Rowlerson A, Rennie MJ, Moxham J, Harridge SD, Hart N, Montgomery HE (2013). Acute skeletal muscle wasting in critical illness. *JAMA* Oct 16; 310(15), 1591-1600.

Reamy BV (1998). Frostbite: Review and Current Concepts. *J Am Board Fam Pract* 11, 34-40.

Rowell LB (1986) *Human circulation: regulation during physical stress*. Oxford University Press, New York.

Severinghaus JW, Chiodi H, Eger II EI, Brandstater B & Hornbein TF (1966). Cerebral blood flow in man at high altitude: role of cerebrospinal fluid pH in normalization of flow in chronic hypocapnia. *Cir Res* 19, 274-282.

Shattock M & Tipton MJ (2012). "Autonomic conflict": a different way to die on immersion in cold water? *Journal of Physiology*. 590(14), 3219-30.

Shilling CW, Carlston CB, Mathias RA (1984). *The physician's guide to diving medicine*. Plenum Press, New York, p.66.

Simonson TS, Yang Y, Huff CD, Yun H, Qin G, Witherspoon DJ, Bai Z, Lorenzo FR, Xing J, Jorde LB, Prchal JT & Ge R (2010). Genetic evidence for high-altitude adaptation in Tibet. *Science* 329(5987), 72-75.

Suzuki K, Kizaki T, Hitomi Y, Nukita M, Kimoto K, Miyazawa N, Kobayashi K, Ohnuki Y & Ohno H (2003). Genetic variation in hypoxia-inducible factor 1 $\alpha$  and its possible association with high altitude adaptation in Sherpas. *Med Hypotheses* 61(3), 385-389.

Taylor NAS (2014). Human heat adaptation. *Compr Physiol* 4(1), 325-365.

Taylor NAS & Cotter JD (2006). Heat adaptation: guidelines for the optimisation of human performance. *International Sport Med Journal* 7, 33-57.

Tipton MJ (1989). The initial responses to cold-water immersion in man. Editorial Review, *Clinical Science* 77, 581-588.

Tipton MJ (2012). A case for combined environmental stressor studies. *Extreme Physiology & Medicine* 1, 7.

Tipton, MJ (2013). Sudden cardiac death during open water swimming. *British Journal of Sports Medicine*. 48(15), 1134-5.

Tipton MJ, Franks CM, Gennser M & Golden FStC (1999). Immersion death and deterioration in swimming performance in cold water. *The Lancet* Vol 354, 626-9.

Tipton MJ & Golden FStC (2011). Decision-making guide for the search, rescue and resuscitation of submerged (head under) victims. *Resuscitation* 82, 819-824.

Tipton MJ (2015). Extreme Temperature Sport and Exercise Medicine. Chapter 14 in: ABC of Sports Medicine 4<sup>th</sup> Edition. Edited by: Whyte G, Loosemore M & Williams C. BMJ Books, Wiley Balckwell.

Tipton MJ, Pandolf K, Sawka M, Werner J & Taylor NAS (2008). Physiological adaptation to hot and cold environments. Chapter 21 in: Physiological bases of human performance during work and exercise. Editors: Taylor NAS, Herbert Groeller H, Churchill Livingstone, London.

Vico L, Collet P, Guignandon A, Lafage-Proust MH, Thomas T, Rehailla M & Alexandre C (2000). Effects of long-term microgravity exposure on cancellous and cortical weight-bearing bones of cosmonauts. *Lancet* 5; 355(9215), 1607-1611.

Wettschureck N & Offermanns S (2002) Rho/Rho-kinase mediated signalling in physiology and pathophysiology. *J Mol Med (Berl)* 80(10), 629-638.

World Health Organisation (2014). Global report on drowning: preventing a leading killer. World Health Organisation.

Xie A, Skatrud JB, Khayat R, Dempsey JA, Morgan B & Russell D (2005). Cerebrovascular response to carbon dioxide in patients with congestive heart failure. *Am J Respir Crit Care Med* 172, 371-378.