

## 7. Chapter Seven: Reference List

- Acornley, R. M. (1999). Water Temperatures within Spawning Beds in Two Chalk Streams and Implications for Salmonid Egg Development. *Hydrological Processes*, 13, 439-446.
- Acornley, R. M., and Sear, D. A. (1999). Sediment Transport and Siltation of Brown Trout (*Salmo trutta* L.) Spawning Gravels in Chalk Streams. *Hydrological Processes*, 13, 447-458.
- Adams, J. N., and Beschta, R. L. (1980). Gravel Bed Composition in Oregon Coastal Streams. *Canadian Journal of Aquatic Fisheries Sciences*, 37, 1514-1521.
- Ahel, M., Evans, K., Fileman, T., and Mantoura, R. (1992). Determination of atrazine and simazine in estuarine samples by high-resolution gas chromatography and nitrogen selective detection. *Analytica Chimica Acta*, 268(2), 195-204.
- Akkanen, J., Penttinen, S., Haitzer, M. and Kukkonen, J. V. K. (2001). Bioavailability of atrazine, pyrene and benzo[a]pyrene in European river waters. *Chemosphere* 45, 453-462.
- Allenbach, D. M., Sullivan, K. B., and Lydy, M. J. (1999). Higher fluctuating asymmetry as a measure of susceptibility to pesticides in fishes. *Environmental Toxicology and Chemistry*, 18(5), 899-905.
- Almeida, D., Almodovar, A., Nicola, G. G., and Elvira, B. (2008). Fluctuating asymmetry, abnormalities and parasitism as indicators of environmental stress in cultured stocks of goldfish and carp. *Aquaculture*, 279, 120-125.
- Andersen, M. E., Gargas, M. L., Hays, S. M., Clewell, H. J., III, and Paustenbach, D. J. (1997). Estimating minimally effective inducing doses (ED01s) of TCDD in livers of rats and humans using physiologically based pharmacokinetic modeling. *Organohalogen Compounds*, 34, 311-316.
- Argent, D. G., and Flebbe, P. A. (1999). Fine Sediment Effects on Brook Trout Eggs in Laboratory Streams. *Fisheries Research*, 39, 253-262.
- Armstrong, J., Kemp, P., Kennedy, G., Ladle, M., and Milner, N. (2003). Habitat requirements of Atlantic salmon and brown trout in rivers and streams. *Fisheries research*, 62(2), 143-170.
- Arsenault, J.T.M., Fairchild, W.L., MacLatchy, D.L., Burridge, L., Haya, K., Brown, S.B. (2004). Effects of water-borne 4-nonylphenol and 17 $\beta$ -estradiol exposures during parr-smolt transformation on growth and plasma IGF-I of Atlantic salmon (*Salmo salar* L.). *Aquatic Toxicology*. 66, 255-265.

- Arthur, C. L., and Pawliszyn, J. (1990). Solid phase microextraction with thermal desorption using fused silica optical fibers. *Analytical Chemistry*, 62(19), 2145-2148.
- Ateeq, B., Farah, M. A. and Ahmad, W. (2006). Evidence of apoptotic effects of 2,4-D and butchlor on walking catfish, *Clarions batrachus*, by transmission electron microscopy and DNA degradation studies. *Life Sciences*, 78, 977–986.
- Barbee, G. C., Barich, J., Duncan, B., Bickham, J. W., Matson, C. W., Hintze, C. J., Autenrieth, R. L., Zhou, G. D., McDonald, T. J., Cizmas, L., Norton, D., and Donnelly, K. C. (2008). *In situ* biomonitoring of PAH-contaminated sediments using juvenile coho salmon (*Oncorhynchus kisutch*). *Ecotoxicology and environmental safety*, 71(2), 454-464.
- Bardonnet, A. and J. Baglinière. (2000). Freshwater habitat of Atlantic salmon (*Salmo salar*). *Canadian Journal of Fisheries and Aquatic Science*, 57, 497-506.
- Barron, M. G., Carls, M.G., Heintz, R., and Rice, S.D. (2004) Evaluation of fish early life stage toxicity models of chronic embryonic exposures to complex polycyclic aromatic hydrocarbon mixtures. *Journal of Toxicological. Science*, 78, 60-67.
- Barrett, K., McDonald, D., and O'Donnell, M. (2001). Changes in ion content and transport during development of embryonic rainbow trout. *Journal of Fish Biology*, 59(5), 1323-1335.
- Battin, T. J., Kaplan, L. A., Newbold, J. D., and Hendricks, S. P. (2003). A mixing model analysis of stream solute dynamics and the contribution of a hyporheic zone to ecosystem function. *Freshwater biology*, 48(6), 995-1014.
- Beall, E. (1994). Les phases de la reproduction. In: J.C, Gueguen and P. Prouzet. (Eds). *Le Saumon Atlantique*. (pp. 123–140). Brest: Editions de l'Ifremer.
- Beaumont, W.R.C., Scott, L. and Ibbotson, A.T. (2010). Salmon Research Report 2009. Game and Wildlife Conservation Trust, Fordingbridge, Hampshire.
- Belfroid, A. C., van Drunen, M., Beek, M. A., Schrap, S. M., van Gestel, C. A. M., and van Hattum, B. (1998). Relative Risks of Transformation Products of Pesticides for Aquatic Ecosystems. *The Science of the Total Environment*, 222, 167-183.
- Benedetti, M., Martuccio, G., Fattorini, D., Canapa, A., Barucca, M., Nigro, M., and regoli, F. (2007). Oxidative and modulatory effects of trace metals on metabolism of polycyclic aromatic hydrocarbons in the Antarctic fish *Trematomus bernacchii*. *Aquatic Toxicology*, 85(3), 167-175.
- Bervoets, L., and Blust, R. (2003). Metal concentrations in water, sediment and gudgeon (*Gobio gobio*) from a pollution gradient: relationship with fish condition factor. *Environmental Pollution*, 126(1), 9-19.
- Beschta, R. L., and Jackson, W. L. (1979). The Intrusion of Fine Sediment into a Stable Gravel Bed. *Journal of Fisheries Research Board of Canada*, 36, 204-210.

- Beyer, J., Aas, E., Borgenvik, H. K., and Ravn, P. (1998). Bioavailability of PAH in effluent water from an aluminium works evaluated by transplant caging and biliary fluorescence measurements of Atlantic cod (*Gadus morhua* L.). *Marine Environmental Research*, 46, 233–236.
- Bhattacharya, B., Sarkar, S. K., and Mukherjee, N. (2003). Organochlorine pesticide residues in sediments of a tropical mangrove estuary, India: implications for monitoring. *Environment international*, 29(5), 587-592.
- Billard, R. (1992). Reproduction in rainbow trout: sex differentiation, dynamics of gametogenesis, biology and preservation of gametes. *Aquaculture*. 100, 263-298.
- Billiard, S. M., Querbach, K., and Hodson, P. V. (1999). Toxicity of retene to early life stages of two freshwater fish species. *Environmental Toxicology and Chemistry*, 18(9), 2070-2077.
- Birge, W. J., Black, J. A., Westerman, A. G., and Ramey, B. A. (1983). Fish and amphibian embryos--a model system for evaluating teratogenicity. *Fundamental and applied toxicology*, 3(4), 237-242.
- Bjornn, T. C., and Reiser, D.W. (1991). Habitat Requirements of salmonids in streams. in W.R. Meehan (ed). Influences of forest and rangeland management on salmonid fishes and their habitat. (pp. 83-138). Special Publication 19. American Fisheries Society, Bethesda, MD.
- Bonislawska, M., and Winnicki, A. (2000). Duration of embryonic development and S/V (surface area/volume) coefficient in fish eggs. *Archives of Polish Fisheries*, 8(2), 161-169.
- Bosch, C., Olivares, A., Faria, M., Navas, J. M., del Olmo, I., Grimalt, J. O., Piña, B., and Barata, B. (2009). Identification of water soluble and particle bound compounds causing sublethal toxic effects. A field study on sediments affected by a chlor-alkali industry. *Aquatic Toxicology*, 94(1), 16-27.
- Bouk, G. R. and Jacobson, J. (1976). Estimation of salmonid sperm concentration by microhematocrit technique. *Transactions of the American Fisheries Society*. **105**, 534-535.
- Brack, W., Klamer, H. J. C., de Alda, M. L., and Barceló, D. (2007). Effect-directed analysis of key toxicants in European river basins. A review. *Environmental Science and Pollution Research International*, 14(1), 30-38.
- Brännäs, E. (1989). The Use of a Simulated Redd for Incubating Baltic Salmon (*Salmo salar*) and Trout (*Salmo trutta*) Alevins. *Aquaculture*, 83, 261-267.
- Brassard, C., Gill, L., Stavola, A., Lin, J., and Turner, L. (2003). atrazine analysis of risks to endangered and threatened salmon and steelhead. U.S. Environmental Protection Agency Policy and Regulatory Services Branch and Environmental Field Branch Office of Pesticide Programs. July 27.

- Brown, J. N., and Peake, B. M. (2006). Sources of heavy metals and polycyclic aromatic hydrocarbons in urban stormwater runoff. *Science of the Total Environment*, 359(1-3), 145-155.
- Brunke, M., and Gonser, T. (1997). The ecological significance of exchange processes between rivers and groundwater. *Freshwater biology*, 37(1), 1-33.
- Burton, D. T., Turley, S. D., and Peters, G. T. (1994). The acute and chronic toxicity of hexahydro-1, 3, 5-trinitro-1, 3, 5-triazine (RDX) to the fathead minnow (*Pimephales promelas*). *Chemosphere*, 29(3), 567-579.
- Cachot, J., Geffard, O., Augagneur, S., Lacroix, S., Le Menach, K., Peluhet, L., Devier, M. H., Couteau, J., Pottier, D., Budzinski, H. (2006) Evidence of genotoxicity associated to high PAH content of sediments in the upper part of the Seine estuary (Normandy, France), *Aquatic Toxicology*, 79, 257-267.
- Cai, Z., Wang, D., and Ma, W. T. (2004). Gas Chromatography/Ion Trap Mass Spectrometry Applied for the Analysis of Triazine Herbicides in Environmental Waters by an Isotope Dilution Method. *Analytica Chimica Acta*, 503, 263-270.
- Cantrell, S. M., Joy-Schlezing, J., Stegeman, J. J., Tillitt, D. E., and Hannink, M. (1998). Correlation of 2,3,7,8-Tetrachlorodibenzo-*p*-dioxin-induced Apoptotic Cell Death in the Embryonic Vasculature with Embryotoxicity. *Toxicology and Applied Pharmacology*, 148, 24-34.
- Capkin, E., Altinok, I., and Karahan, S. (2006). Water quality and fish size affect toxicity of endosulfan, an organochlorine pesticide, to rainbow trout. *Chemosphere*, 64(10), 1793-800.
- Carling, P. A. (1984). Deposition of Fine and Coarse Sand in an Open-Work Gravel Bed. *Canadian Journal of Fish. Aquatic Sciences*, 41, 263-270.
- Carling, P. A., and McCahon, C. P. (1987). Natural Siltation of Brown Trout (*Salmo trutta* L.) Spawning Gravels During Low-Flow Conditions. *Regulated Streams: Advances in Ecology*, 228-244.
- Carls, M. G., Holland, L., Larsen, M., Collier, T. K., Scholz, N. L., and Incardona, J. P. (2008). Fish embryos are damaged by dissolved PAHs, not oil particles. *Aquatic Toxicology*, 88(2), 121-127.
- Carls, M. G., Rice, S. D., and Hose, J. E. (1999). Sensitivity of fish embryos to weathered crude oil: Part I. Low-level exposure during incubation causes malformations, genetic damage, and mortality in larval Pacific herring (*Clupea pallasii*) *Environmental Toxicology and Chemistry*, 18, 481-493.
- Carls, M. G., and Thedinga, J. F. (2010). Exposure of pink salmon embryos to dissolved polynuclear aromatic hydrocarbons delays development, prolonging vulnerability to mechanical damage. *Marine Environmental Research*, 69(5), 318-325.

Carpenter, S.R., Caraco, N.F., Correll, D.L., Howarth, R.W., Sharpley, A.N. and Smith, V.H. (1998). Non-point pollution of surface waters with phosphorus and nitrogen. *Ecological Applications*, 8(3), 559–568.

Carter, A. (2000). How pesticides get into water-and proposed reduction measures. *Pesticide outlook*, 11(4), 149-156.

Casey, H., Clarke, R., and Smith, S. (1993). Increases in nitrate concentrations in the River Frome (Dorset) Catchment related to changes in land use, fertiliser applications and sewage input. *Chemistry and Ecology*, 8(2), 105-117.

Casey, H., and Smith, S.M. (1994). The effects of watercress growing on chalk headwater streams in Dorset and Hampshire. *Environmental Pollution*, 85(2), 217-28.

Cattaneo, R., Loro, V., Spanevello, R., Silveira, F., Luz, L., Miron, D, Fonseca, M. B., Moraes, B. S., and Clasen, B. (2008). Metabolic and histological parameters of silver catfish (*Rhamdia quelen*) exposed to commercial formulation of 2, 4-dichlorophenoxyacetic acid (2, 4-D) herbicide. *Pesticide Biochemistry and Physiology*, 92(3), 133-137.

Cavanagh, J. E., Burns, K. A., Brunskill, G. J., and Coventry, R. J. (1999). Organochlorine Pesticide Residues in Soils and Sediments of the Herbert and Burdekin River Regions, North Queensland- Implications for Contamination of the Great Barrier Reef. *Marine Pollution Bulletin*, 39, 367-375.

Cave, R.R., Andrews, J.E., Jickells, T. and Coombes, E.G. (2005) A review of sediment contamination by trace metals in the Humber catchment and estuary, and the implications for future estuary water quality. *Estuarine, Coastal and Shelf Science*, 62,547-557.

Chandler, G. T., Shipp, M. R., and Donelan, T. L. (1997). Bioaccumulation, growth and larval settlement effects of sediment-associated polynuclear aromatic hydrocarbons on the estuarine polychaete, *Streblospio benedicti* (Webster). *Journal of experimental marine biology and ecology*, 213(1), 95-110.

Chang, X., Zhai, B., Liu, X., and Wang, M. (2007). Effects of temperature stress and pesticide exposure on fluctuating asymmetry and mortality of *Copera annulata* (Selys)(Odonata: Zygoptera) larvae. *Ecotoxicology and environmental safety*, 67(1), 120-127.

Chapman, G. A. (1969). *Toxicity of pentachlorophenol to trout alevins*. Ph. D. Thesis. Oregon State University, Corvallis, Oregon.

Chapman, D. W. (1988). Critical review of variables used to define effects of fines in redds of large salmonids. *Transactions of the American Fisheries Society*, 117, 1-21.

Chen, S. J., Luo, X. J., Mai, B. X., Sheng, G. Y., Fu, J. M., and Zeng, E. Y. (2006). Distribution and mass inventories of polycyclic aromatic hydrocarbons and organochlorine pesticides in sediments of the Pearl River Estuary and the northern South China Sea. *Environmental science and technology*, 40(3), 709-714.

Cho, D, Youn, S. H. and Rhee, I. H. (2004). Interaction of b-cyclodextrin with lipophilic ring compounds deposited in a sandy soil matrix during flushing. *Korean Journal of Chemical Engineering*, 21(2), 389-397.

Clarke, G. M. (1994). The genetic basis of developmental stability. I. Relationships between stability, heterozygosity, and genomic coadaptation. In: T. A. Markow (Ed). *Developmental Instability: its Origins and Evolutionary Implications* (pp. 17–25). Kluwer, Boston.

Collins, A. L., McGonigle, D. F., Evans, R., and Zhang, Y. (2009). Emerging priorities in the management of diffuse pollution at catchment scale. *International Journal of River Basin Management*, 7(3), 179-185.

Couch, J., Schoor, W., Davis, W., and Courtney, L. (1983). *Effects of Carcinogens, Mutagens, and Teratogens on Nonhuman Species (Aquatic Animals)*.

Cox, J. R. (2002). Precautions in the use and handling of pesticides and chemical reagents. In: I. F. Grant and C. C. D. Tingle (Eds) *Ecological Monitoring Methods for the Assessment of Pesticide Impact in the Tropics* (pp. 89-94). Natural Resources Institute, Kent, UK.

Crisp, D. T. (1970). Input and output of minerals for a small watercress bed fed by chalk water. *Journal of Applied Ecology*, 7, 117–140.

Crisp, D. T, and Carling, P. (1989). Observations on siting, dimensions and structure of salmonid redds. *Journal of Fish Biology*, 34(1), 119-134.

Crisp, D. T. (1988). Prediction, from temperature, of eyeing, hatching and 'swim-up' times for salmonid embryos. *Freshwater Biology*, 19, 41-48.

Crisp, D. T. (1991). Stream channel experiments on downstream movement of recently emerged trout, *Salmo trutta* L., and salmon, *S. salar* L., III. Effects of developmental stage and day and night upon dispersal. *Journal of Fish Biology*, 39, 371-381.

Crisp, D. T. (1993). The environment requirements of salmon and trout in fresh water. *Freshwater Forum*, 3, 176-202.

Crisp, D. T. (1996). Environmental requirements of common riverine european salmonid fish species in fresh water with particular reference to physical and chemical aspects. *Hydrobiologia*, 323, 201-221.

Crisp, D. T. and Hurley, M. A. (1991). Stream channel experiments on downstream movement of recently emerged trout, *Salmo trutta* L., and salmon, *S. salar* L. II. Effects of constant and changing velocities and of day and night upon dispersal rate. *Journal of Fish Biology*, 39, 363-370.

- Crozier, W., and Kennedy, G. (1994). Marine exploitation of Atlantic salmon (*Salmo salar* L.) from the River Bush, Northern Ireland. *Fisheries research*, 19(1-2), 141-155.
- Curry, R. A., and MacNeill, W. S. (2004). Population-level responses to sediment during early life in brook trout. *Journal of Information*, 23(1).
- da Fonseca, M. B., Gluszcak, L., Silveira Moraes, B., de Menezes, C. C., Pretto, A., Tierno, M. A., Zanella, R., Ferreira Gonçalves, F., and Loro, V. L. (2008). The 2, 4-D herbicide effects on acetylcholinesterase activity and metabolic parameters of piava freshwater fish (*Leporinus obtusidens*). *Ecotoxicology and environmental safety*, 69(3), 416-420.
- Dagnac, T., Bristeau, S., Jeannot, R., Mouvet, C., and Baran, N. (2005). Determination of chloroacetanilides, triazines and phenylureas and some of their metabolites in soils by pressurised liquid extraction, GC-MS/MS, LC-MS and LC-MS/MS. *Journal of Chromatography A*, 1067(1-2), 225-233.
- Davey, G. W., Doeg, T. J., and Blyth, J. D. (1987). Changes in Benthic Sediment in the Thomson river, SouthEastern Australia, During Construction of the Thomson Dam. *Regulated Rivers*, 1, 71-84.
- Davis, J. C. (1975). Minimal dissolved oxygen requirements of aquatic life with emphasis on canadian species: a review. *Journal of the Fisheries Research Board of Canada*, 32, 2295-2332.
- Davies, P. E., Cook, L. S. J., and Barton, J. L. (1994). Triazine Herbicide Contamination of Tasmanian Streams: Sources, Concentrations and Effects on Biota. *Australian Journal of Marine and Freshwater Research*, 45, 209-226.
- Daye, P., and Glebe, B. (1984). Fertilization success and sperm motility of Atlantic salmon (*Salmo salar* L.) in acidified water. *Aquaculture*, 43(1-3), 307-312.
- Debowski, P., Dobosz, S., Grudniewska, J., and Kuzminski, H. (2006). Impact of egg size and origin on the course and period of sea trout (*Salmo trutta* L.) hatching. *Archives of Polish Fisheries*, 14(1), 41-51.
- De Flora, S., Vigano, L., D'Agostini, F., Camoirano, A., Bagnasco, M., Bennicelli, C., Melodia, F., and Arillo, A. (1993). Multiple genotoxicity biomarkers in fish exposed in situ to polluted river water. *Mutation Research/Genetic Toxicology*, 319(3), 167-177.
- del Carmen Alvarez, M., and Fuiman, L. A. (2005). Environmental levels of atrazine and its degradation products impair survival skills and growth of red drum larvae. *Aquatic Toxicology*, 74(3), 229-241.
- Drummond Sedgwick, S. (1982). *The salmon handbook*. Andre Deutsch Ltd. London.

- Dudgeon, D., Arthington, A. H., Gessner, M. O., Kawabata, Z. I., Knowler, D. J., Lévêque, C., Naiman, R. J., Prieur-Richard, A. H., Soto, D., and Stiassny, M. L. J. (2006). Freshwater biodiversity: importance, threats, status and conservation challenges. *Biological Reviews*, 81(2), 163-182.
- Dumas, J., Bassenave, J., Jarry, M., Barriere, L., and Glise, S. (2007). Effects of fish farm effluents on egg to fry development and survival of brown trout in artificial redds. *Journal of Fish Biology*, 70(6), 1734-1758.
- Environment Agency. (2005). *Groundwater quality review: Hampshire Avon Chalk*. Bristol: Environment Agency.
- Environment Agency. (2005). *Cleaner coasts healthier seas. The state of the marine environment of England and Wales*. Bristol: Environment Agency.
- Environment Agency. (2005). *State of the environment*. Bristol: Environment Agency.
- Environment Agency. (2006b). *Underground, under threat. The state of groundwater in England and Wales*. Bristol: Environment Agency. [http://publications.environment-agency.gov.uk/pdf/GEHO1006BLMT-e-e.pdf?lang=\\_e&lang=\\_e](http://publications.environment-agency.gov.uk/pdf/GEHO1006BLMT-e-e.pdf?lang=_e&lang=_e)
- Environment Agency. (2007). *The unseen threat to water quality. Diffuse water pollution in England and Wales*. Bristol: Environment Agency.
- Environment Agency. (2008). *Better sea trout and salmon fisheries*. Bristol: Environment Agency.
- Edeline, E., Lambert, P., Rigaud, C., and Elie, P. (2006). Effects of body condition and water temperature on *Anguilla anguilla* glass eel migratory behavior. *Journal of experimental marine biology and ecology*, 331(2), 217-225.
- Einum, S., and Fleming, I. A. (2000). Selection Against Late Emergence and Small Offspring in Atlantic Salmon (*Salmo salar*). *Evolution*, 54, 628-639.
- Einum, S., Hendry, A. P., and Fleming, I. A. (2002). Egg-size evolution in aquatic environments: does oxygen availability constrain size? *Proceedings of the Royal Society of London. Series B: Biological Sciences*, 269(1507), 2325.
- Elliott, J. M., and Hurley, M. A. (1998). Predicting fluctuations in the size of newly emerged sea-trout fry in a Lake District stream. *Journal of Fish Biology*, 53, 1120-1133.
- Ellis, J., Sumner, K., Griffiths, A., Bright, D., and Stevens, J. Population genetic structure of Atlantic salmon, *Salmo salar* L., in the River Tamar, southwest England. *Fisheries Management and Ecology*.
- Eriksson, T., and Eriksson, L. O. (1993). The status of wild and hatchery propagated Swedish salmon stocks after 40 years of hatchery releases in the Baltic rivers. *Fisheries research*, 18(1-2), 147-159.



European Community (1998). Quality of drinking water. [http://europa.eu/legislation\\_summaries/environment/water\\_protection\\_management/12807\\_9\\_en.htm](http://europa.eu/legislation_summaries/environment/water_protection_management/12807_9_en.htm)

Evans, K., Gill, R., and Robotham, P. (1990). The PAH and organic content of sediment particle size fractions. *Water, Air, and Soil Pollution*, 51(1), 13-31.

Evans, K., Gill, R., and Robotham, P. (1990). The source, composition and flux of polycyclic aromatic hydrocarbons in sediments of the River Derwent, Derbyshire, UK. *Water, Air, and Soil Pollution*, 51(1), 1-12.

Everest, F. H., R. L. Beschta, J. C. Scrivener, K. V. Koski, J. R. Sedell, and C. J. Cederholm. (1987). Fine sediment and salmonid production: a paradox. In: E. Salo and T. Cundy (Eds). *Streamside management and forestry and fishery interactions*. (p. 98-142). University of Washington, College of Forest Resources, Contribution 57, Seattle, WA.

Fairchild, W. L., Swansburg, E. O., Arsenault, J. T., and Brown, S. B. (1999). Does An Association Between Pesticide Use and Subsequent Declines in Catch of Atlantic Salmon (*Salmo salar*) Represent a Case of Endocrine Disruption? *Environmental Health Perspectives*, 107, 349-357.

Fatima, M., Mandiki, S.N.M., Douxfils, J., Silvestre, F., Coppe, P., and Kestemont, P. (2007). Combined effects of herbicides on biomarkers reflecting immune-endocrine interactions in goldfish immune and antioxidant effects. *Aquatic Toxicology*, 81, 159-167.

Faulkner, H., Edmonds-Brown, V., and Green, A. (2000). Problems of quality designation in diffusely polluted urban streams--the case of Pymme's Brook, north London. *Environmental Pollution*, 109(1), 91-107.

Fausch, K. D., Lyons, J., Karr, J. R., and Angermeier, P. L. (1990). *Fish communities as indicators of environmental degradation*.

Feist, G. W., Webb, M. A. H., Gundersen, D. T., Foster, E. P., Schreck, C. B., Maule, A. G., and Fitzpatrick, M. S. (2005). Evidence of detrimental effects of environmental contaminants on growth and reproductive physiology of white sturgeon in impounded areas of the Columbia River. *Environmental Health Perspectives*, 113(12), 1675.

Finn, R.N. (2007) The physiology and toxicology of salmonid eggs and larvae in relation to water quality criteria. *Aquatic Toxicology*, 81, 337-354.

Fleming, I. A., Ng, S. (1987). Evaluation of Techniques for Fixing, Preserving, and Measuring Salmon Eggs. *Canadian Journal of Fisheries and Aquatic Sciences*, 44, 1957-1962.

Fletcher, J.S., Pflieger, T.G. Ratsch, H.C.and Hayes, R.. (1996). Potential impact of low levels of chlorsulfuron and other herbicides on growth and yield of non-target plants. *Environmental Toxicology and Chemistry*, 15, 1189-1196.

- Forney, D. R., and D. E. Davis. (1981). Effects of low concentrations of herbicides on submersed aquatic plants. *Weed Science*, 29, 677-685.
- Forrest, S., and Caux, P.Y. (1989). Pesticides in tributaries of the St Lawrence River 1987-1988. Program report. Environment Canada.
- Frank, R., Braun, H. E., and Ripley, B. D. (1990). Residues of insecticides and fungicides on Ontario-grown vegetables 1986-1988. *Food Additives and Contaminants*, 7, 545-554.
- Frenzilli, G., Nigro, M., Lyons, B.P. (2008). The Comet assay for the evaluation of genotoxic impact in aquatic environments. *Mutation Research/Reviews in Mutation Research*, 681(1), 80-92.
- Frenzilli, G., Scarcelli, V., Del Barga, I., Nigro, N., Förlin, L., Bolognesi, C., and Sturve, J. (2004). *Mutation Research*, 552 (1-2), 187-195.
- Friberg, N., Lindstrøm, M., Kronvang, B., and Larsen, S. E. (2003). Macroinvertebrate/sediment relationships along a pesticide gradient in Danish streams. *Hydrobiologia*, 494(1), 103-110.
- Friedland, K.D., Hansen, L.P. and Dunkley, D.A. (1998). Marine temperatures experienced by postsmolts and the survival of Atlantic salmon, *Salmo salar* L., in the North sea area. *Fisheries Oceanography*, 7, 22-34.
- Frost, W. E., and Brown, M. E. (1967). *The Trout*. Collins: London.
- Fudge, T. S., Wautier, K.G., Evans, R.E., and Palace, V. P. (2008). Effect of different levels of fine-sediment loading on the escapement success of rainbow trout fry from artificial redds. *North American Journal of Fisheries Management*. 28, 758-765.
- Fyhn, H. J. and Serigstad, B. (1987). Free amino acids as energy substrate in developing eggs and larvae of the cod *Gadus morhua*, *Marine Biology*, 96, 335-341.
- Galbraith, R. V., MacIsaac, E. A., Macdonald, J. S., and Farrell, A. P. (2006). The effect of suspended sediment on fertilization success in sockeye (*Oncorhynchus nerka*) and coho (*Oncorhynchus kisutch*) salmon. *Canadian Journal of Fisheries and Aquatic Sciences*, 63(11), 2487-2494.
- Gamble, D.S., Khan, .S.U. and Tee, Q.S. (1983). Atrazine hydrolysis: Part 1. Proton catalysis at 25°C. *Pesticide Science*, 14, 537-545.
- Gao, J. P., Maguhn, J., Spitzauer, P., and Kettrup, A. (1997). Distribution of pesticides in the sediment of the small teufelsweiher pond (Southern Germany). *Water Research*, 31, 2811-2819.
- Gavin, M., Brooke, D., Howe, P., and Dodson, S. (1996). Environmental Hazard Assessment: Naphthalene. TSD/27, Chemicals and Biotechnology Division, Department of Environment, Garston, Watford.

- Geffard, O., Geffard, A., His, E., and Budzinski, H. (2003). Assessment of the bioavailability and toxicity of sediment-associated polycyclic aromatic hydrocarbons and heavy metals applied to *Crassostrea gigas* embryos and larvae. *Marine Pollution Bulletin*, 46(4), 481-490.
- Geist, D. R., and Dauble, D. D. (1998). Redd site selection and spawning habitat use by fall chinook salmon: the importance of geomorphic features in large rivers. *Environmental Management*, 22, 655-669.
- Geist, D. R., Hanrahna, T. P., Arntzen, E. V., McMichael, G. A., Murray, C. J., and Chien, Y. (2002). Physicochemical characteristics of the hyporheic zone affect redd site selection by chum salmon and fall Chinook salmon in the Columbia River. *North American Journal of Fisheries Management*, 22, 1077-1085.
- Gevao, B., Jones, K.C., Hamilton-Taylor, J. (1998). Polycyclic aromatic hydrocarbon (PAH) deposition to and processing in a small rural lake, Cumbria UK. *The Science of the Total Environment*, 215, 231-242.
- Ghadiri H., and Rose C. W. (1991). Sorbed chemical transport in overland flow: 1. A nutrient and pesticide enrichment mechanism. *Journal of Environmental Quality*, 20, 628-634.
- Giles, N. (1989). Assessing the status of British wild brown trout, *Salmo trutta*, stocks - a pilot-study utilizing data from game fisheries. *Freshwater Biology*, 21 (1), 125-133.
- Goedkoop, W., and Peterson, M. (2003). The fate, distribution, and toxicity of lindane in tests with *Chironomus riparius*: effects of bioturbation and sediment organic matter content. *Environmental Toxicology and Chemistry*, 22, 67-76.
- Gomez, L., Masot, J., Martinez, S., Duran, E., Soler, F., and Roncero, V. (1998). Acute 2,4-D poisoning in tench (*Tinca tinca* L.) lesions in the hematopoietic portion of kidney. *Archives of Environmental Contamination and Toxicology*, 35, 479-483.
- Gomez-Lahoz, C., and Ortega-Calvo, J. J. (2005). Effect of slow desorption on the kinetics of biodegradation of polycyclic aromatic hydrocarbons. *Environmental science and technology*, 39(22), 8776-8783.
- Gorge, G., and Nagel, R. (1990). Toxicity of Lindane, Atrazine, and Deltamethrin to Early Life Stages of Zebrafish (*Brachydanio rerio*). *Ecotoxicology and Environmental Safety*, 20, 246-255.
- Gorodilov, Y. N. (1989). Comparative analysis of the dynamics of early ontogeny of species of the genus *Salmo*. *Journal of Ichthyology*, 29, 19-31.
- Gorodilov, Y. N. (1996). Description of the early ontogeny of the Atlantic salmon, *Salmo salar*, with a novel system of interval (state) identification. *Environmental Biology of Fishes*, 47, 109-127.

- Graham, C., and Harrod, C. (2009). Implications of climate change for the fishes of the British Isles. *Journal of Fish Biology*, 74(6), 1143-1205.
- Greig, S., Sear, D., and Carling, P. (2005). The impact of fine sediment accumulation on the survival of incubating salmon progeny: implications for sediment management. *Science of the Total Environment*, 344(1-3), 241-258.
- Greig, S., Sear, D., Smallman, D., and Carling, P. (2005). Impact of clay particles on the cutaneous exchange of oxygen across the chorion of Atlantic salmon eggs. *Journal of Fish Biology*, 66(6), 1681-1691.
- Grost, R. T., Hubert, W. A., and Wesche, T. A. (1991). Description of brown trout redds in a mountain stream. *Transactions of the American Fisheries Society*, 120(5), 582-588.
- Guerrin, F., and Dumas, J. (2001a). Knowledge representation and qualitative simulation of salmon redd functioning. Part I: qualitative modeling and simulation. *BioSystems*, 59, 75-84.
- Guerrin, F., and Dumas, J. (2001b). Knowledge representation and qualitative simulation of salmon redd functioning. Part II: qualitative model of redds. *BioSystems*, 59, 85-108.
- Gunnes, K. (1979). Survival and development of Atlantic salmon eggs and fry at three different temperatures. *Aquaculture*, 16, 211-218.
- Gutenmann, W. H., Ebel, J. G., Kuntz, H. T., Yourstone, K. S., and Lisk, D. J. (1992). Residues of p, p'-DDE and mercury in lake trout as a function of age. *Archives of Environmental Contamination and Toxicology*, 22(4), 452-455.
- Haigh, S. D. (1996). A review of the interaction of surfactants with organic contaminants in soil. *Science of the Total Environment*, 185(1-3), 161-170.
- Hakanson, L. (1976). A Bottom Sediment Trap for Recent Sedimentary Deposits. *Limnology and Oceanography*, 21, 170-174.
- Hall, L. H., Kier L. B., and Phipps, G. (1984). Structure-activity relationship studies on the toxicities of benzene derivatives: I. An additivity model. *Environmental Toxicology and Chemistry*, 3, 355-365.
- Hamm, J. T., and Hinton, D. E. (2000). The role of development and duration of exposure to the embryotoxicity of diazinon. *Aquatic Toxicology*, 48, 403-418.
- Hancock, P. J. (2002). Human impacts on the stream-groundwater exchange zone. *Environmental Management*, 29(6), 763-781.
- Hansen, E. A. (1975). Some effects of groundwater on brown trout redds. *Transactions of the American Fisheries Society*, 1, 100-110.

Harrad, S. (2001). *Persistent organic pollutants: environmental behaviour and pathways of human exposure*: Springer Netherlands.

Harrad, S., and Mao, H. (2004). Atmospheric PCBs and organochlorines pesticides in Birmingham, UK: concentrations, sources, temporal and seasonal trends. *Atmospheric Environment*, 38, 1437–1445.

Harris, G. S. (1973). A simple egg box planting technique for estimating the survival of eggs deposited in stream gravel. *Journal of Fish Biology*, 5, 85-88.

Hartwell, S. I. (2008). Distribution of DDT and other persistent organic contaminants in canyons and on the continental shelf off the central California coast. *Marine Environmental Research*, 65(3), 199-217.

Harvey, J., Lyons, B., Waldock, M., and Parry, J. (1997). The application of the <sup>32</sup>P-postlabelling assay to aquatic biomonitoring. *Mutation Research/Fundamental and Molecular Mechanisms of Mutagenesis*, 378(1-2), 77-88.

Harwell, M. A., and Gentile, J. H. (2006). Ecological significance of residual exposures and effects from the Exxon Valdez oil spill. *Integrated environmental assessment and management*, 2(3), 204-246.

Hayes, T. B., Case, P., Chui, S., Chung, D., Haeffele, C., Haston, K., Lee, M., Mai, V. P., Marjuoa, Y., and Tsui, M. (2006). Pesticide mixtures, endocrine disruption, and amphibian declines: Are we understanding the impact? *Environmental Health Perspectives*, 114, 40–50.

Heath, A. G. (1995). *Water pollution and fish physiology*. CRC Press Inc. Boca Raton Florida.

Hela, D. G., Lambropoulou, D. A., Konstantinou, I. K., and Albanis, T.A. (2005) Environmental monitoring and ecological risk assessment for pesticide contamination and effects in Lake Pamvotis, northwestern Greece, *Environmental Toxicology and Chemistry*, 24, 1548–1556.

Heintz, R. A., Rice, S. D., Wertheimer, A. C., Bradshaw, R.F., Thrower, F.P., Joyce, J.E, Short, J.W. (2000). Delayed effects on growth and marine survival of pink salmon *Oncorhynchus gorbuscha* after exposure to crude oil during embryonic development. *Marine Ecology Progress Series*, 208, 205-216.

Heintz, R. A., Short, J. W. and Rice, S. D. (1999). Sensitivity of fish embryos to weathered crude oil: Part II. Increased mortality of pink salmon (*Oncorhynchus gorbuscha*) embryos incubating downstream from weathered Exxon Valdez crude oil. *Environmental Toxicology and Chemistry*, 18(3), 494-503.

Helder, T. (1980). Effects of 2,3,7,8-tetrachlorodibenzo-p-dioxin (TCDD) on early life stages of the pike (*Esox lucius* L.). *Science of the Total Environment*, 14, 255-264.

- Hellou, J., Leonard, J., Collier, T., and Ariese, F. (2006). Assessing PAH exposure in feral finfish from the Northwest Atlantic. *Marine Pollution Bulletin*, 52(4), 433-441.
- Hendricks, J., Meyers, T., Shelton, D., Casteel, J., and Bailey, G. (1985). Hepatocarcinogenicity of benzo [a] pyrene to rainbow trout by dietary exposure and intraperitoneal injection. *Journal of the National Cancer Institute*, 74(4), 839.
- Hendry, K., and Cragg-Hine, D. (2000). Ecology of the Atlantic salmon. *Conserving Natura*, 36.
- Hendry, K. and Cragg-Hine, D. (2003). Ecology of the Atlantic salmon. *Conserving Natura 2000 Rivers - Ecology Series No. 7*. English Nature, Peterborough. 32pp.
- Heywood, M., and Walling, D. (2007). The sedimentation of salmonid spawning gravels in the Hampshire Avon catchment, UK: implications for the dissolved oxygen content of intragravel water and embryo survival. *Hydrological Processes*, 21(6), 770-788.
- Heywood, M. J. T., and Walling, D. E. (2003). Suspended sediment fluxes in chalk streams in the Hampshire Avon Catchment, U.K. *Hydrobiologia*, 494, 111-117.
- Hicks, B. J., Hall, J. D., Bisson, P., and Sedell, J. R. (1991). Responses of salmonids to habitat changes. *Influences of Forest and Rangeland Management on Salmonid Habitat: American Fisheries Society Special Publication*, 19.
- Hilscherova, K., Dusek, L., Kubik, V., Cupr, P., Hofman, J., Klanova, J., and Holoubek, I. (2007). Redistribution of organic pollutants in river sediments and alluvial soils related to major floods. *Journal of Soils and Sediments*, 7(3), 167-177.
- Hinkle-Conn, C., Fleeger, J., Gregg, J., and Carman, K. (1998). Effects of sediment-bound polycyclic aromatic hydrocarbons on feeding behavior in juvenile spot (*Leiostomus xanthurus* Lacépède: Pisces). *Journal of experimental marine biology and ecology*, 227(1), 113-132.
- Hoai, P. M., Ngoc, N. T., Minh, N. H., Viet, P. H., Berg, M., Alder, A. C., and Giger, W. (2010). Recent levels of organochlorine pesticides and polychlorinated biphenyls in sediments of the sewer system in Hanoi, Vietnam. *Environmental Pollution*, 158(3), 913-920.
- Hobbs, D. F. (1937). Natural reproduction of quinnat salmon, brown and rainbow trout in certain New Zealand waters. *Fisheries Bulletin*, 6, 1-104.
- Hollert, H., Keiter, S., Konig, N., Rudolf, M., Ulrich, M., Braunbeck, T. (2003). A new sediment contact assay to assess particle-bound pollutants using zebrafish (*Danio rerio*) embryos. *J Soils and Sediments*, 3((3)), 197-207.

- Holth, T. F., Nourizadeh-Lillabadi, R., Blaesbjerg, M., Grung, M., Holbech, H., Petersen, G. I., Alestrom, P., and Hylland, K. ((2008). Differential gene expression and biomarkers in zebrafish (*Danio rerio*) following exposure to produced water components. *Aquatic Toxicology*, 90, 277-291.
- Honkanen, J. O. (2004): *Ecotoxicological testing of organic chemicals on early life stages of salmonid fish at environmentally realistic temperatures*. PhD Thesis, University of Joensuu.
- Honnen, W., Rath, K., Schlegel, T., Schwinger, A., and Frahne, D. (2001). Chemical analyses of water, sediment and biota in two small streams in southwest Germany. *Journal of Aquatic Ecosystem Stress and Recovery (Formerly Journal of Aquatic Ecosystem Health)*, 8(3), 195-213.
- Hose, J. E., Hannaht, J. B., Puffer, H. W., and Landolt, M. L. (1984). Histologic and skeletal abnormalities in benzo (a) pyrene-treated rainbow trout alevins. *Archives of Environmental Contamination and Toxicology*, 13(6), 675-684.
- Hose, J. E., Mcgurk, M. D., Marty, G. D., Hinton, D. E., Brown, E. D., and Baker, T. T. (1996). Sublethal effects of the Exxon Valdez oil spill on herring embryos and larvae: Morphological, cytogenic, and histopathological assessments, 1989-1991. *Canadian Journal of Fisheries and Aquatic Sciences*, 53(10), 2355-2365.
- Hoysak, D. J. and Liley, N. R. (2001). Fertilisation dynamics in sockeye salmon and a comparison of sperm from alternative male phenotypes. *Journal of Fish Biology*. 58, 1286-1300.
- Huber, W. (1993). Ecotoxicological relevance of atrazine in aquatic systems. *Environmental Toxicology and Chemistry*, 12, 1865-1881.
- Huskes, R., and Levsen, K. (1997). Pesticides in rain, *Chemosphere*, 35, 3013 – 3024.
- Hutchings, J. A. and Jones, M. E. B. (1998). Life history variation and growth rate thresholds for maturity in Atlantic salmon, *Salmo salar*. *Canadian Journal Fisheries and Aquatic Sciences*. 55, 22–47.
- Hutchinson, T. H., Shillabeer, N., Winter, M. J., and Pickford, D.B. (2006). Acute and chronic effects of carrier solvents in aquatic organisms: a critical review, *Aquatic Toxicology*, 76, 69–92.
- Hutson, D. H., and Roberts, T.R. (1990) *Environmental fate of pesticides*. Wiley. Chichester, England, pp123-47.
- Imanpoor, M. R., Alvai, S. H., and Cosson, J. (2009). Relationship between biological characteristics of eggs and female brood stocks of Persian sturgeon *Acipenser persicus*: a comparison with other fish species. *Fish Physiology and Biochemistry*, 35, 701-707.

Incardona, J. P., Collier, T. C., and Scholz, N. L. (2004). Defects in cardiac function precede morphological abnormalities in fish embryos exposed to polycyclic aromatic hydrocarbons *Toxicology and Applied Pharmacology*, 196, 191–205.

Ingendahl, D. (2001). Dissolved oxygen concentration and emergence of sea trout fry from natural redds in tributaries of the River Rhine. *Journal of Fish Biology*, 58, 325-341.

Inoue, T., Ebise, S., Numabe, A., Nagafuchi, O., and Matsui, Y. (2002). Runoff characteristics of particulate pesticides in a river from paddy fields. *Water Science and Technology*, 45, 121–126.

Iversen, E.R., Arnesen, R.T., and Knudsen, C.H. (2000). *Chemical treatment of acid mine drainage from the King's mine at Roeros, Norway*. In: ICARD 2000: Proceedings from the Fifth International Conference on Acid Rock Drainage. Littleton, CO: Society for Mining, Metallurgy and Exploration, pp. 1079-1086.

Jankowska, A., Biesaga, M., Drzewicz, P., Trojanowicz, M., and Pyrzynska, K. (2004). Chromatographic separation of chlorophenoxy acid herbicides their radiolytic degradation products in water sample. *Water Research*, 38, 3259–3264.

Jarvie, H. P., Neal, C., Withers, P. J. A., Wescott, C., and Acornley, R. M. (2005). Nutrient hydrochemistry for a groundwater-dominated catchment: The Hampshire Avon, UK. *Science of the Total Environment*, 344(1-3), 143-158.

Jarvie, H.P., Withers, P.J.A., Hodgkinson, R., Bates, A., Neal, M., Wickham, H. D., Harman, S.A. and Armstrong, L. (2008). Influence of rural land use on streamwater nutrients and their ecological significance. *Journal of Hydrology*, 350, 166-186

Jellyman, D., and Chisnall, B. (1999). Habitat preferences of shortfinned eels (*Anguilla australis*), in two New Zealand lowland lakes. *New Zealand Journal of Marine and Freshwater Research*, 33(2), 233-248.

Jensen, J. O. T., and Alderdice, D. F. (1983). Changes in mechanical shock sensitivity of coho salmon (*Oncorhynchus kisutch*) eggs during incubation. *Aquaculture*, 32, 303-312.

Jensen, A. J., and Johnsen, B. O. (1999). The functional relationship between peak spring floods and survival and growth of juvenile Atlantic Salmon (*Salmo salar*) and brown trout (*Salmo trutta*). *Functional Ecology*, 13, 778-785.

Johansson, N., Runn, P., and Milbrink, G. (1977). Early Development of Three Salmonid Species in Acidified Water. *Zoon*, 5, 127-132.

Johnson, A. C., Acreman, M. C., Dunbar, M. J., Feist, S. W., Giacomello, A. M., Gozlan, R. E. (2009). The British river of the future: How climate change and human activity might affect two contrasting river ecosystems in England. *Science of the Total Environment*, 407(17), 4787-4798.

Jones, J.W. (1959). *The salmon*. Collins, London.



- Jonsson, B., and Jonsson, N. (2009). A review of the likely effects of climate change on anadromous Atlantic salmon *Salmo salar* and brown trout *Salmo trutta*, with particular reference to water temperature and flow. *Journal of Fish Biology*, 75(10), 2381-2447.
- Johnstone, R. A. (1997). Evolution of animal signals. In: J. R. Krebs and N. B. Davies (Eds) *Behavioural ecology, 4th ed* (pp. 175-178). London: Blackwell Science.
- Julien, H.P. and Bergeron, N.E. (2006). Effect of fine sediment infiltration on Atlantic salmon (*Salmo salar*) embryo survival. *Hydrobiologia*, 563, 61-71.
- Kammann, U., Bunke, M., Steinhart, H., and Theobald, N. (2001). A permanent fish cell line (EPC) for genotoxicity testing of marine sediments with the comet assay. *Mutation Research/Genetic Toxicology and Environmental Mutagenesis*, 498(1-2), 67-77.
- Karickhoff, S. W., and Morris, K. R. (1985). Impact of tubificid oligochaetes on pollutant transport in bottom sediments. *Environmental science and technology*, 19(1), 51-56.
- Karr, J. R. (1981). Assessment of biotic integrity using fish communities. *Fisheries*, 6(6), 21-27.
- Keiter, S., Rastall, A., Kosmehl, T., Wurm, K., Erdinger, L., Braunbeck, T., and Hollert, H. (2006). Ecotoxicological assessment of sediment, suspended matter and water samples in the upper Danube River - A pilot study in search for the causes for the decline of fish catches. *Environmental Science and Pollution Research*, 13(5), 308-319.
- Kennedy, I., Ahmad, N., Beasley, H., Chapman, J., Hobbs, J., Simpson, B., and Woods, N. (1998). Quality Assurance in Pesticide Sampling and Analysis. *LWRRDC Occasional Paper No 14/98*.
- Kemp, P., Sear, D., Collins, A., Naden, P., and Jones, I. (2011). The impacts of fine sediment on riverine fish. *Hydrological Processes*.
- Killeen, J., McLay, H.A., Johnston, I.A. (1999). Development in *Salmo trutta* at different temperatures, with a quantitative scoring method for intraspecific comparisons. *Journal of Fish Biology*, 55, 382-404.
- Klamer, J. C., Hegeman, W. J. M., and Smedes, F. (1990). Comparison of grain-size correction procedures for organic micropollutants and heavy-metals in marine-sediments. *Hydrobiologia*, 208(3), 213-220.
- Knapp, R. A., and Vredenburg, V. T. (1996). A field comparison of the substrate composition of California golden trout redds sampled with two devices. *North American journal of fisheries management*, 16(3), 674-681.
- Kneusli, B. Berrer, D. Dupuis, G and Esser. H. (1969). S-Triazines. In: P. C. Kearney and D. D. Kaufman, (Eds) *Degradation of Herbicides*. (pp. 51). New York: Marcel Dekker.

- Knezovich, J. P., and Harrison, F. L. (1988). The bioavailability of sediment-sorbed chlorobenzenes to larvae of the midge, *Chironomus decorus*. *Ecotoxicology and environmental safety*, 15(2), 226-241.
- Knezovich, J. P., Harrison, F. L., and Wilhelm, R. G. (1987). The bioavailability of sediment-sorbed organic chemicals: A review. *Water, Air, and Soil Pollution*, 32(1-2), 233-245.
- Koehler, M. E., and Hardy, J. T. (1999). Effects of outboard motor emissions on early development of the killifish *Oryzias latipes*.
- Kondolf, G. M. (2000). Assessing salmonid spawning gravel quality. *Transactions of the American Fisheries Society*, 129(1), 262-281.
- Korn, S., and Rice, S. (1981). Sensitivity to, and accumulation and depuration of, aromatic petroleum components by early life stages of coho salmon *Oncorhynchus hutch.* *Rapports et Proces-Verbaux des Reunions Conseil International pour l'Exploration de la Mer*, 178, 87-92.
- Krein, A., and Schorer, M. (2000). Road runoff pollution by Polycyclic Aromatic Hydrocarbons and its Contribution to River Sediments. *Water Research*, 34(16), 4110 – 4115.
- Kreuger, J., Peterson, M., and Lundgren, E. (1999). Agricultural inputs of pesticide residues to stream and pond sediments in a small catchment in southern Sweden. *Bulletin of environmental contamination and toxicology*, 62(1), 55-62.
- Kristoffersen, J. B., and Magoulas, A. (2009). Fluctuating asymmetry and fitness correlations in two *Engraulis encrasicolus* populations. *Journal of Fish Biology*, 75, 2723-2736.
- Krumbein, W. C., and Pettijohn, F. J. (1938). *Manual of Sedimentary Petrography*. Appleton-Century-Crofts: New York.
- Kuang, S., and Xu, Z. (2009). Pollution characteristics of polycyclic aromatic hydrocarbons (PAHs) in oily sludge from the Zhongyuan Oilfield and its peripheral soils. *Chinese Journal of Geochemistry*, 28(2), 176-183.
- Kuhnhold, W. W., and Busch, F. (1977). On the uptake of three different types of hydrocarbons by salmon eggs (*Salmo salar* L.). *Meeresforsch*, 26, 50-59.
- Kulikova, N. A., and Perminova, I. V. (2002). Binding of atrazine to humic substances from soil, peat, and coal related to their structure. *Environmental science and technology*, 36(17), 3720-3724.

Lachance, S., Berube, P., and Lemieux, M. (2000). In situ survival and growth of three brook trout (*Salvelinus fontinalis*) strains subjected to acid conditions of anthropogenic origin at the egg and fingerling stages. *Canadian Journal of Fish Aquatic Sciences*, 57, 1562-1573.

Lacroix, G. L. (1985). Survival of Eggs and Alevins of Atlantic Salmon (*Salmo salar*) in Relation to the Chemistry of Interstitial Water in Redds in some Acidic Streams of Atlantic Canada. *Canadian Journal of Fish. Aquatic Sciences*, 42, 292-299.

Laetz, C. A., Baldwin, D. H., Collier, T. K., Hebert, V., Stark, J. D., and Scholz, N. L. (2009). The synergistic toxicity of pesticide mixtures: implications for risk assessment and the conservation of endangered Pacific salmon. *Environmental Health Perspectives*, 117(3), 348.

La Farré, M., Pérez, S., Kantiani, L, and Barceló, D. (2008). Fate and toxicity of emerging pollutants, their metabolites and transformation products in the aquatic environment. *Trends in Analytical Chemistry*, 27(11), 991-1007.

Lake, R. G., and Hinch, S. G. (1999). Acute effects of suspended sediment angularity on juvenile coho salmon (*Oncorhynchus kisutch*). *Canadian Journal of Fisheries and Aquatic Sciences*, 56(5), 862-867.

Lapointe, M. F., Bergeron, N.E., Bérubé, F., Pouliot, M. A. and Johnston, P. (2004). Interactive effects of substrate sand and silt contents, redd-scale hydraulic gradients, and interstitial velocities on egg-to-emergence survival of Atlantic salmon (*Salmo salar*). *Canadian Journal of Fisheries and Aquatic Science*, 61.

Larson, D. L., McDonald, S., Fivizzani, A. J., Newton, W. E., and Hamilton, S. J. (1998). Effects of the Herbicide Atrazine on *Ambystoma tigrinum* Metamorphosis: Duration, Larval Growth, and Hormonal Response. *Physiological Zoology*, 7, 671-679.

Larson, S. J., Capel, P. D., and Majewski, M. S. (1997), Pesticides in surface waters: distribution, trends, and governing factors: Chelsea, Michigan, Ann Arbor Press.

Laws, E. A. (1993). Aquatic pollution: An introductory text. John Wiley and Sons, Inc., New York.

Lawson, P., Logerwell, E., Mantua, N., Francis, R., and Agostini, V. (2004). Environmental factors influencing freshwater survival and smolt production in Pacific Northwest coho salmon (*Oncorhynchus kisutch*). *Canadian Journal of Fisheries and Aquatic Sciences*, 61(3), 360-373.

Lebo, J. A., Huckins, J. N., Petty, J. D., Cranor, W. L., and Ho, K. T. (2003). Comparisons of coarse and fine versions of two carbons for reducing the bioavailabilities of sediment-bound hydrophobic organic contaminants. *Chemosphere*, 50(10), 1309-1317.

- Lee, R. F., and Steinert, S. (2003). Use of the single cell gel electrophoresis/comet assay for detecting DNA damage in aquatic (marine and freshwater) animals. *Mutation Research/Reviews in Mutation Research*, 544(1), 43-64.
- Lentza-Rizos, C. (1996). Determination of triazine residues in water: comparison between a gas chromatographic method and an Enzyme-Linked Immunosorbent Assay (ELISA). *Bulletin of Environmental Contamination and Toxicology*, 57, 413-420.
- Leung, B., Forbes, M. R., and Houle, D. (2000). Fluctuating asymmetry as a bioindicator of stress: Comparing efficacy of analyses involving multiple traits. *The American Naturalist*, 155, 101–115.
- Lewis, W. M., and Morris, D. P. (1986). Toxicity of nitrite to fish: a review. *Transactions of the American Fisheries Society*, 115(2), 183-195.
- Liess, M., Schulz, R., Liess, M. H. D., Rother, B., and Kreuzig, R. (1999). Determination of insecticide contamination in agricultural headwater streams. *Water Research*, 33(1), 239-247.
- Lima, A. L. C., Eglinton, T. I., and Reddy, C. M. (2003). High-Resolution record of pyrogenic polycyclic aromatic hydrocarbon deposition during the 20th Century. *Environmental Science and Technology*, 37, 53–61.
- Linde, C.D. (1994). Physico-chemical properties and environmental fate of pesticides. Environmental Hazards Assessment Program, California.
- Linkov, I., Burmistrov, D., Cura, J., and Bridges, T.S. (2002). Risk-based management of contaminated sediments: Consideration of spatial and temporal patterns in exposure modeling. *Environmental Science and Technology*, 36 (2), 238–246.
- Lisle, T. E. (1989). Sediment transport and resulting deposition in spawning gravels, north coastal California. *Water Resources Research*, 25(6), 1303-1319.
- Lisle, T. E., and Eads, R. E. (1991). Methods to measure sedimentation of spawning gravels. *USDA Forest Service Research Note*, 411, 1-7.
- Lisle, T. E., and Lewis, J. (1992). Effects of sediment transport on survival of salmonid embryos in a natural stream: A simulation approach. *Canadian Journal of Fisheries and Aquatic Sciences*, 49, 2337-2344.
- Liu, W., Gan, J. J., Lee, S., and Kabashima, J. N. (2004). Phase distribution of synthetic pyrethroids in runoff and stream water. *Environmental Toxicology and Chemistry*, 23, 7–11.
- Lobpreis, T., Lopuchin, E., Vrana, B., Dercova, K., Mills, Graham and Greenwood, R. (2010) Monitoring of polycyclic aromatic hydrocarbons in the Portsmouth Harbour, United Kingdom, using the Chemcatcher® passive sampling devices. *Acta Chimica Slovaca*, 3(1), 81-94.

- Long, J. L. A., House, W. A., Parker, A., and Rae, J. E. (1998). Micro-organic compounds associated with sediments in the Humber rivers. *The Science of the Total Environment*, 210/211, 229-253.
- Lopes, I., Moreira-Santos, M., da Silva, E. M., Sousa, J. P., Guilhermino, L., Soares, A., Ribeiro, R. (2007). *In situ* assays with tropical cladocerans to evaluate edge-of-field pesticide runoff toxicity. *Chemosphere*, 67, 2250–2256.
- Lorenz, J. M., and Filer, J. H. (1989). Spawning habitat and redd characteristics of sockeye salmon in the glacial Taku River, British Columbia and Alaska. *Transactions of the American Fisheries Society*, 118(5), 495-502.
- Lotspeich, F. B., and Everest, F. H. (1981). A new method for reporting and interpreting textural composition of spawning gravel. *U.S. Department of Agriculture: Forest service*, 1-11.
- Lower, N., & Moore, A. (2003). Exposure to insecticides inhibits embryo development and emergence in Atlantic salmon. *Fish Physiology and Biochemistry* 28, 431-432.
- Luckenbach, T., Ferling, H., Gernhöfer, M., Köhler, H. R., Negele, R. D., Pfefferle, E., and Tribskorn, L. (2003). Developmental and subcellular effects of chronic exposure to sub-lethal concentrations of ammonia, PAH and PCP mixtures in brown trout (*Salmo trutta* f. *fario* L.) early life stages. *Aquatic Toxicology*, 65(1), 39-54.
- Luckenbach, T., Tribskorn, R., Müller, E., and Oberemm, A. (2001). Toxicity of waters from two streams to early life stages of brown trout (*Salmo trutta* f. *fario* L.), tested under semi-field conditions. *Chemosphere*, 45(4-5), 571-579.
- Lyons, G. (1997). Polyaromatic Hydrocarbons (PAHs), WWF-UK.
- Lyytikäinen, M., Jussi V., Kukkonen, K. and Lydy, M. J. (2003). Analysis of pesticides in water and sediment under different storage conditions using gas chromatography. *Archives of Environmental Contamination and Toxicology*, 44(4), 0437-0444
- MacGregor, K., Oliver, I., Duguid, A., and Ridgway. (2001). Persistent organic pollutants in scottish freshwater biota monitoring options, current levels and the way forward. *Scottish Environment Protection Agency Report*, 1-69.
- MacKenzie, C., and Moring J. R. (1988). Estimating survival of Atlantic salmon during the intragravel period. *North American Journal of Fisheries Management*, 8, 45-49.
- Mackay, A. A. and Gschwend, P. M. (2001) Enhanced concentrations of PAHs in groundwater at a coal tar site. *Environmental Science and Technology*, 35, 1320-1328.
- Mäenpää, K. A., Penttinen, O., and Kukkonen, J. V. K. (2004). Pentachlorophenol (PCP) bioaccumulation and effect on heat production on salmon eggs at different stages of development. *Aquatic Toxicology*, 68, 75-85.

- Maitland, P. S., Boon, P. J. and McLusky, D. S. (1994) *The freshwater of Scotland: A National Resource of International Significance*. New York: John Wiley. pp. 191-208.
- Maitland, P. S. and Campbell, R. N. (1992). *Freshwater Fishes of the British Isles*. New Naturalist Series, Harper Collins, London.
- Malcolm, I., Soulsby, C., Youngson, A., Hannah, D., McLaren, I., and Thorne, A. (2004). Hydrological influences on hyporheic water quality: implications for salmon egg survival. *Hydrological Processes*, 18(9), 1543-1560.
- Malcom, I. A., Soulsby, C., Youngson, A. F., and Petry, J. (2003a). Heterogeneity in ground water-surface water interactions in the hyporheic zone of a salmonid spawning stream. *Hydrological Processes*, 17, 601-617.
- Malcom, I. A., Youngson, A. F., and Soulsby, C. (2003b). Survival of salmonid eggs in a degraded gravel-bed stream: Effects of groundwater-surface water interactions. *River Research and Applications*, 19, 303-316.
- Mangani, G., Berloni, A., Bellucci, F., Tatano, F., and Maione, M. (2005). Evaluation of the pollutant content in road runoff first flush waters. *Water, Air, and Soil Pollution*, 160(1), 213-228.
- Mann, R. H. K, Blackburn, J. H. and Beaumont, W. R. C. (1989). The ecology of brown trout *Salmo trutta* in English chalk streams. *Freshwater Biology*, 21, 57-70.
- Marvin, C. H., Lundrigan, J. A., McCarry, B. E., and Bryant, D. W. (1995). Determination and genotoxicity of high molecular mass polycyclic aromatic hydrocarbons isolated from coal-tar contaminated sediment. *Environmental Toxicology and Chemistry* 14, 2059-2066.
- Mayer, P., Tolls, J., Hermens, J. L. M., and Mackay, D. (2003). Peer Reviewed: Equilibrium Sampling Devices. *Environmental science and technology*, 37(9), 184-191.
- Mazzi, D., and Bakker, T. C. M. (2001). Acid stress increases pelvic spine asymmetry in juvenile three-spined sticklebacks. *Journal of Fish Biology*, 59, 582-592.
- McDowall R.M. (2001). Anadromy and homing: two life-history traits with adaptive synergies in salmonid fishes? *Fish and Fisheries*. 2, 78-85.
- McNeil, W. J. (1962). Variations in the dissolved oxygen content of intragravel water in four spawning streams of Southeastern Alaska. *U.S. Fish and Wildlife Service- Special Scientific Report- Fisheries*, 402, 1-15.
- McNeil, W. J. (1964). A method of measuring mortality of pink salmon eggs and larvae. *Fishery Bulletin*, 63, 575-588.
- McNeil, W. J., and Ahnell, W. H. (1964). Success of pink salmon spawning relative to size of spawning bed materials. *U.S. Fish and Wildlife Service Special Scientific Report- Fisheries*, 469, 1-15.

- Meakins N. C., Bubb J. M., and Lester J. N. (1995). The mobility, partitioning and degradation of atrazine and simazine in the salt marsh environment. *Marine Pollution Bulletin* 30(12), 812-819.
- Menon, N. N., and Menon, N. (1999). Uptake of polycyclic aromatic hydrocarbons from suspended oil borne sediments by the marine bivalve *Sunetta scripta*. *Aquatic Toxicology*, 45(1), 63-69.
- Mersie, W., Liu, J., Seybold, C., and Tierney, D. (1998). Extractability and degradation of atrazine in a submerged sediment. *Weed science*, 480-486.
- Metcalf, T. L., Dillon, P. J., Metcalfe, C. D. (2008). Effects of formulations of the fungicide, pentachloronitrobenzene on the early life stage development of the Japanese medaka (*Oryzias latipes*). *Chemosphere*, 71, 1957–1962.
- Meudec, A., Dussauze, J., Deslandes, E., and Poupart, N. (2006). Evidence for bioaccumulation of PAHs within internal shoot tissues by a halophytic plant artificially exposed to petroleum-polluted sediments. *Chemosphere*, 65(3), 474-481.
- Meyer, C. B. (2003). The importance of measuring biotic and abiotic factors in the lower egg pocket to predict coho salmon egg survival. *Journal of Fish Biology*, 62, 534-548.
- Meyer, C. B., Madej, M. A., and Klein, R. D. (1993). Effects of fine sediment on salmonid redds in Prairie Creek, a tributary of Redwood Creek, Humboldt County, California. In: , S. D., Jr, Stohlgren, T. J., and Schonewald-Cox, C., Eds) *Proceedings of the Fourth Conference on Research in California's National Parks, NPS/NRUC/NRTP-93/9* (pp. 47-55).: Natural Resources Publication Office, National Park Service, Denver, Colorado
- Micic, M., Bihari, N., Labura, Z., Muller, W., and Batel, R. (2001). Induction of apoptosis in the blue mussel *Mytilus galloprovincialis* by tri-*n*-butyltin chloride, *Aquatic Toxicology*, 1 (55), 61–73.
- Miller, M. A. (1993). Maternal transfer of organochlorine compounds in salmonines to their eggs. *Canadian Journal of Fisheries and Aquatic Science*, 50, 1405-1413.
- Mills, D. (1989). *Ecology and management of Atlantic salmon*. Chapman and Hall Ltd. London.
- Miltner, R. J. And Rankin, E.T. (1998). Primary nutrients and the biotic integrity of rivers and streams. *Freshwater biology*, 40(1), 145-158.
- Miranda, A. L., Roche, H., Randi, M. A. F., Menezes, M. L., and Oliveira Ribeiro, C. A. (2008). Bioaccumulation of chlorinated pesticides and PCBs in the tropical freshwater fish *Hoplias malabaricus*: histopathological, physiological, and immunological findings. *Environment International*, 34, 939–949.

- Moore, A., Bendall, B., Barry, J., Waring, C., Crooks, N. and Crooks, L. (2012). River temperature and adult anadromous Atlantic salmon *Salmo salar* and brown trout *Salmo trutta*. *Fisheries Management and Ecology*, In Press.
- Moore, A., Scott, A. P., Lower, N., Katsiadaki, I., and Greenword, L. (2003). The effects of 4-nonylphenol and atrazine on Atlantic salmon (*Salmo salar* L.) smolts. *Aquaculture*, 222, 253-263.
- Moore, A., and Waring, C.P. (1996). Sublethal effects of the pesticide Diazinon on olfactory function in mature male Atlantic salmon parr. *Journal of Fish Biology* 48, 758-775.
- Moore, A., and Waring, C. P. (1998). Mechanistic Effects of a Triazine Pesticide on Reproductive Endocrine Function in Mature Male Atlantic Salmon (*Salmo salar* L.) Parr. *Pesticide Biochemistry and Physiology*, 62(1), 41-50.
- Moore, A., and Waring, C. P. (2001). The effects of a synthetic pyrethroid pesticide on some aspects of reproduction in Atlantic salmon (*Salmo salar* L.). *Aquatic Toxicology*, 52(1), 1-12.
- Moore, M. T., Rodgers, J. H., Cooper, C. M., and Smith Jr, S. (2000). Constructed wetlands for mitigation of atrazine-associated agricultural runoff. *Environmental Pollution*, 110, 393-399.
- Morales-Caselles, C., Lewis, C., Riba, I., DelValls, T. Á., and Galloway, T. (2009). A multibiomarker approach using the polychaete *Arenicola marina* to assess oil-contaminated sediments. *Environmental Science and Pollution Research*, 16(6), 618-629.
- Mueller, J. G., Middaugh, D., Lantz, S., and Chapman, P. (1991). Biodegradation of creosote and pentachlorophenol in contaminated groundwater: chemical and biological assessment. *Applied and Environmental Microbiology*, 57(5), 1277.
- Mukhi, S., Pan, X., Cobb, G. P., and Patino, R. (2005). Toxicity of hexahydro-1, 3, 5-trinitro-1, 3, 5-triazine to larval zebrafish (*Danio rerio*). *Chemosphere*, 61(2), 178-185.
- Müller, K. B., Hartmann, M., Spiteller, H., and Frede, M. (2002). Point-and nonpoint-source pesticide contamination in the Zwesten Ohm catchment, Germany. *Journal of environmental quality*, 31(1), 309.
- Naden, P., Smith, B., Jarvie, H., Llewellyn, N., Matthiessen, P, Dawson, H., Scarlett, P. and Hornby, D. (2003). Siltation in Rivers. A review of monitoring techniques. *Conserving Natura 2000 Rivers Conservation Techniques Series No. 6*. English Nature, Peterborough.
- Nagpal, N. K. (1993). Ambient water quality criteria for polycyclic aromatic hydrocarbons. Water Quality Branch, Water Management Division, Ministry of Environment, Lands and Parks, Victoria, British Columbia.



- Neal, C., Jarvie, H.P., Williams, R.J., Pinder, L. C., Collett, G. D., Neal, M., and Bhardwaj, L. (2000). The water quality of the Great Ouse. *Science of the Total Environment*, 251/252, 423–440.
- Neff, J. (1979). Polycyclic aromatic hydrocarbons in the aquatic environment: Sources, fates and biological effects. Applied Science Publishers Ltd: London.
- Nelms, B. E., Moravec, R., and Riss, T. (1997). Measuring apoptosis in individual cells with the comet assay. *Promega Notes*, 64, 13.
- Ngabé, B., and Poissant, L. (2003). Polycyclic aromatic hydrocarbons in the air in the St. Lawrence Basin (Québec). *Environmental science and technology*, 37(10), 2094-2099.
- Nongnutch, K., Cheevaporn, V., Helander, H. F., and Tangkrock-oland, N. (2011). Efficiency of methods to investigate paks exposure in fish (Nile Tilapia). *EnvironmentAsia*, 4(1), 6-11.
- Nuro, A., and Marku, E. (2011). Organochlorine Pesticides Residues for some Aquatic Systems in Albania. In: M, Stoytcheva (Ed). *Pesticides - Formulations, Effects, Fate*. InTech (Open).
- Nwani, C.D., Nagpure, N.S., Kumar, R., Kushwaha, B., Kumar, P., and Lakra, W.S. (2011). Mutagenic and genotoxic assessment of atrazine-based herbicide to freshwater fish *Channa punctatus* (Bloch) using micronucleus test and single cell gel electrophoresis. *Environmental Toxicology and Pharmacology*, 31, 314-322.
- O'Connor, W. C. K., and Andrew, T. E. (1998). The effects of siltation on Atlantic salmon, *Salmo salar* L., embryos in the River Bush. *Fisheries Management and Ecology*, 5, 393-401.
- O'Flynn, F.M., McGeachy, S.A., Friars, G.W., Benfey, T.J., and Bailey, J.K. (1997). Comparisons of cultured triploid and diploid Atlantic salmon (*Salmo salar* L.). *ICES Journal of Marine Science*, 54, 1160-1165.
- Oldershaw, C. (2002). The earth in our hands: Groundwater. Geological Society of London.
- Oliva, M., Garrido, C., Sales, D., and González de Canales, M. L. (2008). Lindane toxicity on early life stages of gilthead seabream (*Sparus aurata*) with a note on its histopathological manifestations. *Environmental Toxicology and Pharmacology*, 25(1), 94-102.
- Oliveira Ribeiro, C. A., Vollaire, Y., Sanchez-Chardi, A., and Roche, H. (2005). Bioaccumulation and the effects of organochlorine pesticides, PAH and heavy metals in the eel (*Anguilla anguilla*) at the Camargue Nature Reserve, France. *Aquatic Toxicology*, 74, 53–69.

- Olsson, T. I., and Persson, B. (1986). Effects of gravel size and peat material concentrations on embryo survival and alevin emergence of brown trout, *Salmo trutta* L. *Hydrobiologia*, 135, 9-14.
- Olsson, T. I., and Persson, B. (1988). Effects of deposited sand on ova survival and alevin emergence in brown trout (*Salmo trutta* L.). *Archeology and Hydrobiology*, 113, 621-627.
- Orghidan, T. (1959). Ein neuer Lebensraum des unterirdischen Wassers: der Hyporheische Biotop. *Archives of Hydrobiology*, 55, 392-414.
- Ortlepp, J., and Mürle, U. (2003). Effects of experimental flooding on brown trout (*Salmo trutta* fario L.): The River Spöl, Swiss National Park. *Aquatic Sciences-Research Across Boundaries*, 65(3), 232-238.
- Østbye, K., Øxnevad, S. A., and Vøllestad, L. A. (1997). Developmental stability in perch (*Perca fluviatilis*) in acidic aluminium rich lakes. *Canadian Journal of Zoology*, 75, 919-928.
- Ostrander, G. K. (2004). *Handbook of Techniques in Aquatic Toxicology-Volume 2*. Lewis Publishers/CRC Press.
- Osterauer, R. and Köhler, H. R. (2008). Temperature dependent effects of the pesticides thiacloprid and diazinon on the embryonic development of zebrafish (*Danio rerio*). *Aquatic Toxicology*, 86, 485-494.
- Ostrander, G. K., Landolt, M. L., and Kocan, R. M. (1988). The ontogeny of coho salmon (*Oncorhynchus kisutch*) behaviour following embryonic exposure to benzo[a]pyrene. *Aquatic Toxicology*, 13, 325-346.
- Øxnevad, S. A., Heibo, E., and Vollestad, L. A. (2002). Is there a relationship between fluctuating asymmetry and reproductive investment in perch (*Perca fluviatilis*)? *Canadian Journal of Zoology*, 80, 120-125.
- Ozcan Oruc, E., Sevgiler, Y and Uner, N. (2004). Tissue-specific oxidative stress responses in fish exposed to 2,4-D and azinphosmethyl. *Comparative Biochemistry and Physiology Part C*, 137, 43-51.
- Paine, J. and Leggett, W. (1992). Effects of Hibernia crude oil on capelin (*Mallotus villosus*) embryos and larvae. *Marine Environmental Research*, 33, 159-187.
- Palma, P., Palma, V., Matos, C., Fernandes, R., Bohn, A., Soares, A., and Barbosa, I. R. (2009). Effects of atrazine and endosulfan sulphate on the ecdysteroid system of *Daphnia magna*. *Chemosphere*, 74(5), 676-681.
- Parker, J. T. C., Fossum, K.D., Ingersoll, T.L. (2000). Environmental auditing: Chemical characteristics of urban stormwater sediments and implications for environmental Management, Maricopa County, Arizona. *Environmental Management*, 26, 99-115.

- Parrish, D. L., Behnke, R. J., Gephard, S. R., McCormick, S. D., and Reeves, G. H. (1998). Why aren't there more Atlantic salmon (*Salmo salar*)? *Canadian Journal of Fisheries and Aquatic Sciences*, 55(Supplement 1), 281-287.
- Peer, M. M., Nirmala, J., and Kutty, M. (1983). Effects of pentachlorophenol (Na PCP) on survival, activity and metabolism in *Rhinomugil corsula* (Hamilton), *Cyprinus carpio* (Linnaeus) and *Tilapia mossambica* (Peters). *Hydrobiologia*, 107(1), 19-24.
- Petersen, G. I., and Kristensen, P. (1998). Bioaccumulation of lipophilic substances in fish early life stages. *Environmental Toxicology and Chemistry*, 17, 1385-1395.
- Peterson, N. P., and Quinn, T. P. (1996a). Persistence of egg pocket architecture in redds of chum salmon, *Oncorhynchus keta*. *Environmental Biology of Fishes*, 46, 243-253.
- Peterson, N. P., and Quinn, T. P. (1996b). Spatial and temporal variation in dissolved oxygen in natural egg pockets of chum salmon, in Kennedy Creek, Washington. *Journal of Fish Biology*, 48, 131-143.
- Peterson, R. H., and Metcalfe, J. L. (1981). Emergence of atlantic salmon fry from gravels of varying composition: A laboratory study. *Fisheries and Oceans*, 1020, 1-15.
- Peterson, R. H., Spinney, H. C. E., and Sreedharan, A. (1976). Development of Atlantic salmon (*Salmo salar*) eggs and alevins under varied temperature regimes. *Journal of Fisheries Research Board of Canada*, 34, 31-43.
- Petit, V., Cabridenc, R., Swannell, R. P. J., and Sokhi, R. S. (1995). Review of strategies for modelling the environmental fate of pesticides discharged into riverine systems. *Environment International*, 21, 167-176.
- Petts, G. E., Thoms, M. C., Brittan, K., and Atkin, B. (1989). A freeze-coring technique applied to pollution by fine sediments in gravel-bed rivers. *Science of the Total Environment*, 84, 259-272.
- Petty, J. D., Huckins, J. N., Orazio, C. E., Lebo, J. A., Poulton, B. C., Gale, R. W., Charbonneau, C. S., and Kaiser, E. M. (1995). Determination of waterborne bioavailable organochlorine pesticide residues in the lower Missouri River. *Environmental science and technology*, 29(10), 2561-2566.
- Phyu, Y., Warne, M. S. and Lim, R. P. (2005). The toxicity and bioavailability of atrazine and molinate to *Chironomus tepperi* larvae in laboratory and river water in the presence and absence of sediment. *Chemosphere*, 58(9), 1231-1239.
- Phyu, Y. L., Warne, M. S. and Lim, R.P. (2006). Toxicity and bioavailability of atrazine and molinate to the freshwater fish (*Melanotenia fluviatilis*) under laboratory and simulated field conditions. *Science of the Total Environment*, 356, 86-99.
- Pitt, R., Clark, S., and Field, R. (1999). Groundwater contamination potential from stormwater infiltration practices. *Urban Water*, 1(3), 217-236.

- Pitt, R., Field, R., Lalor, N., and Brown, M. (1994). *Potential groundwater contamination from intentional and nonintentional stormwater infiltration*. Washington, US: Environmental Protection Agency.
- Pittinger, C. A., Woltering, D. M., and Masters, J. A. (1989). Bioavailability of sediment sorbed and aqueous surfactants to *Chironomus riparius* (midge). *Environmental Toxicology and Chemistry*, 8(11), 1023-1033.
- Pollard, R. A. (1955). Measuring seepage through salmon spawning gravel. *Journal of Fisheries Research Board of Canada*, 12, 706-741.
- Portz, D. E., Woodley, C. M., and Cech, J. J. (2006). Stress-associated impacts of short-term holding on fishes. *Reviews in Fish Biology and Fisheries*, 16(2), 125-170.
- Potter, I. C. (1980). Ecology of larval and metamorphosing lampreys. *Canadian Journal of Fisheries and Aquatic Science*, 37, 1641-1657.
- Potts, W. T. W. and Rudy, P. P. JR (1969). Water balance in the eggs of the Atlantic salmon *Salmo salar*. *Journal of experimental Biology*, 50, 223-37.
- Pyefinch, K. A. (1972). Atlantic salmon in the sea. *Proceedings of the Royal Society of Edinburgh (B)*, 423-27.
- Qin, X., Sun, H., Wang, C., Yu, Y., and Sun, T. (2010). Impacts of crab bioturbation on the fate of polycyclic aromatic hydrocarbons in sediment from the Beitang estuary of Tianjin, China. *Environmental Toxicology and Chemistry*, 29(6), 1248-1255.
- Rae, J. E., Cooper, C. S., Parker, A., and Peters, A. (1998). Pesticide Sorption onto Aquifer Sediments. *Journal of Geochemical Exploration*, 64, 263-276.
- Rao, D. M. R., and Murty, A. S. (1980). Toxicity, biotransformation and elimination of endosulfan in *Anabas testudineus* (Bloch). *Indian Journal of experimental Biology*. 18, 664-665.
- Reaney, S. M., Lane, S. N., Heathwaite, A. L., and Dugdale, L. J. (2010). Risk-based modelling of diffuse land use impacts from rural landscapes upon salmonid fry abundance. *Ecological Modelling*.
- Regetz, J. (2003). Landscape level constraints on recruitment of chinook salmon (*Oncorhynchus tshawytscha*) in the Columbia River basin, USA. *Aquatic Conservation: Marine and Freshwater Ecosystems*, 13(1), 35-49.
- Reible, D., Popov, V., Valsaraj, K., Thibodeaux, L., Lin, F., Dikshit, M., Todaro, M. A., and Fleeger, J. W. (1996). Contaminant fluxes from sediment due to tubificid oligochaete bioturbation. *Water Research*, 30(3), 704-714.

- Reiser, D. W., and White, R. G. (1988). Effects of two sediment size-classes on survival of steelhead and chinook salmon eggs. *North American Journal of Fisheries Management*, 8, 432-437.
- Reynoldson, T. B. (1987). Interactions between sediment contaminants and benthic organisms. *Hydrobiologia*, 149(1), 53-66.
- Rhodes, S., Farwell, A., Hewitt, L. M., MacKinnon, M., Dixon, D. G. (2005). The effects of dimethylated and alkylated polycyclic aromatic hydrocarbons on the embryonic development of Japanese medaka. *Ecotoxicology and Environmental Safety*, 60, 247-258.
- Rice, S. D., and Bailey, J. E. (1980). Survival, size, and emergence of pink salmon, *Oncorhynchus gorbuscha*, alevins after short- and long-term exposures to ammonia. *Fisheries Bulletin U.S.* 78, 641-648.
- Riley, W.D., Ives, M.J., Pawson, M.G., and Maxwell, D.L. (2006). Seasonal variation in habitat use by salmon, *Salmo salar*, trout, *Salmo trutta*, and grayling, *Thymallus thymallus*, in a chalk stream. *Fisheries Management and Ecology*, 13, 221-236.
- Rollinson, N., and Hutchings, J.A. (2010). Why does egg size increase with maternal size? Effects of egg size and egg density on offspring phenotypes in Atlantic salmon (*Salmo salar*). *Evolutionary Ecology Research*, 12, 949-960.
- Rombough, P. J. (2007). Oxygen as a constraining factor in egg size evolution of salmonids. *Canadian Journal of Fisheries and Aquatic Sciences*, 64, 692-699.
- Ross, P. S., De Swart, R. L., Reijnders, P., Van Loveren, H., Vos, J. G., and Osterhaus, A. (1995). Contaminant-related suppression of delayed-type hypersensitivity and antibody responses in harbor seals fed herring from the Baltic Sea. *Environmental Health Perspectives*, 103(2), 162.
- Rubin, J. F., and Glümsater, C. (1996). Egg-to-fry survival of the sea trout in some streams of Gotland. *Journal of Fish Biology*, 48, 585-606.
- Runkel, R. L., McKnight, D. M., and Rajaram, H. (2003). Modeling hyporheic zone processes. *Advances in Water Resources*, 26, 901-905.
- Russo, R. C., and Thurston, R. V. (1991). Toxicity of ammonia, nitrite, and nitrate to fishes. *Aquaculture and water quality*. World Aquaculture Society, Baton Rouge, Louisiana, 58-89.
- Sarikaya, R., and Yilmaz, M. (2003). Investigation of acute toxicity and the effect of 2,4-D (2,4-dichlorophenoxyacetic acid) herbicide on the behavior of the common carp (*Cyprinus carpio* L., 1758; Pisces, Cyprinidae). *Chemosphere*, 52(1), 195-201.
- Saunders, J., and Smith, M. (1965). *Changes in a stream population of trout associated with increased silt*: Fisheries Research Board of Canada.

- Sawyer, H. W. (1959) Burrowing activities of the larval lampreys. *Copeia*, 1959, 256- 257
- Schafer, R. B., Mueller, R., Brack, W., Wenzel, K. D, Streck, G., Ruck, W., and Liess, M. (2008). Determination of 10 particle-associated multiclass polar and semi-polar pesticides from small streams using accelerated solvent extraction. *Chemosphere*, 70, 1952-1960.
- Schulz, R. (2001). Rainfall-induced sediment and pesticide input from orchards into the Lourens River, western cape, South Africa: Importance of a single event. *Water Research*, 35, 1869-1876.
- Scorecard- The pollution information website (2005). *Organochlorine Pesticides*. [http://www.scorecard.org/about/txt/organochlorine\\_pesticides.html](http://www.scorecard.org/about/txt/organochlorine_pesticides.html).
- Schriever, C. A., von der Ohe, P. C., and Liess, M. (2007). Estimating pesticide runoff in small streams. *Chemosphere*, 68, 2161–2171.
- Sear, D. A. (1993). Fine sediment infiltration into gravel spawning beds within a regulated river experiencing floods: Ecological implications for salmonids. *Regulated Rivers: Research and Management*, 8, 373-390.
- Segerson, K. (1990). Liability for groundwater contamination from pesticides. *Journal of Environmental Economics and Management*, 19(3), 227-243.
- Sethajintanin, D., and Anderson, K. A. (2006). Temporal bioavailability of organochlorine pesticides and PCBs. *Environmental Science and Technology*, 40(12), 3689-3695.
- Shirazi, M. A., and Seim. W. K. (1979). A stream system evaluation - an emphasis on spawning habitat for salmonids. *U.S. Environmental Protection Agency, EPA -600/3-79-109*, Corvallis Research Laboratory. Corvallis, Oregon, 36 p.
- Singh, N. P., McCoy, M. T., Tice, R. R., and Schneider, E. L. (1988): A simple technique for quantitation of low levels of DNA damage in individual cells. *Experimental Cell Research*, 175(1), 184-191.
- Sipaúba-Tavares, L., Guariglia, C., and Braga, F. (2007). Effects of rainfall on water quality in six sequentially disposed fishponds with continuous water flow. *Brazilian Journal of Biology*, 67(4), 643-649.
- Skei, J., Larsson, P., Rosenberg, R., Jonsson, P., Olsson, M., and Broman, D. (2000). Eutrophication and contaminants in aquatic ecosystems. *Ambio*, 184-194.
- Smith, V. H. (2006). Responses of estuarine and coastal marine phytoplankton to nitrogen and phosphorus enrichment. *Limnology and Oceanography*, 51, 377-384.
- Smith, V. H., Tilman, G. D., and Nekola, J. C. (1999). Eutrophication: impacts of excess nutrient inputs on freshwater, marine and terrestrial ecosystems. *Environmental Pollution*, 100, 179–196.

Solbakken, J. E., Tilseth, S., and Palmork, K.H. (1984) Uptake and elimination of aromatic hydrocarbons and chlorinated biphenyl in eggs and larvae of cod *Gadus morhua*. *Marine Ecology Progress Service*, 16, 297-301.

Solomon, K. R., Baker, D. B., Richards, R. P., Dixon, K. R., Klaine, S. J., La Point, T. W., Kendall, R. J., Weisskopf, C. P., Giddings, J. M., and Giesy, J. P. (1996). Ecological risk assessment of atrazine in North American surface waters. *Environmental Toxicology and Chemistry*, 15, 31-76.

Somarakis, S., Kostikas, I., Peristeraki, N., Tsimenides, N. (1997). Fluctuating asymmetry in the otoliths of larval anchovy *Engraulis encrasicolus* and the use of developmental instability as an indicator of condition in larval fish. *Marine Ecology Progress Series*, 151, 191– 203.

Soto-Galera, E., Díaz-Pardo, E., López-López, E., and Lyons, J. (1998). Fish as indicators of environmental quality in the Río Lerma Basin, México. *Aquatic Ecosystem Health and Management*, 1(3), 267-276.

Soulsby, C., Malcolm, I., and Youngson, A. (2001a). Hydrochemistry of the hyporheic zone in salmon spawning gravels: a preliminary assessment in a degraded agricultural stream. *Regulated Rivers: Research and Management*, 17(6), 651-665.

Soulsby, C., Youngson, A. F., Moir, H. J., and Malcom, I. A. (2001b). Fine sediment influence on salmonid spawning habitat in lowland agricultural stream: a preliminary assessment. *The Science of the Total Environment*, 265, 295-307.

Sowden, T. K., and Power, G. (1985). Prediction of rainbow trout embryo survival in relation to groundwater seepage and particle size of spawning substrate. *Transactions of the American Fisheries Society*, 114, 804-812.

Srivastava, R. K., Brown, J. A., and Shahidi, F. (1995). Changes in the amino acid pool during embryonic development of cultured and wild Atlantic salmon (*Salmo salar*). *Aquaculture*, 131, 115–124.

Ssebugere, P., Kiremire, B. T., Kishimba, M., Wandiga, S. O., Nyanzi, S. A., and Wasswa, J. (2009). DDT and metabolites in fish from Lake Edward, Uganda. *Chemosphere*, 76(2), 212-215.

Stackelberg, P. E. (1997). Presence and distribution of chlorinated organic compounds in streambed sediments, New Jersey. *Journal of the American Water Resources Association*, 33(2), 271-284.

Standridge, J.H., and Lesar, D.J. (1977). Comparison of four-hour and twenty-four-hour refrigerated storage of nonpotable water for fecal coliform analysis. *Applied Environmental Microbiology*, 34, 398-402.

- Steele, S. L., Chadwick, T. D. and Wright, P. A. (2001). Ammonia detoxification and localization of urea cycle enzyme activity in embryos of the rainbow trout (*Oncorhynchus mykiss*) in relation to early tolerance to high environmental ammonia levels. *Journal of Experimental Biology*, 204, 2145-2154.
- Stoker, T.E., and Cooper, R. L. (2007). Distribution of <sup>14</sup>C-atrazine following an acute lactational exposure in the Wistar rat. *Reproductive Toxicology*, 23, 607–610.
- Strmac, M., Oberemm, A. and Braunbeck, T. (2002). Effects of sediment eluates and extracts from differently polluted small rivers on zebrafish embryos and larvae. *Journal of Fish Biology*, 61, 24–38.
- Stuer-Lauridsen, F. (2005). Review of passive accumulation devices for monitoring organic micropollutants in the aquatic environment. *Environmental Pollution*, 136(3), 503-524.
- Summers, D. W., Giles, N., Stubbing, D. N. (2005). The effect of riparian grazing on brown trout, *Salmo trutta*, and juvenile Atlantic salmon, *Salmo salar*, in an English chalk stream. *Fisheries Management and Ecology*, 12(6), 403–405.
- Sun, Y., Yu, H., Zhang, J., Yin, Y., Shi, H., and Wang, X. (2006). Bioaccumulation, depuration and oxidative stress in fish *Carassius auratus* under phenanthrene exposure. *Chemosphere*, 63(8), 1319-1327.
- Talley, J. W., Ghosh, U., Tucker, S. G., Furey, J. S., and Luthy, R.G. (2002). Particle-scale understanding of the bioavailability of PAHs in sediment, *Environmental Science and Technology*, 36, 477–483.
- Tautz, A.F. and Groot, C. (1975). Spawning behavior of chum salmon (*Oncorhynchus keta*) and rainbow trout (*Salmo gairdneri*). *Journal of the Fisheries Research Board of Canada*. 32, 633–642.
- Taylor, K. G., Owens, P. N., Batalla, R. J., and Garcia, C. (2008). Sediment and contaminant sources and transfers in river basins. In N. O. Dr. Philip (Ed.), *Sustainable Management of Sediment Resources* (Vol. Volume 4, pp. 83-135): Elsevier.
- Thorpe, J. E., Miles, M. S., and Keay, D. S. (1984). Developmental rate, fecundity and egg size in Atlantic salmon, *Salmo salar* L. *Aquaculture*, 43, 289-305.
- Tillitt, D. E., Papoulias, D. M., Whyte, J. J., and Richter, C. A. (2010). Atrazine reduces reproduction in fathead minnow (*Pimephales promelas*). *Aquatic Toxicology*, 99(2), 149-159.
- Timmons, M. B., Ebeling, J. M., Wheaton, F. W., Summerfelt, S. T., and Vinci, B.J. (2001). *Recirculating Aquaculture systems*. Cayuga Aqua Ventures. Ithaca, New York.
- Tomlin, C. (2000). *The pesticide manual: a world compendium*. Twelfth ed. Farnham (UK): The British Crop Protection Council.



- Turnpenny, A. W. H., and Williams, R. (1980). Effects of sedimentation on the gravels of an industrial river system. *Journal of Fish Biology*, 17, 681-693.
- Umlauf, G., and Bierl, R. (1987). Distribution of organic micropollutants in different size fractions of sediment and suspended solid particles of the River Rotmain. *Zeitschrift für Wasser-und Abwasser-Forschung*, 20(6), 203-209.
- Uriarte, A., and Borja, A. (2009). Assessing fish quality status in transitional waters, within the European Water Framework Directive: setting boundary classes and responding to anthropogenic pressures. *Estuarine, Coastal and Shelf Science*, 82(2), 214-224.
- Vagi, M. C., Petsas, A. S., Kostopoulou, M. N., Karamanoli, M. K., and Lekkas, T. B. (2007). Determination of organochlorine pesticides in marine sediments samples using ultrasonic solvent extraction followed by GC/ECD. *Desalination*, 210, 146-156.
- Van Leeuwen, C., Griffioen, P., Vergouw, W., and Maas-Diepeveen, J. (1985). Differences in susceptibility of early life stages of rainbow trout (*Salmo gairdneri*) to environmental pollutants. *Aquatic Toxicology*, 7(1-2), 59-78.
- Vasquez, M. Z. (2010). Combining the in vivo comet and micronucleus assays: A practical approach to genotoxicity testing and data interpretation. *Mutagenesis*, 25, 187-199.
- Viant, M. R., Pincetich, C. A., and Tjeerdema, R. S. (2006). Metabolic effects of dinoseb, diazinon and esfenvalerate in eyed eggs and alevins of Chinook salmon (*Oncorhynchus tshawytscha*) determined by 1H NMR metabolomics. *Aquatic Toxicology*, 77(4), 359-371.
- Vigano, L., Arillo, A., Falugi, C., Melodia, F. (1998). Histochemical and biochemical markers in trout larvae exposed to river sediments. *Chemosphere*, 37, 2797-2807.
- Vink, J. P. M., and Van der Zee, S. E. (1997). Pesticide biotransformation in surface waters: multivariate analyses of environmental factors at field sites. *Water Research*, 31(11), 2858-2868.
- Vitousek, P. M., Mooney, H. A., Lubchenco, J., and Melillo, J. M. (1997). Human domination of Earth's ecosystems. *Science*, 277(5325), 494.
- Vives, I., Grimalt, J. O., Fernandez, P., and Rosseland, B. (2004). Polycyclic aromatic hydrocarbons in fish from remote and high mountain lakes in Europe and Greenland. *Science of the Total Environment*, 324(1-3), 67-77.
- Vogue, P. A., Kerle, E. A. and Jenkins, J. J. (1998) OSU extension pesticide properties database. EM 8709.
- Vollestad, L. A., and Hindar, K. (1997). Developmental stability and environmental stress in *Salmo salar* (Atlantic salmon). *Heredity*, 78, 215-222.

Voorspoels, S., Covaci, A., Maervoet, J., De Meester, I., and Schepens, P. (2004). Levels and profiles of PCBs and OCPs in marine benthic species from the Belgian North Sea and the Western Scheldt Estuary. *Marine Pollution Bulletin*, 49(5-6), 393-404.

Vrzoc, M., and Petras, M. L. (1997). Comparison of alkaline single cell gel (Comet) and peripheral blood micronucleus assays in detecting DNA damage caused by direct and indirect acting mutagens. *Mutation Research*, 1381(1), 31-40.

Waddington, C. H. (1942). Canalization of development and the inheritance of acquired characters. *Nature*, 150, 563-565.

Wagner, E. J. (1996). History and fluctuating asymmetry of Utah salmonid broodstocks. *The Progressive Fish-Culturist*, 58, 92-103.

Wakefield, A., Cunjak, R., and Kieffer, J. (2004). Metabolic recovery in Atlantic salmon fry and parr following forced activity. *Journal of Fish Biology*, 65(4), 920-932.

Walling, D. E., and Amos, C. M. (1999). Source, storage and mobilisation of the fine sediment in a chalk stream system. *Hydrological Processes*, 13, 323-340.

Wan, M. T., Buday, C., Schroeder, G., Kuo, J., and Pasternak, J. (2006). Toxicity to *Daphnia magna*, *Hyalella azteca*, *Oncorhynchus kisutch*, *Oncorhynchus mykiss*, *Oncorhynchus tshawytscha*, and *Rana catesbeiana* of atrazine, metolachlor, simazine, and their formulated products. *Bulletin of Environmental Contamination and Toxicology*, 76, 52-58.

Waring, C., and Moore, A. (1997). Sublethal effects of a carbamate pesticide on pheromonal mediated endocrine function in mature male Atlantic salmon (*Salmo salar* L.) parr. *Fish Physiology and Biochemistry*, 17(1), 203-211.

Waring, C. P., and Moore, A. (2004). The effect of atrazine on Atlantic salmon (*Salmo salar*) smolts in fresh water and after sea water transfer. *Aquatic Toxicology*, 66(1), 93-104.

Waring, C. P., Moore, A., Best, J. H., Crooks, N., and Crooks, L. E. (2011). Do trout farm effluents affect Atlantic salmon smolts? Preliminary studies using caged salmon smolts, *Aquaculture*, (*In Press*).

Warren, N., Allan, I. J., Carter, J. E., House, W. A., and Parker, A. (2003). Pesticides and other micro-organic contaminants in freshwater sedimentary environments- A review. *Applied Geochemistry*, 18, 159-194.

Weeber, M. A. (2006). *Effects of Kokanee (Oncorhynchus nerka) Redd Superimposition on Bull Trout (Salvelinus confluentus) Reproductive Success in the Deschutes River Basin, Oregon*. PhD Thesis. Oregon State University.

Weis, J. S., and Weis, P. (1989). Effects of environmental pollutants on early fish development. *Aquatic Sciences*, 1, 45-73.

- Welton, J. S., Beaumont, W. R. C., and Ladle, M. (1999). Timing of migration and changes in age structure of Atlantic salmon, *Salmo salar* L., in the River Frome, a Dorset chalk stream, over a 24-year period. *Fisheries Management and Ecology*, 6, 437-458.
- Wentworth, C.K. (1922). A scale of grade and class terms for clastic sediments, *Journal of Geology*, 30, 377-392.
- Wessel, N., Rousseau, S., Caisey, X., Quiniou, F., and Akcha, F. (2007). Investigating the relationship between embryotoxic and genotoxic effects of benzo [a] pyrene, 17 [alpha]-ethinylestradiol and endosulfan on *Crassostrea gigas* embryos. *Aquatic Toxicology*, 85(2), 133-142.
- Westin, D. T., and B. A. Rogers. (1978). *Synopsis of the biological data on the striped bass*. University of Rhode Island Marine Technical Report No. 67, Kingston, Rhode Island.
- Wheeldon, J. (2003) *River Avon cSAC Conservation Strategy*. English Nature, Peterborough.
- White, H. C. (1942). Atlantic salmon redds and artificial spawning beds. *Journal of Fisheries Research Board of Canada*, 6, 37-44.
- Wiegand, C., Krause, E., Steinberg, C., and Pflugmacher, S. (2001). Toxicokinetics of atrazine in embryos of the zebrafish (*Danio rerio*). *Ecotoxicology and Environmental Safety*, 49, 199-205.
- Wiegand, C., Pflugmacher, S., Giese, M., Frank, H., and Steinberg, C. (2000). Uptake, toxicity, and effects on detoxication enzymes of atrazine and trifluoroacetate in embryos of zebrafish. *Ecotoxicology and Environmental Safety*, 45, 122-131.
- Wijbenga A, and Hutzinger, O. (1984) Chemicals, man and the environment: a historic perspective of pollution and related topics. *Naturwissenschaften*. 71(5):239-46.
- Wild, S. R., and Jones, K. C. (1995). Polynuclear aromatic hydrocarbons in the United Kingdom environment: a preliminary source inventory and budget. *Environmental Pollution*, 88(1), 91-108.
- Witzel, L. D., and MacCrimmon, H. R. (1981). Role of gravel substrate on ova survival and alevin emergence of rainbow trout, *Salmo gairdneri*. *Canadian Journal of Zoology*, 59, 629-636.
- Wohl, N. E., and Carline. R. F. (1996). Relations among riparian grazing, sediment loads, macroinvertebrates, and fishes in three central Pennsylvania streams. *Canadian Journal of Fisheries and Aquatic Sciences*, 53(1) 260-266.
- Wood, P. J., and Armitage, P. D. (1997). Biological effects of fine sediment in the lotic environment. *Environmental Management*, 21(2), 203-217.

- Wu, F. C. (2000). Modeling embryo survival affected by sediment deposition into salmonid spawning gravels: Application to flushing flow prescriptions. *Water Resources Research*, 36(6), 1595-1603.
- Xue, N., Zhang, D., and Xu, X. (2006). Organochlorinated pesticide multiresidues in surface sediments from Beijing Guanting reservoir. *Water Research*, 40(2), 183-194.
- Yang, G.P. (2000). Polycyclic aromatic hydrocarbons in the sediments of the South China Sea. *Environmental Pollution*, 108, 163–171.
- Yang, R., Yao, T., Xu, B., Jiang, G., and Xin, X. (2007). Accumulation features of organochlorine pesticides and heavy metals in fish from high mountain lakes and Lhasa River in the Tibetan Plateau. *Environment international*, 33(2), 151-156.
- Young, M. K., Hubert, W., and Wesche, T. (1990a). Effect of substrate composition and stock origin on the survival to emergence of brown trout: A laboratory study. *Northwest Science*, 64(4), 224-231.
- Young, M. K., Hubert, W. A., and Wesche, T. A. (1990b). Fines in redds of large salmonids. *Transactions of the American Fisheries Society*, 119(1), 156-162.
- Younger, P. L., Mackey, R., and Connorton, B. J. (1993). Streambed sediment as a barrier to groundwater pollution: Insights from fieldwork and modelling in the River Thames basin. *Journal of Institution of Water and Environment Management*, 7, 577-585.
- Zakharov, V. M. (1992) Population phenogenetics: Analysis of developmental stability in natural populations. *Acta Zoologica Fennica*, 191, 7-30
- Zhang, W., Ye, Y., Tong, Y., Ou, L., Hu, D., and Wang, X. (2011). Modeling time-dependent toxicity to aquatic organisms from pulsed exposure of PAHs in urban road runoff. *Environmental Pollution*.
- Zhou, R., Zhu, L., Yang, K., and Chen, Y. (2006). Distribution of organochlorine pesticides in surface water and sediments from Qiantang River, East China. *Journal of Hazardous Material*, 137, 68-75.