

Palynological records of the Permian Ecca Group (South Africa): Utilizing climatic icehouse-greenhouse signals for cross basin correlations

Katrin Ruckwied¹, Annette E. Götz² and Phil Jones³

¹Shell International Exploration and Production, 200 North Dairy Ashford, Houston, Texas 77079, USA, E-mail:

katrin.ruckwied@shell.com

²Department of Geology, Rhodes University, Grahamstown 6140, P.O. Box 94, South Africa, E-mail:

a.gotz@ru.ac.za

³Ichron Ltd, Century House, Northwich, Cheshire, CW9 7TL, UK, E-mail: phil.jones@ichron.com

Abstract

The Permian formations of the South African Karoo Basin play a crucial role in understanding Gondwana's climate history during this time of major global changes. In this paper we present two data sets, one from the coal-bearing Vryheid Formation (Witbank Basin) and one from the Whitehill and Upper Prince Albert formations of the DP 1/78 core (NE Karoo).

Main goal was to study the vegetation changes during this period of global warming and test if the climatic signals could be used to correlate the basinal Ecca group facies with the fluvio-deltaic coal-bearing strata of the Witbank Basin. The palynological record of the No. 2 Coal Seam of the Vryheid Formation indicates a cold climate, fern wetland community, characteristic of lowland alluvial plains, and an upland conifer community in the lower part of the coal seam. Up section, these communities are replaced by a cool-temperate cycad-like lowland vegetation and gymnospermous upland flora. The data set of the DP 1/78 core is interpreted to represent a cool to warm temperate climate represented by a high amount of Gangamopteris and Glossopteris elements.

Both data sets are very different in their composition, which can be explained by the differences in depositional environment, however our findings reveal a different age of the studied assemblages and thus

also suggest that both data sets represent different stages in the transition from icehouse to greenhouse during Permian times. As the stratigraphic correlation between the Main Karoo Basin and the peripheral basins is still under discussion, this paper provides new data to underpin the stratigraphic placement of the Whitehill Formation relative to the coal-bearing Vryheid Formation.

Keywords: Palynology, Palaeoclimate, Permian, Karoo, South Africa

1. Introduction

Our knowledge of the Permian palynology of the South African Karoo Basin is based on fundamental research carried out in the 1970s and 1980s by Anderson (1977) and Falcon (see 1989 and references therein). No recent works address high-resolution palynostratigraphy of the Permian-Triassic coal-bearing formations in the South African Karoo, whereas new palynostratigraphic zonation schemes were established in other parts of southern Africa (D'Engelbronner 1996; Nyambe and Utting 1997; Stephenson and McLean 1999; Modie and Le Hérisse 2009). In general, Gondwanan land plant communities changed rapidly during Permian times due to the dramatic climate change subsequent to the melting of the Dwyka ice. This prominent change in vegetation, displayed in palynomorph assemblages, enables a high-resolution correlation.

From this background, it seems imperative to study the palynological record of the Permian succession in much more detail with respect to establish a high-resolution stratigraphic framework and climate history of the Karoo. Here, we report on new palynological data from the No. 2 Coal Seam of the northern Witbank coal field and a core (DP 1/78) drilled in the northeastern part of the Main Karoo Basin with the aim of using climatic signals for basin-wide correlation.

2. Geological setting

The Late Carboniferous to Middle Jurassic Karoo Basin covers approximately one third (i.e. 700 000 km²) of South Africa (Johnson et al., 2006), extending into Lesotho and in parts of Swaziland and Mozambique (Cole, 1992). A total thickness of 12 km sediment infill is reached within the southeastern part of the Main Karoo Basin. This sedimentary succession, known as the Karoo Supergroup, is subdivided into the Dwyka Group (Late Carboniferous (Pennsylvanian) to Early Permian (Artinskian); Visser, 1996), the Ecca Group of Permian age, the Permian-Triassic Beaufort Group (Johnson et al., 2006), and the Stormberg Group (Late Triassic to Early Jurassic; Catuneanu et al., 1998). The Karoo succession is capped by 1.4 km of basaltic lavas of the Drakensberg Group and intruded mafic dyke swarms (Veevers et al., 1994; Johnson et al., 1996).

The Karoo Basin forms part of a major series of Gondwanan foreland basins, including the Paraná Basin in South America, the Beacon Basin in Antarctica, and the Bowen Basin in Australia (De Wit and Ransome, 1992; Veevers et al., 1994; Catuneanu et al., 1998; Catuneanu and Elango, 2001; Catuneanu and Bowker, 2002). These basins formed mainly as result of collision and terrain accretion tectonics along the southern edge margin of Gondwana (De Wit and Ransome, 1992; Veevers et al., 1994). Recent interpretations of the basin evolution and tectonic setting of the Karoo range from a retro-arc foreland basin (Johnson et al., 2006), a transtensional foreland system created by subsidence and tilting in a strike-slip regime (Tankard et al., 2009) to a thin-skinned fold belt that developed from collisional tectonics and distant subduction to the south (Lindeque et al., 2011).

The Karoo Basin hosts important coal resources of South Africa (Johnson et al., 1997; Cairncross, 2001). The Witbank coal field comprises the basin's northeastern and most economic coal deposits (Snyman, 1998) with 5 coal seams (coal seams 1 – 5 in ascending order) that belong stratigraphically to the Vryheid Formation. One of the most important coal seams of the Witbank coal field, the No. 2 Coal Seam, represents postglacial fluvio-deltaic deposits of the Lower Ecca Group, pointing to a highly proximal setting of the northern basin margin (Falcon, 1989). Further south in the Main Karoo Basin organic-rich

shales of the Whitehill Formation have been regarded as time-equivalent to the coal-bearing Vryheid Formation (Viljoen, 1994; Johnson et al. 1996, 1997).

3. Material and methods

During a field campaign in 2011, we sampled an eight meter thick succession of the No. 2 Coal Seam exposed in the Inyanda Coal Mine south of Witbank (Fig. 1) to study palynofacies and palynostratigraphy. Lithologies of 22 samples include fine- to medium-grained sandstones, organic-rich siltstones and coals. We also present here data from the DP 1/78 core (Fig. 1) that was sampled during a data gathering campaign by Shell Exploration & Production. In total, 11 samples of a black shale interval have been analysed.

For palynofacies analysis samples were prepared using standard palynological processing techniques, including HCl (33%) and HF (73%) treatment for dissolution of carbonates and silicates, and saturated ZnCl₂ solution (D≈2.2 g/ml) for density separation. Residues were sieved at 15 µm mesh size. Slides have been mounted in Eukitt, a commercial, resin-based mounting medium.

Sedimentary organic matter is grouped into sporomorphs, fresh water algae, acritarchs, degraded organic matter, amorphous organic matter and phytoclasts. The phytoclast group has been subdivided into opaque and translucent particles. Equidimensional and needle-shaped phytoclasts have been counted separately. The relative percentage of these components is based on counting at least 300 particles per slide.

4. Sedimentology, palynofacies and palaeoenvironmental interpretation

The northern Witbank coal field is characterized by proximal deposits including coarse-grained to pebbly sandstones, partially with an abrupt upward transition into fine-grained sediments and coal, trough cross-stratified medium- to coarse-grained sandstones, and horizontally laminated, fine- to medium-grained sandstones and organic-rich siltstones. An up to 1 m thick sandstone marker bed separates the No. 2 Coal

Seam into a Lower and Upper Coal Seam. In this fluvial-alluvial depositional setting, peat swamps developed on broad abandoned alluvial plains.

Sedimentary organic matter of these proximal deposits is highly degraded and poorly preserved, however relative abundance changes in the palynomorph assemblage and variations in the amount and size and shape of plant debris are clearly documented. Generally, palynofacies is dominated by a high amount of opaque and translucent phytoclasts showing a high variability in size and shape throughout the section.

The lower part of the section (Lower Coal Seam) is characterized by a high amount of opaque particles, whereas the upper part (Upper Coal Seam) comprises high amount of translucent plant debris.

Amorphous Organic Matter (AOM) is characteristic of laminated, organic-rich siltstones and coals.

Degraded organic matter (DOM) is preserved in all sampled lithologies. Freshwater algae (*Tetraporina* spp., *Botryococcus* spp.) are common in the Upper Coal Seam. The lower part of the section (Lower Coal Seam) is dominated by horsetail and fern spores and gymnospermous monosaccate pollen grains whereas the upper part of the section (Upper Coal Seam) is totally dominated by gymnospermous bisaccate (taeniate and non-taeniate) pollen grains with only very few fern spores preserved. The sporomorph assemblages indicate a fern wetland community, characteristic of lowland alluvial plains, and an upland conifer community dominated by monosaccate-producers in the lower part of the coal seam. Up section, these communities are replaced by a cycad-like and lycopsid lowland vegetation and a gymnospermous upland flora of bisaccate-producers, indicating a more temperate climate than the cooler floras in the lower part (Götz and Ruckwied, 2013).

The sampled interval of the DP1/78 well is composed of organic-rich, carbonaceous black shales. The lower part of this interval is interpreted to be part of the Prince Albert Formation and contains a higher amount of silt than the upper part of the interval, which is interpreted to represent deposits of the Whitehill Formation. Palynofacies of the Prince Albert Formation is characterized by rich bleached kerogens with abundant and diverse palynomorphs. Palynomorphs comprise approximately 85% of the sedimentary organic matter. Bisaccate pollen grains dominate the assemblage. Though fully marine

indications were not recorded, the assemblages are considered to have been deposited in a distal subaqueous, probably marine environment with ventilated bottom waters.

Samples at the base of the Whitehill Formation are characterized by the frequent occurrence of *Leiosphaeridia* spp. and peak abundance of bisaccate pollen grains including *Vittatina* spp. and *Vittatina saccata*. Such an assemblage together with the occurrence of AOM is interpreted to indicate a distal (restricted) marine environment.

A decrease up-hole in abundance of leiospheres together with an influx of the freshwater/brackish algae *Botryococcus* spp. tentatively indicates an increase in terrestrial (freshwater) influence. This interpretation is supported by the increase (up-hole) in spore types including *Verrucosisporites andersonii*, *Anapiculatisporites* spp. and *Cyclogranisporites* spp., and a general increase in phytoclasts. These particles are thought to be deposited more proximally to source than airborne bisaccate pollen and thus indicate a landward shift in depositional environment. The presence of rare specimens of *Cymatiosphaera gondwanensis* is suggestive of (weak?) marine influences. In summary, the Whitehill Formation therefore is interpreted to represent a distal marine setting with increasing proximity to shore and terrestrial (freshwater) influences up-hole. The high amount of striate and non-striate bisaccates is indicative for a warmer climate during time of deposition relative to the No. 2 Coal Seam assemblages, which imply a cool to temperate climate.

5. Palynostratigraphy

Palynostratigraphy of the Witbank sub-basin and the Northern Main Karoo Basin is well understood. Two schemes have been published by Anderson (1977) and Falcon et al. (1984). In the present study, we focus on the latter (Fig. 2), as it was generated on data sets very close to the Inyanda coal mine, which is analysed herein. To date, there are no palynological data published from the Whitehill Formation, the relatively high maturity in wide parts of the Main Karoo Basin prevents to apply this method further towards the basin centre and the southern parts of the basin.

The palynological data from the Witbank No. 2 Coal Seam document a climatic shift from cold to cool-temperate conditions. A postglacial monosaccate-producing flora is replaced by a warmer climate bisaccate-producing plant community. The lower part of the No. 2 Coal Seam (Lower Coal Seam) is dominated by *Virkkipollenites* spp., *Potonieisporites* spp. and *Plicatipollenites*, and horsetail and fern spores such as *Calamospora* spp., *Microbaculispora* spp., *Punctatisporites* spp., *Verrucosisporites* spp. The additional presence of *Florinites* spp. places this assemblage into the Virkkipollenites-Plicatipollenites-Potonieisporites-Florinites sub-assemblage, i.e. Biozone Ic (Falcon et al., 1984). The assemblage of the Upper Coal Seam is dominated by *Sulcatisporites* spp., *Protohaploxypinus* spp., and *Alisporites* spp., with monosaccate elements still present. This part of the seam is within the Sulcatisporites-Virkkipollenites-Plicatipollenites-Potonieisporites sub-assemblage. The presence of key marker species *Striatoabieites multistriatus*, *Protohaploxypinus limpidus* and *Sulcatisporites splendens* implies a correlation with Falcon's Biozone IId (Fig. 3).

Samples of the upper Prince Albert Formation in the DP 1/78 well are dominated by non-striate bisaccate pollen grains, but striate pollen grains, mainly *Protohaploxypinus* spp. and *Striatoabieites* spp. are common. The frequent occurrence of *Lundbladispora braziliensis* as well as the presence of *Marsupipollenites striatus* places this interval into Falcon's Biozone IIIf, i.e. the Sulcatisporites-Protohaploxypinus sub-assemblage (Fig. 4). Samples of the Whitehill Formation are characterized by a high amount of *Vittatina* spp., the general increase of the striate bisaccate group, the consistent presence of *Guttulapollenites* aff. *hanonicus* and *Lueckisporites virkkiae* suggest to place this sediments into the Protohaploxypinus-Vittatina-Guttulapollenites-Lueckisporites sub-assemblage, i.e. biozones IVg and IVh (Falcon et al., 1984).

6. Age Interpretation, correlation and conclusions

The Vryheid Formation was previously dated as Artinskian, with possible extensions into the Kungurian by various authors (McLachlan and Anderson, 1973; MacRae, 1988; Catuneanu et al., 2002). In general

the authors stated that the Vryheid Formation was deposited during a more extended period of time and that deposition continued at least during the early Middle Permian. Floral composition of the lower two coal seams (No. 1 and No. 2) clearly show a Lower Permian signature which matches with the previous age interpretation, however, the upper coal seams (No. 3, 4, and 5), especially coal seam No. 5, clearly indicate a Middle Permian age. Samples of the Inyanda Coal Mine from the No. 2 Coal Seam are very comparable to the assemblages described by Falcon et al. (1984). However, the samples of the Lower Coal Seam (No. 2) show clearly the signature of Falcon's Biozone Ic. This differs from the results of Falcon's original work where all data from the No. 2 Coal Seam match with Biozone IId. Due to the fact that palynofacies suggest a swamp dominated palaeoenvironment, the authors assume that a diachronous sedimentation of the coals within the Witbank Basin is most likely the explanation for this difference. The age interpretation for these samples is Artinskian to Kungurian. Data from the DP 1/78 well represent the first palynological data for the upper Prince Albert Formation and Whitehill Formation. Samples of the Prince Albert Formation show a very similar composition as samples from the No. 4 Coal Seam of the Witbank Basin (Falcon et al., 1984). Retallack (2006) reports *Lundbladispora braziliensis* not above the Guadalupian. Souza et al. (2005) describe a similar assemblage from the Paraná Basin in Brazil. Their *Vittatina costabilis* interval is defined as a zone with the first appearance of *Vittatina* spp. at the base and the first appearance of *Lueckisporites virkkiae* at the top. Due to the similarities to this assemblage we suggest to place the Prince Albert Formation chronostratigraphically into the uppermost Early Permian to lowermost Middle Permian. Samples from the Whitehill Formation seem to yield similar palynomorph assemblages as described from the No. 5 Coal Seam of the Witbank Basin (Falcon et al., 1984). The presence of *Lueckisporites virkkiae* suggests a correlation with the *Lueckisporites virkkiae* Interval zone of the Brazilian Paraná Basin (Souza et al., 2005), which is interpreted to be Guadalupian in age. An absolute age obtained from a tuff in the overlying Collingham Formation in the Namibian part of the Karoo, dated as 270 My (Stollhofen et al., 2000), supports a Roadian age.

The comparison of the two assemblages from the Witbank Basin and the DP 1/78 well strongly suggests that the sediments of the Vryheid Formation and the uppermost Prince Albert Formation and Whitehill

Formation have been deposited at the same time (Fig. 5). This agrees with previous correlations by Viljoen (1994) and Johnson et al. (1996, 1997), but disagrees with Catuneanu et al. (2002) who suggested that the Whitehill Formation is significantly older and equivalent to the Pietermaritzburg Formation.

Acknowledgements

We kindly acknowledge Exxaro Resources, South Africa for giving us permission to sample the No. 2 Coal Seam exposed in the Inyanda Mine. We thank Stuart Clague (Exxaro) and Wlady Altermann (University of Pretoria) who encouraged us to work on the South African Gondwana coals and supported our field campaign in 2011. The authors wish to thank the Council for Geoscience (CGS) of South Africa for the supply of well DP 1/78 material. We also want to thank Shell Exploration and Production for permission to publish data of the DP 1/78 well, especially Manuel Poupon and Hermann Läufer. This work is based on the research supported in part by the National Research Foundation of South Africa (Grant No. 85354). The comments of two anonymous reviewers greatly improved the manuscript.

References

- Anderson, J.M. 1977. The biostratigraphy of the Permian and Triassic. Part 3. A review of Gondwana palynology with particular reference to the northern Karoo Basin, South Africa. *Memoirs of the Botanical Survey of South Africa* 4, 1–33.
- Cairncross, B. 2001. An overview of the Permian (Karoo) coal deposits of southern Africa. *Journal of African Earth Sciences* 33, 529–562.
- Catuneanu, O., Elango, H.N. 2001. Tectonic control on fluvial style: the Balfour Formation of the Karoo Basin, South Africa. *Sedimentary Geology* 140, 291–313.
- Catuneanu, O., Bowker, D. 2002. Sequence stratigraphy of the Koonap and Middleton fluvial formations in the Karoo foredeep, South Africa. *Journal of African Earth Sciences* 33, 579–595.

- Catuneanu, O., Hancox, P.J., Rubidge, B.S. 1998. Reciprocal flexural behaviour and contrasting stratigraphies: a new basin development model for the Karoo retroarc foreland system, South Africa. *Basin Research* 10, 41–439.
- Catuneanu, O., Wopfner, H., Eriksson, P.G., Cairncross, B., Rubidge, B.S., Smith, R.M.H., Hancox, P.J. 2005. The Karoo basins of south-central Africa. *Journal of African Earth Sciences* 43, 211–253.
- Cole, D.I. 1992. Evolution and development of the Karoo Basin. In: De Wit, M.J., Ransome, I.G.D. (Eds.), *Inversion Tectonics of the Cape Fold Belt, Karoo and Cretaceous Basins of Southern Africa*. A.A. Balkema, Rotterdam, pp. 87–99.
- De Wit, M.J., Ransome, I.G.D. 1992. Regional inversion tectonics along the southern margin of Gondwana. In: De Wit, M.J., Ransome, I.G.D. (Eds.), *Inversion Tectonics of the Cape Fold Belt, Karoo and Cretaceous Basins of Southern Africa*. A.A. Balkema, Rotterdam, pp. 15–21.
- D’Engelbronner, E.R. 1996. New palynological data from Karoo sediments, Mana Pools basin, northern Zimbabwe. *Journal of African Earth Sciences* 23(1),17–30.
- Falcon, R.M.S 1986. A brief review of the origin, formation and distribution of coal in Southern Africa. In: Anhaeusser, C.R., Maske, S. (Eds.), *Mineral Deposits of Southern Africa*. Geological Society of South Africa 2, 1879–1898.
- Falcon, R.M.S. 1989. Macro and micro-factors affecting coal-seam quality and distribution in southern Africa with particular reference to the No. 2 seam, Witbank Coalfield, South Africa. *International Journal of Coal Geology* 12, 681–731.
- Falcon, R.M.S., Pinheiro, H., Sheperd, P. 1984. The palynobiostratigraphy of the major coal seams in the Witbank Basin with lithostratigraphic, chronostratigraphic and palaeoclimatic implications. *Comunicações dos Serviços Geológicos de Portugal* 70, 215–243.
- Johnson, M.R., Van Vuuren, C.J., Hegenberger, W.F., Key, R., Shoko, U. 1996. Stratigraphy of the Karoo Supergroup in southern Africa: an overview. *Journal of African Earth Sciences* 23(1), 3–15.

- Johnson, M.R., Van Vuuren, C.J., Visser, J.N.J., Cole, D.I., Wickens, H. de V., Christie, A.D.M., Roberts, D.L. 1997. The foreland Karoo Basin, South Africa. In: Selley, R.C. (Ed.), *Sedimentary Basins of Africa*. Elsevier, Amsterdam, pp. 269–317.
- Johnson, M.R., Van Vuuren, C.J., Visser, J.N.J., Cole, D.I., Wickens, H. de V., Christie, A.D.M., Roberts, D.L., Brandl, G. 2006. Sedimentary rocks of the Karoo Supergroup. In: Johnson, M.R., Anhaeusser, C.R., Thomas, R.J. (Eds.), *The Geology of South Africa*. Geological Society of South Africa, Council for Geoscience, Pretoria, pp. 461–499.
- Lindeque, A., De Wit, M.J., Ryberg, T., Weber, M., Chevallier, L. 2011. Deep crustal profile across the southern Karoo Basin and Beattie Magnetic Anomaly, South Africa : An integrated interpretation with tectonic implications. *South African Journal of Geology* 114(3/4), 265–292.
- McLachlan, I.R., Anderson, A. 1973. A review of the evidence for marine conditions in southern Africa during Dwyka times. *Palaeontologia Africana*, 15, 38–64.
- MacRae, C.S. 1988. Palynostratigraphical correlation between the Lower Karoo sequence of the Waterburg and Pafuri coal basins and the Hammanskraal plant macrofossil locality, Republic of South Africa. *Memoirs of the Geological Survey of South Africa* 75, 1–217.
- Modie, B.N., Le Hérisse, A. 2009. Late Palaeozoic palynomorph assemblages from the Karoo Supergroup and their potential for biostratigraphic correlation, Kalahari Karoo Basin, Botswana. *Bulletin of Geosciences* 84(2), 337–358.
- Nyambe, I.A., Utting, J. 1997. Stratigraphy and palynostratigraphy, Karoo Supergroup (Permian and Triassic), mid-Zambezi Valley, southern Zambia. *Journal of African Earth Sciences* 24(4), 563–583.
- Retallack, G.J., Metzger, C.A., Greaver, T., Jahren, A.H., Smith, R.M.H., Sheldon, N.H. 2006. Middle-Late Permian mass extinction on land. *GSA Bulletin* 118 (11-12), 1398–1411.
- Snyman, C.P. 1998. Coal. In: Wilson, M.G.C., Anhaeusser, C.R. (Eds.), *The Mineral Resources of South Africa*. Council for Geoscience Handbook 16, 136–205.
- Souza, P.A., Merques-Toigo, M. 2005. Progress on the palynostratigraphy of the Permian strata in Rio Grande do Sul State, Paraná basin, Brazil. *Anais da Academia brasileira de Ciencias* 77 (2), 353–365.

- Stephenson, M.H., McLean, D. 1999. International correlation of Early Permian palynofloras from the Karoo sediments of Morupule, Botswana. *South African Journal of Geology* 102, 3–14.
- Stollhofen, H., Stanistreet, I.G., Bangert, B., Grill, H. 2000. Tuffs, tectonism and glacially related sea-level changes, Carboniferous-Permian, southern Namibia. *Palaeogeography, Palaeoclimatology, Palaeoecology* 161, 127–150.
- Tankard, A., Welsink, H., Aukes, P., Newton, R., Stettler, E. 2009. Tectonic evolution of the Cape and Karoo basins of South Africa. *Marine and Petroleum Geology* 26(8), 1379–1412.
- Veevers, J.J., Cole, D.I., Cowan, E.J. 1994. Southern Africa: Karoo Basin and Cape Fold Belt. In: Veevers, J.J., Powell, C.McA. (Eds.), *Permian-Triassic Pangean Basins and fold belts along the Panthalassan Margin of Gondwanaland*. GSA Memoir 184, 223–279.
- Viljoen, M.J., 1994, A review of regional variations in facies and grade distribution of the Merensky Reef, western Bushveld Complex, with some mining implications. *Proceedings of the 15th CMMI Congress, SAIMM*, 183–194.
- Visser, J.N.J. 1996. Controls on Early Permian shelf deglaciation in the Karoo Basin of South Africa. *Palaeogeography, Palaeoclimatology, Palaeoecology* 125, 129–139.

Figure captions

Figure 1. Location of the studied well section (DP 1/78) and the No. 2 Coal Seam exposed in the Inyanda mine (Witbank). Map modified from Johnson et al. (2006).

Figure 2. Palynostratigraphy of the coal-bearing Ecca Group (Lower-Middle Permian) in the northeastern Main Karoo Basin (after Falcon et al., 1984).

Figure 3. Palynomorph distribution chart of the No. 2 Coal Seam, Inyanda Mine (Witbank). Biozones Ic and IId after Falcon et al. (1984).

Figure 4. Palynomorph distribution chart of the DP 1/78 core (NE Karoo). Biozones IIIf and IVg/h after Falcon et al. (1984).

Figure 5. Correlation of the Prince Albert and Whitehill formations (N Main Karoo) and Vryheid Formation (Witbank Basin, NE Karoo) based on new palynological data from the DP 1/78 well and the No. 2 Coal Seam.




Figure 1




Figure 2




Figure 3




Figure 4

N Karoo (DP 1/78)	Biozones Falcon et al. (1984)	NE Karoo (Witbank)
Tierberg Fm.	IV	h ¹ Coal Seam No. 6
Whitehill Fm.		h Coal Seam No. 5
Prince Albert Fm.	III	g Coal Seam No. 4
		f Coal Seam No. 3
	II	e Coal Seam No. 2
		d Coal Seam No. 1
I	c	
	b	
a		
?		Pieter-maritzburg Fm.
Dwyka Group		

Vryheid Fm.

Figure 5