

Testing gravity at large scales with H I intensity mapping

Alkistis Pourtsidou^{*}

Institute of Cosmology & Gravitation, University of Portsmouth, Burnaby Road, Portsmouth PO1 3FX, UK

Accepted 2016 June 9. Received 2016 June 9; in original form 2015 December 14

ABSTRACT

We investigate the possibility of testing *Einstein's* general theory of relativity (GR) and the standard cosmological model via the E_G statistic using neutral hydrogen (H I) intensity mapping. We generalize the Fourier space estimator for E_G to include H I as a biased tracer of matter and forecast statistical errors using H I clustering and lensing surveys that can be performed in the near future, in combination with ongoing and forthcoming optical galaxy and cosmic microwave background (CMB) surveys. We find that fractional errors < 1 per cent in the E_G measurement can be achieved in a number of cases and compare the ability of various survey combinations to differentiate between GR and specific modified gravity models. Measuring E_G with intensity mapping and the Square Kilometre Array can provide exquisite tests of gravity at cosmological scales.

Key words: gravitational lensing; weak – cosmology; observations – cosmology; theory – large-scale structure of Universe.

1 INTRODUCTION

During the last two decades, observational cosmology has entered an era of unprecedented precision. The standard cosmological model (Λ CDM) fits the data extremely well, but requires that General Relativity (GR) is the correct description of gravity on all scales and that the matter-density of our Universe is dominated by the two constituents of the dark sector, i.e. dark energy in the form of a cosmological constant and cold dark matter. Dark energy is thought to be responsible for the accelerated expansion of the Universe (Riess et al. 1998; Perlmutter et al. 1999) and uncovering its nature is arguably the most exciting challenge in modern cosmology. Alternative explanations to the cosmological constant have been proposed, for example a dynamically evolving scalar field playing the role of dark energy (see Copeland, Sami & Tsujikawa 2006, for a review). A different point of view suggests that late time cosmic acceleration could be due to modifications to the laws of gravity on the largest (cosmological) scales [see Clifton et al. (2012) for a review]. In general, exotic dark energy and modified gravity (MG) theories modify the background and perturbation evolution and dynamics of the Universe and they have distinct and detectable observational effects.

In this paper we are going to investigate the possibility of testing Λ CDM and the laws of gravity at large scales using the E_G statistic, which was first introduced in Zhang et al. (2007). The definition of E_G in Fourier space is

$$E_G(k, z) = \frac{c^2 k^2 (\phi - \psi)}{3H_0^2 (1+z)\theta(k)}, \quad (1)$$

where (ϕ, ψ) are the scalar potentials in the perturbed Friedmann–Robertson–Walker (FRW) metric $ds^2 = (1 + 2\psi)dt^2 - a^2(1 + 2\phi)d\mathbf{x}^2$, $\theta \equiv \nabla \cdot \mathbf{v}/H(z)$ is the peculiar velocity perturbation field, and H_0 is the value of the Hubble parameter today.

From the above definition it is clear that E_G depends on how gravity behaves on large scales. In GR, assuming that the background Universe is described by a flat FRW metric and in the absence of anisotropic stress, we can show that the Poisson and anisotropy equations can be written as

$$\begin{aligned} k^2 \psi &= -4\pi G a^2 \rho \delta \\ \phi &= -\psi, \end{aligned} \quad (2)$$

where a is the scale factor, ρ is the background matter density and δ is the matter density perturbation. In MG we can use two scale- and time-dependent functions $\mu(k, a)$ and $\gamma(k, a)$ to parametrize possible departures from Λ CDM and write (Hojjati, Pogosian & Zhao 2011)

$$\begin{aligned} k^2 \psi &= -4\pi G a^2 \mu(k, a) \rho \delta \\ \phi &= -\gamma(k, a) \psi. \end{aligned} \quad (3)$$

Substituting these expressions in the E_G definition (1) and using the fact that on linear scales we can write $\theta = f\delta$, where f is the linear growth rate, we find (Pullen, Alam & Ho 2015a)

$$E_G(k, z) = \frac{\Omega_{m,0} \mu(k, a) [1 + \gamma(k, a)]}{2f}. \quad (4)$$

For GR, we set $\mu = \gamma = 1$ and find $E_G(k, z) = \Omega_{m,0}/f(z)$, with $\Omega_{m,0}$ the matter density today relative to the critical density.

In order to measure E_G , the estimator in Zhang et al. (2007) involved the ratio between the cross-correlation power spectrum

* E-mail: alkistis.pourtsidou@port.ac.uk

of galaxies and weak lensing convergence $P_{g\kappa}$ and the cross-correlation power spectrum of galaxies and velocities $P_{g\theta}$. The latter is equivalent to the galaxy autocorrelation times the redshift-space distortion (RSD) parameter β , i.e. βP_{gg} , where $\beta = f/b_g$ with b_g the galaxy clustering bias. In Reyes et al. (2010) and Blake et al. (2015) E_G was measured using a real space estimator and the results were fully consistent with GR. Recently, forecasts for future galaxy surveys were presented in Leonard, Ferreira & Heymans (2015).

In Pullen et al. (2015a), a Fourier estimator for E_G was constructed in terms of the galaxy- cosmic microwave background (CMB) lensing and galaxy clustering angular power spectra, also including the RSD parameter β . Very recently, they reported the largest-scale measurement of E_G using the aforementioned method; they found $E_G(z = 0.57) = 0.243 \pm 0.060$ (stat) ± 0.013 (sys), a result in 2.6σ tension with the GR prediction at this redshift (Pullen et al. 2015b).

In the next section we will closely follow the approach by Pullen et al. (2015a) and generalize their E_G Fourier space estimator to include a different dark matter tracer than galaxies, namely neutral hydrogen (H I). In Section 3 we describe the various optical galaxy, intensity mapping (IM) and CMB surveys we are going to use to forecast future E_G measurements. We present our forecasts in Section 4 and demonstrate how they can be used to discriminate between GR and specific MG models. We conclude in Section 5.

2 FORMALISM

The Fourier space estimator for E_G is constructed as (Pullen et al. 2015a)

$$\hat{E}_G(\ell, \bar{z}) = \frac{c^2 \hat{C}_\ell^{g\kappa}}{3H_0^2 \hat{C}_\ell^{g\theta}}, \quad (5)$$

and it can be expressed in terms of the galaxy-convergence angular cross-power spectrum $C_\ell^{g\kappa}$, the galaxy angular auto-power spectrum C_ℓ^{gg} , and the RSD parameter $\beta = f/b_g$ (see also Reyes et al. (2010)). Note that it is galaxy bias free in the linear regime, which evades the issue of possible degeneracies between MG effects and bias.

In this work, we are going to utilize the above estimator but using H I instead of galaxies. We will forecast statistical errors for E_G using H I IM clustering and lensing surveys which can be performed in the near future using the Square Kilometre Array (SKA).¹ Intensity mapping (Battye, Davies & Weller 2004; Chang et al. 2008; Loeb & Wyithe 2008; Mao et al. 2008; Peterson et al. 2009; Seo et al. 2010; Ansari et al. 2012; Battye et al. 2013; Switzer et al. 2013; Bull et al. 2015) is an innovative technique which uses H I to map the large-scale structure of the Universe in three dimensions. Instead of detecting individual galaxies like the conventional galaxy surveys, intensity mapping surveys use H I as a dark matter tracer by measuring the intensity of the redshifted 21-cm line across the sky and along redshift, treating the 21-cm sky as a diffuse background.

Using H I instead of galaxies we can write the Fourier space E_G estimator as

$$\hat{E}_G(\ell, \bar{z}) = \frac{c^2 \hat{C}_\ell^{\delta_{H1}\kappa}}{3H_0^2 \hat{C}_\ell^{\delta_{H1}\theta}}, \quad (6)$$

where δ_{H1} is the H I density contrast which traces the matter density as $\delta_{H1} = b_{H1}\delta$, with b_{H1} the H I bias. The H I-convergence angular cross-power spectrum using the Limber approximation (Limber

1954) for scales $\ell \geq 10$ is given by

$$C_\ell^{\delta_{H1}\kappa} = \frac{3\Omega_{m,0}H_0^2}{2c^2} b_{H1}(\bar{\chi})(1 + \bar{z}) P_{\delta\delta}(\ell/\bar{\chi}, \bar{\chi}) \bar{\chi}^{-1} (1 - \bar{\chi}/\chi_s), \quad (7)$$

where $\bar{\chi}$ is the comoving radial distance to redshift \bar{z} , χ_s is the comoving radial distance to the background sources, and $P_{\delta\delta}$ is the matter power spectrum. When deriving this formula we assumed that both the foreground lenses and the background source distributions can be approximated by delta functions. This is going to be a valid assumption in the cases we are going to study, as our chosen foreground redshift bins are always going to be small enough ($\Delta z = 0.1$), and for the background sources we use either the CMB plane or 21-cm sources, both of which can be well approximated by a delta function distribution.

Then, following the formalism Pullen et al. (2015a) used for the galaxy clustering case, we construct the velocity-H I angular cross-power spectrum, to be

$$C_\ell^{\delta_{H1}\theta} = \frac{\bar{\chi}(1 - \bar{\chi}/\chi_s)\beta(\bar{z})(1 + \bar{z})}{2\Delta\chi^{-1}} \hat{C}_\ell^{\delta_{H1}\delta_{H1}}, \quad (8)$$

with $\Delta\chi$ the comoving width of the redshift bin with central redshift \bar{z} , and

$$C_\ell^{\delta_{H1}\delta_{H1}} = \frac{1}{\Delta\chi} \bar{\chi}^{-2} b_{H1}^2(\bar{\chi}) P_{\delta\delta}(\ell/\bar{\chi}, \bar{\chi}). \quad (9)$$

Note that the RSD parameter is $\beta = f/b_{H1}$.

The fractional error of E_G can be written as

$$\frac{\sigma^2[E_G(\ell, \bar{z})]}{E_G^2} = \left[\left(\frac{\sigma(C_\ell^{\delta_{H1}\kappa})}{C_\ell^{\delta_{H1}\kappa}} \right)^2 + \left(\frac{\sigma(\beta)}{\beta} \right)^2 + \left(\frac{\sigma(C_\ell^{\delta_{H1}\delta_{H1}})}{C_\ell^{\delta_{H1}\delta_{H1}}} \right)^2 \right]. \quad (10)$$

The error in the measurement of $C_\ell^{\delta_{H1}\kappa}$ is

$$\sigma^2(C_\ell^{\delta_{H1}\kappa}) = \frac{(C_\ell^{\delta_{H1}\kappa})^2 + (C_\ell^{\kappa\kappa} + N_\ell^{\kappa\kappa})(C_\ell^{\delta_{H1}\delta_{H1}} + N_\ell^{\delta_{H1}\delta_{H1}})}{(2\ell + 1)f_{\text{sky}}}, \quad (11)$$

where $C_\ell^{\kappa\kappa}$ is the lensing convergence power spectrum, $N_\ell^{\kappa\kappa}$ the lensing reconstruction noise, $N_\ell^{\delta_{H1}\delta_{H1}}$ the noise in the measurement of the H I clustering angular power spectrum $C_\ell^{\delta_{H1}\delta_{H1}}$, and f_{sky} the (overlapping) fraction of the sky scanned by the surveys used.

Note that in the following we are also going to consider combinations of galaxy-CMB lensing surveys, like the ones presented in Pullen et al. (2015a). The formulae for this case can be recovered by setting $\delta_{H1} \rightarrow \delta_g$ and $b_{H1} \rightarrow b_g$ in the above equations. We should also mention that equation (10) is exact only if β and $C_\ell^{\delta_{H1}\delta_{H1}}$ (or C_ℓ^{gg} in the optical galaxy case) are measured by different, independent surveys – otherwise they should be correlated. However, we can safely use equation (10) for the cases we are going to consider in this paper. That is because the dominant errors are the lensing and β ones, and the clustering error due to thermal noise (shot noise) for the upcoming IM (photometric galaxy) surveys we will use is very small so it can be neglected. Thus, the combined (βC^{H1-H1}) error increases only slightly via the covariance and our results are practically unchanged [see Pullen et al. (2015a) for a detailed discussion of this issue for the optical galaxy case].

Before we move on to our forecasts, we will dedicate the next section to the specifications and noise properties of the various surveys we are going to use.

¹ www.skatelescope.org

3 THE SURVEYS

3.1 Galaxy and H I IM clustering

For our forecasts involving galaxy clustering we will consider two photometric surveys: the ongoing Dark Energy Survey (DES)² and the planned Large Synoptic Survey Telescope (LSST).³ Both of them aim to investigate the nature of cosmic acceleration and are able to perform precision galaxy clustering measurements.

The DES survey parameters are $A_{\text{sky}} = 5000 \text{ deg}^2$ (equivalently, $f_{\text{sky}} \simeq 0.1$), number density of galaxies $n_g = 10 \text{ arcmin}^{-2}$, and redshift range $0 < z < 2$ with median redshift $z_0 = 0.7$ (Becker et al. 2015). The LSST survey parameters are assumed to be $f_{\text{sky}} = 0.5$, number density of galaxies $n_g = 40 \text{ arcmin}^{-2}$, and redshift range $0 < z < 2.5$ with median redshift $z_0 = 1$ (Abell et al. 2009). For our forecasts we model the redshift distribution of galaxies as

$$\frac{dn}{dz} \propto z^2 \exp[-(z/z_0)^{3/2}] \quad (12)$$

and use $b_g(z) = \sqrt{1+z}$ for the galaxy clustering bias. The noise term $N_{\ell}^{\delta_g \delta_g} = 1/\bar{n}_i$, where \bar{n}_i is the number of galaxies per steradian in the i th redshift bin.

For H I clustering using the IM technique we will consider the SKA_Mid instrument. This can operate in two observing modes, the single-dish (autocorrelation) mode and the interferometer mode. Their noise properties have been described in detail in Bull et al. (2015) and Pourtsidou et al. (2016), so we are just going to state the relevant formulae here. The thermal noise angular power spectrum for the single-dish mode is given by

$$C_{\ell}^{\text{N}} = \Omega_{\text{pix}} (\sigma_{\text{pix}})^2 \exp[\ell(\ell+1)(\theta_{\text{B}}/\sqrt{8\ln 2})^2], \quad (13)$$

where the pixel thermal noise is $\sigma_{\text{pix}} = T_{\text{sys}}/\sqrt{2Bt_{\text{obs}}}$, and $\Omega_{\text{pix}} \simeq 1.13\theta_{\text{B}}^2$. T_{sys} is the system temperature, B the bandwidth of observation, t_{obs} the observation time, and $\theta_{\text{B}} \sim \lambda/D_{\text{dish}}$ the beam full width at half-maximum of a dish with diameter D_{dish} at wavelength λ . The observation time is given by $t_{\text{obs}} = t_{\text{tot}}(\Omega_{\text{pix}}/\Omega_{\text{tot}})N_{\text{dishes}}$, where t_{tot} is the total survey time, Ω_{tot} is the sky area the survey scans, and N_{dishes} is the number of available dishes. For the interferometer mode the thermal noise power spectrum is given by (Pourtsidou et al. 2016)

$$C_{\ell}^{\text{N}} = \frac{T_{\text{sys}}^2 [\text{FOV}]^2}{B t_{\text{tot}} n(\ell)} \quad (14)$$

with $\text{FOV} \simeq (\lambda/D_{\text{dish}})^2$ and $n(\ell)$ the number density of baselines.

The H I clustering noise term will be calculated as

$$N_{\ell}^{\delta_{\text{H I}} \delta_{\text{H I}}} = \frac{C_{\ell}^{\text{N}}}{\bar{T}^2}, \quad (15)$$

where the mean brightness temperature $\bar{T}(z)$ is given by

$$\bar{T}(z) = 180 \frac{\Omega_{\text{H I}}(z) h(1+z)^2}{H(z)/H_0} \text{ mK}. \quad (16)$$

For our forecasts we will assume that the H I density evolves with redshift as $\Omega_{\text{H I}}(z) = 4 \times 10^{-4} (1+z)^{0.6}$ which has been suggested in Crighton et al. (2015). We are also going to use the H I bias model $b_{\text{H I}}(z)$ from Camera et al. (2013) and consider Phase 1 of the SKA_Mid instrument (SKA1_Mid), consisting of 130 dishes with 15-m diameter according to the recently updated specifications ('re-baselining') in order to meet budget constraints (McPherson 2015).

² <http://www.darkenergysurvey.org/>

³ <http://www.lsst.org>

The redshift range is $0.35 < z < 3$ (Band 1). The system temperature is given by (Dewdney 2013)

$$T_{\text{sys}} = 28 + 66 \left(\frac{\nu}{300 \text{ MHz}} \right)^{-2.55} \text{ K}, \quad (17)$$

with ν the observing frequency. The $n(\ell)$ distribution is taken from Bull et al. (2015). The bandwidth B will be determined by the width of our chosen redshift bins, which is going to be $\Delta z = 0.1$ for all cases. Finally, we will consider a survey strategy with $t_{\text{tot}} = 4000 \text{ h}$ and $f_{\text{sky}} = 0.5$. An important point is the difference in the range of angular scales that the dish and interferometer mode can observe. We will analyse this in detail when we present our forecasts in Section 4.

3.2 CMB and 21-cm lensing

Following Pullen et al. (2015a) we will study the case where CMB is the background source plane for the lensing convergence measurements. We will calculate the lensing reconstruction noise using the formalism by Hu & Okamoto (2002). The instrumental noise for a CMB survey is given by

$$C_{\ell}^{\text{N}} = \Delta_{\text{T}}^2 \exp[\ell(\ell+1)\sigma^2/8\ln 2]. \quad (18)$$

We will consider a future CoRE-like satellite with full width at half-maximum $\sigma = 3.0 \text{ arcmin}$ and temperature noise $\Delta_{\text{T}} = 1 \mu\text{K}'$. We will also consider the full *Planck* lensing map (i.e. including temperature and polarization) using the sensitivities given by the *Planck* Collaboration (Tauber et al. 2006).

For the 21-cm lensing case we will use two different estimators depending on the chosen source redshift. The first one assumes the temperature distribution is Gaussian, which is a reasonable approximation at the Epoch of Reionization (EoR), at least while the ionized regions are small. This Fourier space estimator was developed in Zahn & Zaldarriaga (2006) and it is a 3D extension of the CMB lensing estimator by Hu & Okamoto (2002). More specifically, the 21-cm brightness temperature fluctuations are divided into wave vectors perpendicular to the line of sight $\mathbf{k}_{\perp} = l/\mathcal{D}$, with \mathcal{D} the angular diameter distance to the source redshift, and a discretized version of the parallel wave vector $k_{\parallel} = \frac{2\pi}{\mathcal{L}} j$ where \mathcal{L} is the depth of the observed volume. Considering modes with different j as independent, an optimal estimator can be found by combining the individual estimators for different j modes without mixing them. The three-dimensional lensing reconstruction noise is then found to be (Zahn & Zaldarriaga 2006)

$$N_{\ell}^{\kappa\kappa} = (\ell^4/4) \times \left[\sum_{j=j_{\text{min}}}^{j_{\text{max}}} \int \frac{d^2 \ell'}{(2\pi)^2} \frac{[l' \cdot l C_{\ell',j} + l \cdot (l-l') C_{|l'-\ell|,j}]^2}{2C_{\ell',j}^{\text{tot}} C_{|l'-\ell|,j}^{\text{tot}}} \right]^{-1}, \quad (19)$$

where

$$C_{\ell,j} = [\bar{T}(z)]^2 \frac{P_{\delta\delta}(\sqrt{(\ell/\mathcal{D})^2 + (j2\pi/\mathcal{L})^2})}{\mathcal{D}^2 \mathcal{L}}$$

and

$$C_{\ell,j}^{\text{tot}} = C_{\ell,j} + C_{\ell}^{\text{N}}.$$

For the EoR case we will use an SKA_Low-like interferometer array with uniform antennas distribution for which the thermal noise

power spectrum is given by (Zaldarriaga, Furlanetto & Hernquist 2004)

$$C_\ell^N = \frac{(2\pi)^3 T_{\text{sys}}^2}{B t_{\text{tot}} f_{\text{cover}}^2 \ell_{\text{max}}(\nu)^2}, \quad (20)$$

where $\ell_{\text{max}}(\nu)$ is the highest multipole that can be measured by the array at frequency ν (wavelength λ) and is related to D_{tel} , the maximum baseline of the core array, by $\ell_{\text{max}}(\lambda) = 2\pi D_{\text{tel}}/\lambda$; f_{cover} is the total collecting area of the core array, A_{coll} divided by $\pi(D_{\text{tel}}/2)^2$.

Finally, for post-reionization redshifts we will use SKA_Mid in interferometer mode and the lensing estimator developed in Poursidou & Metcalf (2014, 2015) using the IM technique. This estimator takes into account the discreteness of galaxies, and models the H I distribution as a Poisson distribution drawn from a Gaussian distribution that represents the clustering of galaxies. Further details and results for the lensing reconstruction using this technique and the updated SKA_Mid instrument have been analysed in detail in recent work (Poursidou et al. 2016) so we will not repeat them here. We just note that $N_\ell^{\kappa\kappa}$ for this case involves the underlying dark matter power spectrum $P_{\delta\delta}$, the H I density $\Omega_{\text{H I}}(z)$ as well as the H I mass (or luminosity) moments up to fourth order and, of course, the thermal noise of the instrument.

4 FORECASTS

In this section we will present our forecasts for the constraining power of combinations of IM, optical galaxy, and CMB surveys using the E_G statistic. Our goal is to show which combinations can measure E_G to very high statistical accuracy (<1 per cent) and provide exquisite tests of gravity at cosmological scales. For our fiducial cosmology we set the *Planck* Λ CDM cosmological parameters (Ade et al. 2014) and assume GR when calculating uncertainties.

We are going to consider two MG models. The first one is Chameleon gravity type models (Brax et al. 2004; Khoury & Weltman 2004), in which the scalar field which drives the cosmic acceleration couples to matter and acquires an environmentally dependent mass allowing consistence with local tests of gravity. In these theories the μ and γ coefficients in equation (4) can be parametrized using a three-parameter set (B_0, s, β_1) (Bertschinger & Zúkin 2008). We calculate the theoretical value of E_G for chameleon gravity using MGCAMB (Zhao et al. 2009; Lewis, Challinor & Lasenby 2000; Hojjati, Pogosian & Zhao 2011). We are also going to consider the popular growth index parametrization, where deviations from Λ CDM are expressed via the γ_L parameter and the linear growth factor can be written as $f = \Omega_m(z)^{\gamma_L}$ (Linder & Cahn 2007), with $\gamma_L = 6/11$ the GR value. Our chosen parameter set for the chameleon model is $(B_0, s, \beta_1) = (0.4, 4, 1.2)$. For the modified growth parametrization we take $\gamma_L = 0.65$ – note that we use γ_L as a trigger parameter, i.e. as a phenomenological parameter that is designed to indicate departures from GR, without the need to specify a particular theory.

As in Pullen et al. (2015a) we quantify our results by calculating the signal-to-noise ratio (SNR) defined as

$$\text{SNR}^2(E_G) = \sum_{\ell, z_i} \frac{[E_G^{\text{GR}}(z_i)]^2}{\sigma^2[E_G(\ell, z_i)]} \quad (21)$$

where z_i are the foreground redshift bins. The χ^2 values to discriminate GR and MG models are written as

$$\chi^2(E_G) = \sum_{\ell, z_i} \frac{[E_G^{\text{MG}}(\ell, z_i) - E_G^{\text{GR}}(z_i)]^2}{\sigma^2[E_G(\ell, z_i)]}, \quad (22)$$

with $E_G^{\text{MG}}(\ell, z_i)$ the E_G prediction for the specific MG models under consideration.

Below we present our results for the various combinations of surveys we have considered; the results are also summarized in Table 1.

4.1 Galaxy clustering × CMB lensing

We will start by considering photometric galaxy clustering surveys (DES and LSST) combined with CMB lensing with the *Planck* and CoRE-like satellites. CMB lensing is ‘cleaner’ than galaxy–galaxy lensing, avoiding issues like intrinsic alignments and source redshift uncertainties. Photometric galaxy clustering surveys have been found to be able to discriminate between GR and MG models more effectively than spectroscopic surveys (Pullen et al. 2015a). That is because the RSD error, which is better measured by spectroscopic surveys, is not the dominant source of error in equation (10). Hence, reducing the shot noise error by having higher number densities with a photometric survey is more important than precise RSD measurements. In our forecasts, we will assume a 17 per cent RSD error for DES (Ross et al. 2011), i.e. $\sigma(\beta)/\beta = 0.17$ in equation (10), and 10 per cent for LSST, and consider the wavenumber range $100 \leq \ell \leq 500$, since most of the signal for E_G comes from linear to quasi-linear scales for this case (Pullen et al. 2015a).

We combine the DES and LSST galaxy clustering photometric surveys with CMB lensing measurements using the *Planck* and CoRE-like satellites. Our forecasts for the measurement errors are shown in Fig. 1. The DES × *Planck* cross-correlation gives $\text{SNR} = 41$, while DES × CoRE gives $\text{SNR} = 85$. That is because the CoRE-like satellite has a noise level more than an order of magnitude lower than *Planck*. Taking LSST instead of DES we expect the results to improve significantly, as LSST covers a much bigger sky area and has increased number density of galaxies (the RSD measurement is also better). Indeed, we get $\text{SNR} = 95$ using *Planck* and $\text{SNR} = 198$ using the CoRE-like satellite. In the LSST × CoRE case we reach fractional errors smaller than 1 per cent. We also note that the SNR results using DES/LSST and the full *Planck* lensing map provide a consistency check of our calculations as they are in agreement with the ones found in Pullen et al. (2015a) using the same surveys.

4.2 Galaxy clustering × 21-cm lensing

In this section we will first consider the combination of galaxy clustering surveys with the weak gravitational lensing of the 21-cm emission from the EoR. This has similar advantages to using the CMB, for example precise source redshifts and the fact that we can use the full z -range of galaxy tracers. An additional advantage of 21-cm lensing is that one is able to combine information from multiple redshift slices, which makes the lensing reconstruction noise calculated using an SKA_Low-like instrument significantly lower than the one using the 2D CMB lensing estimator and a *Planck*-like satellite (Zahn & Zaldarriaga 2006). The possibility of measuring the lensing signal from the EoR has been studied in the past (Zahn & Zaldarriaga 2006; Metcalf & White 2009; Poursidou & Metcalf 2015). Here, we repeat these calculations assuming that the brightness temperature follows a Gaussian distribution, which is a reasonable approximation at the EoR, at least while the ionized regions are small.

We consider an SKA_Low-like instrument with collecting area $A_{\text{coll}} = 0.5 \text{ km}^2$ and maximum baseline $D_{\text{tel}} = 4 \text{ km}$ which scans

Table 1. Forecasts of the SNR and $\chi_{\text{rms}} = \sqrt{\chi^2}$ between GR and the MG models under consideration for the various survey combinations we consider. For the chameleon gravity model we set $(B_0, s, \beta_1) = (0.4, 4, 1.2)$, while for the modified growth model we use $\gamma_L = 0.65$ (see text for further details).

Survey	z_c	z_s	SNR	$\chi_{\text{rms}}[\text{Cham}]$	$\chi_{\text{rms}}[\gamma_L]$
DES \times <i>Planck</i> (full)	0.0–2.0	z_{cmb}	41	4.3	1.5
DES \times CoRE-like	0.0–2.0	z_{cmb}	85	8.9	3.0
LSST \times <i>Planck</i> (full)	0.0–2.5	z_{cmb}	95	10.1	3.1
LSST \times CoRE-like	0.0–2.5	z_{cmb}	198	21.1	6.4
LSST \times SKA_Low-like	0.0–2.5	$z_{\text{EoR}} = 7$	238	25.0	8.9
LSST \times SKA1_Mid	0.0–2.5	3	47	4.8	2.1
LSST \times SKA2_Mid	0.0–2.5	3	127	12.9	5.8
SKA1_Mid ^(sd) \times <i>Planck</i> (full)	0.35–3.0	z_{cmb}	34	3.5	1.3
SKA1_Mid \times <i>Planck</i> (full)	0.35–3.0	z_{cmb}	92	10.6	2.0
SKA1_Mid \times CoRE-like	0.35–3.0	z_{cmb}	200	23.1	4.6
SKA1_Mid \times SKA_Low-like	0.35–3.0	$z_{\text{EoR}} = 7$	227	25.3	6.0




Figure 1. E_G forecasts for the DES (top) and LSST (bottom) photometric optical galaxy surveys cross-correlated with the final *Planck* lensing map and with the CoRE-like lensing map. The Chameleon and modified growth predictions are also shown.

half of the sky in $t_{\text{tot}} = 10\,000$ hrs. Current SKA_Low plans include scanning a very small sky area so our chosen survey strategy is very optimistic, but we feel it is worth demonstrating the science potential of such an instrument. Another very interesting possibility if the proposed survey strategy is followed is constraining the standard cosmological parameters using 21-cm lensing measurements (Metcalf & White 2009).

In our forecasts the reionization redshift is assumed to be $z_{\text{EoR}} = 7$ and the observation bandwidth is $B = 8$ MHz. An important issue




Figure 2. E_G forecasts for the LSST photometric optical galaxy survey cross-correlated with 21-cm lensing measurements from the EoR at redshift $z_{\text{EoR}} = 7$ with an SKA_Low-like instrument, and with SKA2_Mid 21-cm lensing measurements at source redshift $z_s = 3$ using the IM method. The Chameleon and modified growth predictions are also shown.

with high-redshift (low-frequency) observations is the large foreground contamination. It has been shown that foreground subtraction techniques will remove the first parallel k modes (McQuinn et al. 2006). We model this effect by using $j_{\text{min}} = 4$ instead of $j_{\text{min}} = 1$ in equation (19). Combining the above with galaxy clustering and β measurements with LSST we find $\text{SNR} = 238$ and reach fractional errors smaller than 1 per cent. That is indeed much better than *Planck* and even exceeds the performance of a CoRE-like satellite. Considering a pessimistic case for the foreground contamination with $j_{\text{min}} = 10$, the lensing reconstruction noise doubles and we find $\text{SNR} = 208$, which still exceeds the performance of a CoRE-like satellite.

We are also going to consider the lensing of 21-cm emission from post-reionization redshifts (in particular $z_s = 3$) probed by the SKA_Mid array. In order to get results competitive with CMB lensing, we need to consider Phase 2 of the array – we model the thermal noise for this case like the one of Phase 1 but one order of magnitude smaller. We also take $f_{\text{sky}} = 0.5$, total observation time $t_{\text{tot}} = 4000$ h and $B = 20$ MHz – note that these numbers are realistic considering the current SKA_Mid plans and the possibility of commissioning it to perform an IM survey. Combining with LSST we find $\text{SNR} = 127$. Our forecasts for the measurement errors are shown in Fig. 2. Note that if we use SKA1_Mid (i.e. Phase 1 of the array) we find $\text{SNR} = 47$.

4.3 H I IM clustering × CMB/21-cm lensing

In this section we will investigate the combination of H I clustering surveys using the IM method with CMB and 21-cm lensing surveys. As shown in Pourtsidou et al. (2016), there are exciting prospects for performing clustering measurements using IM surveys with the SKA and its pathfinders. Note that we will assume a 10 per cent error in the RSD parameter β measurement, which is a very conservative estimate of the level of precision that should be achievable with an IM survey using SKA1_Mid (Bull 2016; Raccanelli et al. 2015).

We will start by considering the SKA1_Mid instrument in single-dish (sd) mode. An important point we need to stress is that the range of scales probed depending on the mode the instrument operates in (single dish or interferometer) is different. Using the single-dish mode we can probe very large scales, hence we are going to use $\ell_{\min} = 10$ and $\ell_{\max} = 2\pi D_{\text{dish}}/\lambda$ for our forecasts (Bull et al. 2015). For example, $\ell_{\max} \sim 220$ at $z \sim 1$. Using the full *Planck* map we find $\text{SNR} = 34$. Since the results are not competitive with our previous forecasts, and because of the issue of possibly severe systematic uncertainties when probing ultra-large angular scales (like in the single-dish mode), we are going to move on to consider the interferometer mode.

Using SKA1_Mid in interferometer mode we get much better results. In this case we can let $\ell_{\max} = 500$ like in the case of the optical galaxy surveys we analysed above, while the minimum multipole at each redshift is $\ell_{\min} = 2\pi D_{\text{dish}}/\lambda$. This gives $\ell_{\min} \sim 330$ at $z \sim 0.3$ and $\ell_{\min} \sim 110$ at $z \sim 3$, while $\ell_{\min} = 100$ for the optical galaxy surveys. Using the full *Planck* map we find $\text{SNR} = 92$, while with CORe we reach $\text{SNR} = 200$, achieving fractional errors < 1 per cent in the E_G measurements. This result implies that (assuming the problem of foreground contamination is alleviated) Phase 1 of the SKA can perform an IM survey with H I clustering measurements that are directly competitive with the galaxy clustering precision measurements by LSST. Note that considering Phase 2 of the SKA (which we model like SKA1 but with the noise level decreased by an order of magnitude) does not considerably improve the results, as the CMB lensing and β errors dominate. Our forecasts for the measurement errors using the interferometer mode are shown in the top panel of Fig. 3.

Finally, we consider the case where H I clustering measurements performed with SKA1_Mid are combined with the 21-cm EoR lensing case we studied previously using an SKA_Low-like instrument. We find $\text{SNR} = 227$, with fractional errors in the E_G measurements below 1 per cent. Our forecasts for the measurement errors for this case are shown in the bottom panel of Fig. 3.

As we have already mentioned, our SNR and χ_{rms} results are summarized in Table 1. We can see that we are able to differentiate between GR and MG at the level of several σ in a number of cases. The discriminating power of the measurements we have considered is larger for the Chameleon model, as it does not converge to the GR value at high redshifts ($z > 1$), while the modified growth model does. We will further comment on our results in Section 5. Before we conclude, we will show the clustering and noise terms for the various surveys we have considered.

4.4 Noise terms comparison

To consolidate our results, we compare the noise terms used for the various survey combinations studied in this work.

The top panel of Fig. 4 compares the tracer density power spectra and noise terms for DES (dot-dashed magenta line), LSST (dashed red line), SKA1_Mid single-dish mode (dotted green line)




Figure 3. E_G forecasts using the SKA1_Mid instrument cross-correlated with the final *Planck* lensing map and with the CORe-like lensing map (top) and the SKA_Low-like 21-cm lensing EoR map (bottom). The Chameleon and modified growth predictions are also shown.

and SKA1_Mid interferometer mode (solid black line) for the bin with central redshift $z_c = 1$. Here, we note that because of the non-uniform $n(\ell)$ antennas distribution the SKA1_Mid (int) noise curve is flat at large scales while at smaller scales (which we do not show here as they are not used) it increases as $\sim \ell^2$. Therefore, at the scales of interest for E_G the SKA1_Mid (int) instrument has its minimum thermal noise value; however, the minimum ℓ -scale it can probe is larger than the one of the optical galaxy surveys considered. The solid black curve is the angular power spectrum $C^{\delta\delta} = C^{\delta_{\text{H I}}, \delta_{\text{H I}}}/b_{\text{H I}}^2 = C^{\delta_{\text{g}}, \delta_{\text{g}}}/b_{\text{g}}^2$ at $z_c = 1$.

The bottom panel of Fig. 4 compares the lensing convergence power spectra and noise terms for the CORe-like satellite (blue dot-dashed line) and the SKA_Low-like interferometer (dashed black line). As we have already mentioned, the *Planck* noise is approximately one order of magnitude higher than CORe, while the SKA2_Mid noise level for H I sources at $z_s = 3$ is about three times higher than CORe (and, of course, the lensing convergence power spectrum is also lower at lower redshifts).

5 DISCUSSION AND CONCLUSIONS

In this work we considered H I IM clustering and lensing as probes of the clustering bias-free E_G statistic which can be used to test GR on cosmological scales. We forecasted the ability of various




Figure 4. Top: a comparison of the clustering noise terms for the various surveys considered for the bin with central redshift $z_c = 1$. Note that $C^{\delta\delta} = C^{\delta_{H1}, \delta_{H1}}/b_{H1}^2 = C^{\delta_g, \delta_g}/b_g^2$. Bottom: a comparison of the lensing noise terms for various surveys considered. We show the most competitive lensing measurements, i.e. using CMB lensing with a CORe-like satellite, and 21-cm lensing from the EoR with an SKA_Low-like instrument. See text for further details.

survey combinations – including IM, (photometric) optical galaxy and CMB lensing surveys – to test GR and constrain MG theories, in particular Chameleon gravity and modified growth index parametrization models.

Our results show that fractional errors < 1 per cent in the E_G measurement can be achieved in a number of cases, namely SKA1_Mid (LSST) H I IM (galaxy) clustering combined with CORe CMB lensing or, alternatively, with a SKA_Low-like EoR 21-cm lensing survey. Looking at the corresponding χ_{rms} values in Table 1, which are well above unity, we see that these surveys can provide strong constraints on the MG models parameters. The modified growth index model is more difficult to constrain as its expectation E_G value becomes indistinguishable from GR at $z > 1$ – that is why the $\chi_{rms}(\gamma_L)$ values are larger when using LSST (instead of SKA1_Mid), as LSST covers the $z < 0.35$ range where the differences between the growth parametrization model and GR are more pronounced (the optical galaxy surveys also cover a bigger ℓ -range than SKA1_Mid in interferometer mode.). It would therefore be very beneficial if the SKA1_Mid clustering and RSD measurements were extended to include the low-redshift regime, $0 < z < 0.35$ (Band 2).

The E_G statistic is clustering bias free, but its statistical error is affected by the bias and it is useful to comment on how sensitive it is to changes in the analytical formulae we have used. For example,

considering a factor of 2 smaller bias (i.e. $\sqrt{1+z}/2$ for the galaxy case and the analogous expression for the H I case), we still get less than a per cent fractional errors in our most competitive cases and the total SNR is reduced by about 3 per cent (on the other hand, a larger bias increases the SNR very slightly). That is because a smaller (larger) bias increases (decreases) the contribution of the shot/thermal noise terms.

In terms of the required precision in the RSD measurements (~ 10 per cent), the IM method appears to be quite advantageous. We already mentioned that in order to produce competitive E_G measurements using optical galaxy surveys one needs to use photometric instead of spectroscopic surveys, but measuring RSD this way is difficult (Pullen et al. 2015a). On the other hand, using IM and the SKA excellent redshift information is automatically provided and one can get results competitive with the ones by upcoming spectroscopic surveys like Euclid (Bull 2016; Raccanelli et al. 2015). Furthermore, a very recent paper on measuring E_G using number counts (Dizgah & Durrer 2016) showed that the usually neglected lensing contribution to galaxy number counts (which affects the g and $g\kappa$ spectra) is important – especially at high redshifts – and renders E_G bias dependent. IM does not suffer from this problem, because there is no magnification term at linear order as surface brightness is conserved.

Another feature of MG (and exotic dark energy) theories can be scale dependence. For example, E_G is found to be strongly scale dependent in the case of $f(R)$ gravity theories (Pullen et al. 2015a), so one could also use $E_G(k)$ measurements to constrain scale-dependent gravity. In this work we have mild scale dependence only in the case of Chameleon gravity, so we just averaged over the wavenumber range at each redshift bin for our predictions. However, this is a potentially very interesting subject and we plan to investigate it in future work.

An important point we need to stress is the need to control systematics. Future measurements will reach an unprecedented level of statistical precision (< 1 per cent) and if systematic effects are not correctly identified and removed the total error will be much larger. Details about systematics when combining galaxy surveys with CMB lensing can be found in Pullen et al. (2015b). In H I IM clustering and lensing surveys the biggest problem is the presence of galactic and extragalactic foregrounds. These can be orders of magnitude brighter than the H I signal but they have a smooth, power-law frequency dependence, in contrast to the fluctuating signal, so they can be removed (Morales, Bowman & Hewitt 2006; Liu & Tegmark 2011; Alonso et al. 2015). In order to identify systematics and test the various foreground removal techniques it is essential to perform auto- and cross-correlation clustering and lensing studies using IM and optical galaxy surveys. For this purpose we can exploit SKA pathfinders like MeerKAT (Poursidou et al. 2016). These studies will also give us precise measurements of the mean H I brightness temperature $\bar{T}(z)$, which is assumed to be known in our forecasts.

To conclude, the IM technique, although still in its infancy, is in principle ideal for testing GR and the standard cosmological model on large scales. IM surveys performed with the SKA have the advantage of excellent redshift resolution and they can map a large fraction of the sky across a wide range of scales and redshift, achieving very high SNR measurements. At large scales in particular, an IM survey with SKA1 can be directly competitive with state-of-the-art photometric optical galaxy surveys like LSST. Combining this with CMB lensing using CORe-like satellites or 21-cm lensing from the EoR with an SKA_Low-like array, we can perform exquisite tests of gravity and, consequently, help unravel the secrets of the dark sector of the Universe.

ACKNOWLEDGEMENTS

This work was supported by STFC grant ST/H002774/1. I would like to thank David Bacon, Davide Bianchi, Robert Crittenden, Dida Markovič and Matthew Withers for useful discussions.

REFERENCES

- Abell P. A. et al., 2009, preprint ([arXiv:0912.0201](https://arxiv.org/abs/0912.0201))
- Ade P. A. R. et al., 2014, *A&A*, 571, A16
- Alonso D., Bull P., Ferreira P. G., Santos M. G., 2015, *MNRAS*, 447, 400
- Ansari R. et al., 2012, *A&A*, 540, A129
- Battye R. A., Davies R. D., Weller J., 2004, *MNRAS*, 355, 1339
- Battye R., Browne I., Dickinson C., Heron G., Maffei B., Pourtsidou A., 2013, *MNRAS*, 434, 1239
- Becker M. R. et al., 2015, preprint ([arXiv:1507.05598](https://arxiv.org/abs/1507.05598))
- Bertschinger E., Zukin P., 2008, *Phys. Rev. D*, 78, 024015
- Blake C. et al., 2015, *MNRAS*, 456, 2806
- Brax P., van de Bruck C., Davis A.-C., Khoury J., Weltman A., 2004, *Phys. Rev. D*, 70, 123518
- Bull P., 2016, *ApJ*, 817, 26
- Bull P., Ferreira P. G., Patel P., Santos M. G., 2015, *ApJ*, 803, 21
- Camera S., Santos M. G., Ferreira P. G., Ferramacho L., 2013, *Phys. Rev. Lett.*, 111, 171302
- Chang T.-C., Pen U.-L., Peterson J. B., McDonald P., 2008, *Phys. Rev. Lett.*, 100, 091303
- Clifton T., Ferreira P. G., Padilla A., Skordis C., 2012, *Phys. Rep.*, 513, 1
- Copeland E. J., Sami M., Tsujikawa S., 2006, *Int. J. Mod. Phys. D*, 15, 1753
- Crighton N. H. M. et al., 2015, *MNRAS*, 452, 217
- Dewdney P., 2013, SKA Project Documents, p. 1, available at: https://www.skatelescope.org/wp-content/uploads/2012/07/SKA-TEL-SKO-DD-001-1_BaselineDesign1.pdf
- Dizgah A. M., Durrer R., 2016, preprint ([arXiv:1604.08914](https://arxiv.org/abs/1604.08914))
- Hojjati A., Pogosian L., Zhao G.-B., 2011, *J. Cosmol. Astropart. Phys.*, 1108, 005
- Hu W., Okamoto T., 2002, *ApJ*, 574, 566
- Khoury J., Weltman A., 2004, *Phys. Rev. Lett.*, 93, 171104
- Leonard C. D., Ferreira P. G., Heymans C., 2015, *JCAP*, 1512, 051
- Lewis A., Challinor A., Lasenby A., 2000, *ApJ*, 538, 473
- Limber D. N., 1954, *ApJ*, 119, 655
- Linder E. V., Cahn R. N., 2007, *Astropart. Phys.*, 28, 481
- Liu A., Tegmark M., 2011, *Phys. Rev. D*, 83, 103006
- Loeb A., Wyithe S., 2008, *Phys. Rev. Lett.*, 100, 161301
- Mao Y., Tegmark M., McQuinn M., Zaldarriaga M., Zahn O., 2008, *Phys. Rev. D*, 78, 023529
- McPherson A., 2015, Report and Options for Re-baselining of SKA-1, available at: https://www.skatelescope.org/wp-content/uploads/2014/03/SKA-TEL-SKO-0000229_Report-and-Options-for-Re-Baselining-of-SKA-1-Word_Amm-signed.pdf
- McQuinn M., Zahn O., Zaldarriaga M., Hernquist L., Furlanetto S. R., 2006, *ApJ*, 653, 815
- Metcalf R. B., White S. D. M., 2009, *MNRAS*, 394, 704
- Morales M. F., Bowman J. D., Hewitt J. N., 2006, *ApJ*, 648, 767
- Perlmutter S. et al., 1999, *ApJ*, 517, 565
- Peterson J. B. et al., 2009, preprint ([arXiv:0902.3091](https://arxiv.org/abs/0902.3091))
- Pourtsidou A., Metcalf R. B., 2014, *MNRAS*, 439, L36
- Pourtsidou A., Metcalf R. B., 2015, *MNRAS*, 448, 2
- Pourtsidou A., Bacon D., Crittenden R., Metcalf R. B., 2016, *MNRAS*, 459, 863
- Pullen A. R., Alam S., Ho S., 2015a, *MNRAS*, 449, 4326
- Pullen A. R., Alam S., He S., Ho S., 2015b, preprint ([arXiv:1511.04457](https://arxiv.org/abs/1511.04457))
- Raccanelli A. et al., 2015, PoS, Cosmology Chapter, Advancing Astrophysics with the SKA (AASKA14), 14, 031
- Reyes R., Mandelbaum R., Seljak U., Baldauf T., Gunn J. E., Lombriser L., Smith R. E., 2010, *Nature*, 464, 256
- Riess A. G. et al., 1998, *AJ*, 116, 1009
- Ross A. J., Percival W. J., Crocce M., Cabre A., Gaztanaga E., 2011, *MNRAS*, 415, 2193
- Seo H.-J., Dodelson S., Marriner J., McGinnis D., Stebbins A., Stoughton C., Vallinotto A., 2010, *ApJ*, 721, 164
- Switzer E. et al., 2013, *MNRAS*, 434, L46
- Tauber J. et al., 2006, preprint ([astro-ph/0604069](https://arxiv.org/abs/astro-ph/0604069))
- Zahn O., Zaldarriaga M., 2006, *ApJ*, 653, 922
- Zaldarriaga M., Furlanetto S. R., Hernquist L., 2004, *ApJ*, 608, 622
- Zhang P., Liguori M., Bean R., Dodelson S., 2007, *Phys. Rev. Lett.*, 99, 141302
- Zhao G.-B., Pogosian L., Silvestri A., Zylberberg J., 2009, *Phys. Rev. D*, 79, 083513

This paper has been typeset from a $\text{\TeX}/\text{\LaTeX}$ file prepared by the author.