



Antifouling bastadin congeners target blue mussel phenoloxidase and complex copper(II) ions

Journal:	<i>Marine Biotechnology</i>
Manuscript ID:	Draft
Manuscript Type:	Original Article
Date Submitted by the Author:	n/a
Complete List of Authors:	Bayer, Mirko; Heinrich-Heine-Universität Düsseldorf, Institut für Pharmazeutische Biologie Hellio, Claire; Portsmouth University, School of Biological Sciences Marechal, Jean-Philippe; Portsmouth University, School of Biological Sciences Lin, Wenhan; Peking University, State Key Laboratory of Natural and Biomimetic Drugs Frank, Walter; Heinrich-Heine-Universität Düsseldorf, Institut für Anorganische Chemie und Strukturchemie Weber, Horst; Heinrich-Heine-Universität Düsseldorf, Institut für Pharmazeutische und Medizinische Chemie Proksch, Peter; Heinrich-Heine-Universität Düsseldorf, Institut für Pharmazeutische Biologie
Keywords:	<i>Mytilus edulis</i> , mussel phenoloxidase, antifouling, bastadins, hemibastadin, dibromohemibastadin

1
2
3 **Antifouling bastadin congeners target blue mussel phenoloxidase**
4
5
6 **and complex copper(II) ions**
7
8
9

10
11
12
13
14
15
16 Mirko Bayer^{1,6}, Claire Hellio³, Jean-Philippe Marechal³, Wenhan Lin⁵, Walter Frank⁴,
17
18 Horst Weber², Peter Proksch^{1a}
19

20
21
22
23
24
25
26
27
28 ¹Institut für Pharmazeutische Biologie und Biotechnologie, Heinrich-Heine-Universität
29
30 Düsseldorf, Universitätsstr. 1, 40225 Düsseldorf, Germany
31

32
33 ²Institut für Pharmazeutische und Medizinische Chemie, Heinrich-Heine-Universität
34
35 Düsseldorf, Universitätsstr. 1, 40225 Düsseldorf, Germany
36

37
38 ³School of Biological Sciences, King Henry Building, Portsmouth University,
39
40 Portsmouth PO1 2DY, United Kingdom
41

42
43 ⁴Institut für Anorganische Chemie und Strukturchemie, Heinrich-Heine-Universität
44
45 Düsseldorf, Universitätsstr. 1, 40225 Düsseldorf, Germany
46

47
48 ⁵State Key Laboratory of Natural and Biomimetic Drugs, Peking University, PR China

49
50 ⁶Caesar & Loretz GmbH, Herderstr. 31, 40721 Hilden, Germany
51

52
53
54
55
56 ^acorresponding author: Prof. Dr. Peter Proksch, proksch@uni-duesseldorf.de, 0049-
57
58 211-81-14170
59
60

Abstract

Synthetically prepared congeners of sponge-derived bastadin derivatives such as DBHB that suppress the settling of barnacle larvae were identified in this study as strong inhibitors of blue mussel phenoloxidase that is involved in firm attachment of mussels and of other marine fouling organisms to a given substrate. The IC_{50} value of DBHB as the most active enzyme inhibitor encountered in this study amounts to 0.84 μ M. Inhibition of phenoloxidase by DBHB is likely due to complexation of copper(II) ions from the catalytic centre of the enzyme by the α -oxo-oxime moiety of the compound as shown here for the first time by structure activity studies and by X-ray structure determination of a copper(II) complex of DBHB.

Abbreviations: 2,3-BMO (2,3-butanedione monoxime), DBHB (6,6'-dibromohemibastadin-1), DMF (dimethylformamide), NBHB (norbromohemibastadin-1), TBT (tri-n-butyl tin oxide), TT (tyrosinyltyramine)

Keywords: *Mytilus edulis*, mussel phenoloxidase, antifouling, bastadins, hemibastadin, dibromohemibastadin, norbromohemibastadin, tyrosinyltyramine, 2,3-butanedione monoxim, x-ray, copper(II), copper complex, natural products

Introduction

Man-made as well as natural surfaces in the marine ecosystem are rapidly colonized by fouling organisms that include microorganisms, algae, barnacles, mussels and others. Biofouling causes significant economical wastage worldwide e.g. by reducing boat speeds and increasing fuel consumption (Wahl 1989, Yebra et al. 2004). Considering that up to 90% of global trade rely on ships serious problems due to biofouling arise for the maritime industry (Liu et al. 1997, Voulvoulis et al. 1999, Marechal and Hellio 2009). Until recently, coatings containing tri-*n*-butyl tin oxide (TBT) were used for preventing biofouling. Since 2008 usage of TBT is banned by the International Maritime Organisation (IMO 2001, Voulvoulis et al. 1999, Lau 1991, Pereira and Ankjaergaard 2009) and new antifouling strategies, involving chemical, physical, and mechanical mechanisms, are urgently needed to replace TBT and other potentially toxic or harmful ingredients (Hellio and Yebra 2009).

The frequently made observation that many sponges are conspicuously free of overgrowth caused by fouling organisms has stimulated the search for naturally occurring antifouling leads from marine sponges (e.g. Tsoukatou et al 2002, Fusetani 2004, Tsoukatou et al 2007, Hellio et al 2009). In this context we demonstrated recently significant antifouling activity for sponge-derived bastadin and hemibastadin derivatives and for several synthetically derived analogues such as DBHB using larvae of the barnacle *Amphibalanus improvisus* (Darwin, 1854) as a model organism (Ortlepp et al., 2007). Bastadins are tyrosine-derived, brominated, oxime bearing peptides that can be cyclic or linear and are typical constituents of marine sponges from the genus *lanthella* such as *I. basta* (Ortlepp et al. 2007). Hemibastadin-1 is the simplest bastadin derivative known and can be considered as a biogenetic precursor of the larger bastadins including two or more hemibastadin moieties that are usually

1
2
3 linked through ether functions. Hemibastadin-1 as well as its closely related
4
5 synthetically derived analogue DBHB suppresses settling of barnacle larvae at low
6
7 micromolar concentrations making these compounds interesting candidates in the
8
9 search for new antifouling constituents from nature especially as they are easily
10
11 accessible by organic synthesis (Ortlepp et al. 2007).
12
13
14
15
16

17 It is well known that phenoloxidases of mussels and of other marine invertebrates
18
19 such as barnacle larvae are highly relevant for settling of these fouling organisms
20
21 since they are involved in polymerization of secreted proteins called foot proteins
22
23 (Aladaileh et al. 2007, Hellio et al. 2000, Zentz et al. 2001). The latter contain
24
25 numerous residues of the aromatic amino acids phenylalanine and tyrosine.
26
27 Phenoloxidases oxidize these aromatic amino acids and form highly reactive ortho-di-
28
29 quinone bearing systems that cross-react with each other as well as with other
30
31 proteins and generate a biopolymer, which firmly attaches the fouling organism to a
32
33 suitable surface (Suci and Geesey 2000, Aladaileh et al. 2007, Kamino 2001,
34
35 Cheung et al. 1977). In order to prevent a spontaneous polymerization of aromatic
36
37 amino acids phenoloxidase is stored intracellularly as enzymatically inactive pro-
38
39 phenoloxidase. Chemical transformation and activation of pro-phenoloxidase to
40
41 phenoloxidase is proteolytically catalyzed during secretion by different proteases that
42
43 are controlled by Ca^{2+} dependant signalling cascades (Asokan et al. 1997, Hellio et al.
44
45 2007). It is known that intracellular Ca^{2+} concentration is important for settling
46
47 (Rittschof et al. 1986, Clare 1996, Yamamoto et al. 1999). Thus, by reduction of Ca^{2+}
48
49 influx the signalling cascades that are involved in proteolytic conversion of pro-
50
51 phenoloxidase and in secretion of the active phenoloxidase can be inhibited. TBT
52
53 interferes with the cellular calcium regulator calmodulin and thus reduces mussel
54
55
56
57
58
59
60

1
2
3 phenoloxidase activity (Cima and Ballarin 2000, Cima et al. 1998a,b, Tujula et al.
4
5 2001).

6
7
8
9
10 A further strategy for inhibiting the attachment of fouling organisms is the search for
11
12 direct enzyme inhibitors of phenoloxidase. This family of oxygen activating enzymes
13
14 is distinguished by containing copper(II) ions that are bound to histidin moieties in the
15
16 catalytic centre playing an essential role for the activity of these enzymes (Kim and
17
18 Uyama 2005; Olivares et al. 2002). If a suitable phenolic substrate draws close to the
19
20 active site of the enzyme, hydroxylation and further oxidation is performed by
21
22 electron transfer from copper via oxygen to the substrate (Decker et al. 2000).
23
24
25
26
27 Complexation of copper ions would thus be a powerful way of inhibiting the
28
29 enzymatic activity of phenoloxidase and as a result attachment of fouling organisms.
30
31
32
33

34 Several inhibitors of phenoloxidases are already known from natural sources, like
35
36 quercetin and other flavonoids, mimosine or kojic acid (Kubo et al. 2000, Chen et al.
37
38 1991, Kim and Uyama 2005, Kim et al. 2006). Some synthetically derived
39
40 halogenated phenols have also been reported to inhibit phenoloxidases (Nagasawa
41
42 et al. 1981). Hemibastadin, however, has so far not been investigated as an inhibitor
43
44 of marine-derived phenoloxidases. In this study we explore the mode of action of
45
46 hemibastadins and of related compounds with regard to their antifouling activity and
47
48
49
50 report on:

- 51
52
- 53 • Inhibition of purified blue mussel phenoloxidase by DBHB and by some of its
54 derivatives *in vitro*
55
 - 56 • Ability of these compounds to complex copper(II) ions
57
 - 58 • X-ray analysis of a distinct complex of DBHB with copper(II) chloride,
59
60 crystallized from DMF and dichloromethane

Results

Several synthetically prepared hemibastadin-1 congeners that included DBHB, NBHB, and TT that differs from the former two compounds by absence of bromine atoms in the aromatic rings and by replacement of the oxime function of the hemibastadin-congeners by an amine substituent were selected for *in vitro* studies with phenoloxidase prepared from the blue mussel *Mytilus edulis*. 2,3-Butanedione monoxime (2,3-BMO) that is likewise of synthetic origin was also included in the experiments as it features an α -oxo-oxime moiety and thus resembles an important characteristic functionality of DBHB and NBHB but lacks the aromatic rings of the latter two compounds. Nearly all compounds analyzed proved to be inhibitors of mussel phenoloxidase albeit at very different concentrations. DBHB was by far the strongest inhibitor encountered with an IC_{50} value of 0.84 μ M. Next in activity was NBHB with an IC_{50} of 2.41 μ M followed by 2,3-BMO with an IC_{50} value of 8.70 μ M thereby indicating a clear enhancing effect of the phenolic rings and of the bromine substituents on the enzyme inhibitory activity of the compounds. TT, if at all, was by far the weakest inhibitor detected in this study, its IC_{50} value was $> 83 \mu$ M (**Fig. 1**). This latter finding underlines the striking importance of the α -oxo-oxime function for inhibition of mussel phenoloxidase especially when viewed in comparison with the data achieved for the structurally closely related NBHB.

Complexation of free copper(II) ions by DBHB was determined photometrically in methanolic solution. Addition of increasing amounts of copper(II) sulphate to an arbitrarily chosen concentration of DBHB resulted in the formation of a distinct bathochromic band (green colour) at 685 nm in the VIS absorption spectrum (**Fig. 2**). Absorption intensity was found to increase upon addition of further amounts copper(II) sulphate. Maximum absorption was reached after a five to tenfold molar excess of

1
2
3 the copper salt in relation to DBHB (**Fig. 3**). NBHB and 2,3-BMO were likewise able
4 to complex copper(II) ions as shown by the green coloration of a methanolic solution
5 of the compounds following addition of copper(II) sulphate (data not shown) whereas
6 no effect was observed for TT.
7
8
9
10
11

12
13
14
15 Subsequently we were able to prepare a neutral complex of DBHB with Cu(II) ions.
16 When a solution of DBHB in methanol is mixed with an aqueous copper(II) acetate
17 solution and the pH is adjusted to >9 by addition of sodium acetate, a brown
18 precipitate is formed that can be separated. After drying a cocoa like powder remains
19 that is insoluble in all common solvents but readily soluble in dilute HCl under
20 decolourization and precipitation of DBHB. Addition of dilute NaOH forms a clear
21 deep green coloured solution. Chemical composition of the brown powder was
22 determined by elementary analysis showing that one copper(II) ion is bound to two
23 molecules of DBHB (**Fig. 4**). Whereas this amorphous material was unsuitable for X-
24 ray analysis, we were successful in preparing a macrocrystalline complex of DBHB
25 with copper(II) chloride, which was crystallized from DMF and dichloromethane.
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42

43 In parallel the crystal structure of free DBHB was likewise analyzed by X-ray analysis.
44 Both crystal structures are reported here for the first time. Crystal data and details of
45 the structure determination processes are presented in the materials and methods
46 section. Representative geometrical data for both structures are summarized in **Tab.**
47 **1** and **Tab. 2**. These data in combination with **Fig. 5** allow for a detailed analysis of
48 copper coordination and of the ligand properties of DBHB. The metal centres of the
49 centrosymmetric dinuclear complex exhibit the typical 4+2-coordination mode of a
50 Jahn-Teller distorted octahedral complex with a terminal (Cl₂) and a bridging (Cl_{1a})
51 chloride ligand occupying the axial positions. As to be expected the corresponding
52
53
54
55
56
57
58
59
60

1
2
3 Cu-Cl bond lengths are much larger than the length of the equatorial Cu1-Cl1 bond.
4
5 According to the observed Cu-O and Cu-N distances strong bonding between the
6
7 copper centre and the chelating DBHB ligand as well as the DMF ligand has to be
8
9 assumed. Comparison of the relevant geometrical parameters of free and complexed
10
11 DBHB (right half of **Tab. 1**) gives evidence for the high complexation potential of
12
13 DBHB towards copper(II) ions. From the coordination chemistry point of view
14
15 dinuclear copper complexes with a $\text{Cu}_2(\mu\text{-Cl})_2$ core and a $\mu\text{-Cl,Cl}_2,\text{O}_2,\text{N}$ -coordination
16
17 motif are very rare species (Kapoor et al. 2002, Kapoor et al. 2004) and we here
18
19 present the first example of such a complex containing chlorido ligands in both the
20
21 axial positions of the distorted hexacoordinated copper centers. In principle the
22
23 closest structural relationship is given to di- μ -chloridobis((N,N''-2-pyridyl-2'-pyridinium
24
25 ketone oxime)aquachlorido-copper(II)) (Sommerer et al. 1995), a $\text{Cu}_2(\mu\text{-Cl})_2$ core-
26
27 complex with copper centers that are hexacoordinated in a $\mu\text{-Cl, Cl}_2,\text{N}_2,\text{O}$ -mode with
28
29 a $\mu\text{-Cl}$ and an aqua ligand in axial positions and the N,N'-chelating pyridyl ketone
30
31 oxime ligand as well as two Cl ligands in equatorial positions. Comparison of the
32
33 relevant bond lengths gives further evidence that bonding of DBHB to copper centers
34
35 is comparatively strong. As can be seen from **Fig. 5** and the hydrogen bond
36
37 parameters listed in **Tab. 2**, DBHB bonded to the copper centers is further involved in
38
39 supramolecular interactions. It has to be assumed that its H bridging donor and
40
41 acceptor properties also play an important role in the receptor binding process.
42
43
44
45
46
47
48
49
50

51
52
53 Complexation of copper(II) ions is not restricted to DBHB or other synthetic
54
55 congeners such as NBHB or 2,3-BMO but extends also to naturally occurring
56
57 bastadin derivatives that share the same important structural elements (α -oxo-oxime
58
59 group) as shown by treatment of a crude aqueous methanolic extract of the sponge *I.*
60
basta with an excess of copper(II) sulphate. A brown precipitate was formed after

1
2
3 addition of sodium carbonate solution which was separated and disintegrated with
4 dilute HCl. The acidic solution was extracted with ethyl acetate followed by HPLC and
5
6 LC-MS analysis. It revealed an enriched complex mixture of several known bastadin
7
8 derivatives (data not shown) that had been formerly described from the same sponge
9
10 (Ortlepp et al., 2007) thereby indicating that also naturally occurring bastadins have
11
12 the capacity to complex copper(II) ions.
13
14
15
16
17
18
19
20
21
22

23 **Materials and Methods**

24 1. Improved synthesis of NBHB and DBHB

25
26 1.1. NBHB: In contrast to the earlier reported low yielding melting procedure (Ortlepp
27
28 et al. 2007) we now describe an improved synthesis of NBHB. Methyl-[2-
29
30 hydroximino-3-(4-hydroxyphenyl)]-propionate (4.18 g, 0.02 mol) was triturated with
31
32 tyramine (6.03 g, 0.044 mol). This mixture was transferred into a small wide-mouth
33
34 flask and DMF (10 g) was added. The open flask was heated in an oil bath up to
35
36 100°C. The resulting amber syrup was magnetically stirred at 135 – 145 °C for 20
37
38 minutes and then cooled to room temperature. After addition of water (20 ml) and
39
40 ethyl acetate (50 ml) the mixture was stirred until nearly all has been dispensed and
41
42 only small amounts of insoluble polymeric material remained. The liquid was
43
44 transferred to a separation funnel using additional ethyl acetate (200 ml). The organic
45
46 layer was washed with water and this phase was extracted exhaustively with ethyl
47
48 acetate. The combined organic phase was washed successively with 1% HCl
49
50 (rejected) and 1% aqueous sodium carbonate solution (rejected). The finally
51
52 remaining ethyl acetate extract was worked up as usual. The after evaporation
53
54 remaining brown syrup (7 g) was diluted with ethyl acetate (5 ml) and
55
56 dichloromethane was added drop by drop until turbidness. This mixture was
57
58
59
60

1
2
3 fractionated on a silica gel column (40 x 5 cm, silica gel 0.064 – 0.2) with
4 dichloromethane/ethyl acetate (1400 + 600) to give after crystallization of the
5 evaporated residue 4.8 g (76.3%) of NBHB as colourless crystals, melting point 178 –
6
7
8
9
10
11 179 °C.

12
13
14
15 1.2. DBHB: NBHB (9.43 g, 0.03 mol) was dissolved in DMF (30 ml) and diluted with
16 dichloromethane (200 ml). After cooling on crashed ice a solution of bromine (150 ml
17 1M-Br₂ in dichloromethane) was added during 15 minutes under stirring an cooling
18 away from light. Complete bromination was controlled by thin layer chromatography.
19
20 The excess of bromine was reduced by adding aqueous 10% sodium hydrogen
21 sulphite solution until the brown colour was converted to a pale yellow. The water
22 phase was separated and extracted exhaustively with ethyl acetate. Both organic
23 phases were separately washed with water, then combined, dried over anhydrous
24 sodium sulphate, filtered and concentrated under vacuum. Addition of
25 dichloromethane to the residue resulted in crystallization of 8.5 g pure DBHB.
26
27 Column chromatography of the mother liquor on silica gel with dichloromethane/ethyl
28 acetate/methanol (1000+100+10) followed by crystallization gave additional 7.7 g of
29 pure DBHB (overall yield 85.7%), melting point 155 – 158 °C. C₁₇H₁₄Br₄N₂O₄ (629.92):
30
31 C calcd. 32.41, found 32.59, H calcd. 2.24, found 2.31, N calcd. 4.45, found 4.40.
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
This and the following elementary analyses were carried out using a Perkin-Elmer PE
2400 CHN Elemental Analyzer (Perkin-Elmer Instruments GmbH, Rodgau-
Jügesheim, Germany). The analytically pure compound was crystallized for x-ray
analysis from hot ethyl acetate by slowly cooling down to room temperature. Crystals
were separated, washed with cold ethyl acetate and dried in vacuum at room
temperature.

1
2
3 2. Inhibition of phenoloxidase from the blue mussel *Mytilus edulis*: Phenoloxidase
4 activity was measured spectrophotometrically as described by Hellio et al. (2000a):
5
6 the purified enzyme was incubated at 25 °C with 10 mM L-dopa or catechol in 50 mM
7
8 phosphate buffer of pH 6.8. Phenoloxidase activity was determined by monitoring the
9
10 increase of absorbance at a wavelength of 475 nm. One unit enzyme activity was
11
12 defined as the amount of enzyme that catalyzes the formation of 1 µmol dopachrome
13
14 per minute under the described experimental conditions. Enzyme inhibitors such as
15
16 DBHB were added to the assay at concentrations up to 50 µg/ml. In addition, the
17
18 biocide TBTO (10 µg/ml) was used as a positive standard (Hellio et al., 2000a).
19
20 Aliquots of pure enzyme were incubated for 2 hours with DBHB or other compounds
21
22 tested as inhibitors, then the enzyme activity was recorded with L-dopa or catechol
23
24 (10 mM) as substrates. All assays were run in triplicate.
25
26
27
28
29
30
31
32
33

34 3. Measurement of copper(II)-complex formation of DBHB in solution: DBHB and
35
36 CuSO₄ were dissolved in methanol (Merck p.a.) and added to a 1 cm-cuvette
37
38 resulting in final concentrations of each 333 µmol DBHB with increasing
39
40 concentrations of CuSO₄ (167, 333, 555, 1111, 1667, and 3333 µmol per ml. The
41
42 molar ratios between DBHB and CuSO₄ were thus 2:1, 1:1.7, 1:3.3, 1:5, and 1:10.
43
44 Assays were measured photometrically in a cuvette with a thickness of 1 cm in a
45
46 Perkin Elmer Lambda 25 UV/VIS spectrometer connected with the computer software
47
48 Perkin Elmer UV winlab 5.1.3.0626. Each sample was measured against methanol
49
50 containing the same molar concentration of CuSO₄ as present in the assay.
51
52
53
54
55
56

57 4. Neutral copper precipitate of DBHB: DBHB (630 mg, 1 mmol) was dissolved in
58
59 methanol (20 ml) and copper acetate (182 mg, 1 mmol) in water (20 ml) was added.
60
After addition of excess sodium acetate solution in water the pH was adjusted to 9

1
2
3 and the brown precipitate was removed by suction, washed with water and dried in
4
5 vacuum at 100°C. The resulting cocoa like powder could be identified as a neutral
6
7 copper complex where one Cu^{2+} is bound to two deprotonated DBHB molecules.
8
9 $\text{C}_{34}\text{H}_{26}\text{Br}_8\text{CuN}_4\text{O}_8$ (1321.37): C calcd. 30.90, found 30.72, H calcd. 1.98, found 2.02,
10
11 N calcd. 4.24, found 4.13.
12
13
14
15
16

17 5. Selective bastadin precipitation and recovery from a crude extract of the sponge

18 *lanthella basta*: To an aliquot of a crude methanolic extract of the sponge *l. basta* a
19
20 surplus of CuSO_4 solution in methanol was added. By addition of sodium carbonate
21
22 the pH of the solution was titrated to pH 11. The formed brown precipitate was
23
24 recovered by centrifuging (1,000 g) whereas the supernatant was rejected. The pellet
25
26 was washed twice with sodium carbonate solution and with water followed by
27
28 addition of 10 ml of diluted HCl. The acidic aqueous phase was partitioned three
29
30 times against 10 ml ethyl acetate followed by evaporation of the ethyl acetate soluble
31
32 part using a rotary evaporator. The residue was dissolved in 2 ml methanol, injected
33
34 into a HPLC machine and analyzed for bastadin congeners as described previously
35
36 (Ortlepp et al., 2007).
37
38
39
40
41
42
43
44
45

46 6. Crystallization of the DBHB complex with copper(II) chloride [Di- μ -chloro-bis(6,6'-

47 dibromohemibastadin-1)(dichloro)bis(dimethylformamido)dicopper(II)]: DBHB (630
48
49 mg, 1 mmol) was dissolved in 2-propanol (3 ml). The colourless solution was
50
51 combined with the light green solution of anhydrous CuCl_2 (134 mg, 1 mmol) in 2-
52
53 propanol (5 ml). The colour of the mixture turned to an intensive dark green.
54
55 Precipitation of an intensive green powder was completed by cooling over night in a
56
57 refrigerator. After separation with a suction filter the precipitate was washed with cold
58
59 2-propanol and dried in vacuum. Yield 750 mg (98%). This microcrystalline powder,
60

1
2
3 however, was not suitable for x-ray analysis. It was nearly insoluble in all common
4 solvents except DMF. Sufficient crystallisation was achieved by slowly dropping
5 dichloromethane into a DMF-solution of the precipitate at room temperature. The light
6 green crystals were separated, washed with dichloromethane and dried in vacuum at
7 100°C. They were insoluble in water or in dichloromethane but more or less soluble
8 in methanol, ethanol and 2-propanol. Elemental analysis showed DMF and
9 dichloromethane as constituents: [1 DBHB x 1 CuCl₂ x 1 DMF x 1 CH₂Cl₂],
10 C₂₁H₂₃Br₄Cl₄CuN₃O₅ (922.4): C calcd. 27.34, found 27.30; H calcd. 2.51, found 2.48;
11 N calcd. 4.56, found 4.47. If a solution of this compound in methanol is treated with
12 an excess of aqueous sodium carbonate solution a brown precipitate is formed which
13 was identical with the neutral copper precipitate of DBHB. Successful X-ray-analysis
14 of the complex of DBHB with CuCl₂/DMF/dichloromethane could only be achieved
15 with those crystals, which only were carefully dried at room temperature. This
16 material contains more than 1 equivalent of the latter solvent (see below: X-ray
17 crystallographic study). However, if the drying procedure is performed at higher
18 temperature, the required crystalline structure of the material is lost.

19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43 7. X-ray Crystallographic Study: Crystal Structure Determinations of compounds
44 C₁₇H₁₄Br₄N₂O₄ (DBHB) and (C₄₀H₄₀Br₈Cl₄Cu₂N₆O₁₀ · 4CH₂Cl₂ ((DBHB(DMF)CuCl₂)₂ ·
45 4CH₂Cl₂): Crystals suitable for X-ray study were selected by means of a polarisation
46 microscope and investigated on a STOE Imaging Plate Diffraction System using
47 graphite monochromatized MoK α radiation ($\lambda = 0.71073 \text{ \AA}$). To avoid loss of CH₂Cl₂
48 and deterioration, crystals of the copper compound had to be enclosed in thin walled
49 glass capillaries and the study had to be performed at 173 K. Unit cell parameters
50 were determined by least-squares refinements on the positions of 6074 and 8000
51 reflections in the range $1.95^\circ < \theta < 25.85^\circ$ and $6.0^\circ < \theta < 20.75^\circ$, respectively. Space
52
53
54
55
56
57
58
59
60

1
2
3 group type No. 14 was uniquely determined in the case of DBHB. For crystals of the
4
5 copper compound systematic extinctions were consistent with space group types *Cc*
6
7 and *C2/c*. In accordance with E-statistics significantly better results were observed
8
9 with a disordered structural model in the centrosymmetric type *C2/c*. Corrections for
10
11 Lorentz and polarization effects and multi-scan absorption corrections were applied
12
13 for both compounds. The structures were solved by direct methods (Sheldrick, 1990)
14
15 and subsequent ΔF -syntheses. Approximate positions of all the hydrogen atoms of
16
17 DBHB and of the dinuclear copper complex were found in different stages of
18
19 refinements by full-matrix least-squares calculations on F^2 (Sheldrick, 1997). The
20
21 hydrogen atom positions of the non-coordinated CH_2Cl_2 molecules in the copper
22
23 compound needed to be calculated. Refinement suffered from the high mobility of
24
25 these solvent molecules and appropriate distance and displacement parameter
26
27 restraints and constraints had to be applied to achieve convergence. Anisotropic
28
29 displacement parameters were refined for all non-hydrogen atoms. For the H atoms
30
31 at N2, O1 and O3 of $\text{C}_{17}\text{H}_{14}\text{Br}_4\text{N}_2\text{O}_4$ positional and isotropic displacement parameters
32
33 were refined. With idealized bonds lengths and angles assumed for all the other OH
34
35 and NH groups and for all the CH_3 , CH_2 , and CH groups of both compounds, the
36
37 riding model was applied for the corresponding H atoms. In addition, the H atoms of
38
39 the OH groups were given allowance for rotation around the neighbouring O-C axis,
40
41 and the H atoms of the CH_3 groups were allowed to move collectively around the
42
43 neighbouring C-C axis. An isotropic displacement parameter was refined for the H
44
45 atom at N2 of the copper complex. The isotropic displacement parameters of the H
46
47 atoms not explicitly mentioned here were constrained to 120% of the equivalent
48
49 isotropic displacement parameters of the parent O, N and C atoms for the NH, CH
50
51 and CH_2 groups and equal to 150% for the OH and the CH_3 groups. CCDC-782628
52
53 ($\text{C}_{17}\text{H}_{14}\text{Br}_4\text{N}_2\text{O}_4$) and CCDC-782629 ($\text{C}_{40}\text{H}_{40}\text{Br}_8\text{Cl}_4\text{Cu}_2\text{N}_6\text{O}_{10} \cdot 4\text{CH}_2\text{Cl}_2$) contain the
54
55
56
57
58
59
60

1
2
3 supplementary crystallographic data (excluding structure factors) for this paper.
4
5 These data can be obtained free of charge from The Cambridge Crystallographic
6
7 Data Centre *via* www.ccdc.cam.ac.uk/data_request/cif.
8
9

10 11 12 13 14 15 **Discussion**

16
17 DBHB, NBHB and 2,3-BMO proved to be strong inhibitors of blue mussel
18 phenoloxidase (**Fig. 1**). The IC₅₀ value of the most active compound DBHB (0.84 μM)
19
20 compares favourably with those of other reported marine-derived blue mussel
21
22 phenoloxidase inhibitors such as the meroditerpenoids (3a*S*,7a*R*)-1,2,3,3a,7,7a-
23
24 hexahydro-5-(2-hydroxy-2-methylpropyl)-6-((*R*)-6-methoxy-2,8-dimethyl-2H-chromen-
25
26 2-yl)-3a,7a-dimethylinden-4-one (IC₅₀: 2.3 μM) or (1*Z*)-1-((5a*R*,8a*S*)-
27
28 2,3,5,5a,6,7,8,8a-octahydro-2,2,5a,8a-tetramethylindeno[4,5-*b*]furan-4-
29
30 ylidene)propan-2-one (IC₅₀: 4.0 μM). Both compounds were isolated previously from
31
32 the brown alga *Cystoseira baccata* (Mokrini et al. 2008).
33
34
35
36
37
38
39
40

41
42 Comparison of the structure activity relations of the compounds tested in this study
43
44 unequivocally proves the importance of the α-oxo-oxime moiety that is the uniting
45
46 structural feature and the pharmacophore of compounds DBHB, NBHB and or 2,3-
47
48 BMO but is lacking in the nearly inactive active compound TT. Presence of the
49
50 phenolic rings and of the bromine substituents as observed for compounds DBHB
51
52 and NBHB enhances the enzyme inhibitory activity in comparison to compound 2,3-
53
54 BMO by factors of 3.6 (2,3-BMO vs. NBHB) and 10.3 (2,3-BMO vs. DBHB) whereas
55
56 replacement of the α-oxo-oxime group by an α-oxo-amino group as present in TT
57
58 causes a dramatic drop in activity (by a factor of > 34 compared to the closest
59
60 structural analogue NBHB, **Fig 1**). Interestingly, the differential degree of inhibition of

1
2
3 blue mussel phenoloxidase by DBHB, NBHB and TT is paralleled by their antifouling
4 activity against larvae of the barnacle *Amphibalanus improvisus* that was reported
5 earlier (Ortlepp et al., 2007). In a settling bioassay DBHB proved to be the most
6 active compound with an IC₅₀ value between 1 – 10 µM followed by NBHB (IC₅₀: 10 –
7 100 µM) whereas TT (4) was inactive up to 100 µM (2,3-BMO had not been tested in
8 the original experiment with *B. improvisus*) (Ortlepp et al., 2007) thus indicating that
9 the *in vitro* inhibition of phenoloxidase as documented in this study is apparently also
10 of relevance under *in vivo* conditions. Hemibastadin derivatives as analyzed in this
11 study are not the only sponge-derived compounds featuring an α-oxo-oxime group
12 known from nature. Besides other compounds belonging to the bastadin family we
13 were previously able to show that for example aplysamine-2 isolated from the sponge
14 *Pseudoceratina purpurea* and psammaphin A obtained from the sponge *Aplysinella*
15 *rhax* likewise suppress settling of barnacle larvae at concentrations similar to those of
16 bastadin derivatives or of DBHB (Ortlepp et al., 2007).
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38

39 No enzyme inhibitory studies were conducted for the latter compounds. However,
40 based on the obvious structural similarities with DBHB, especially with regard to the
41 important α-oxo-oxime group it is very probable that these as well as other natural
42 products featuring this pharmacophoric group (see e.g. Blunt et al., 2009 and
43 previous reviews in this series) will likewise inhibit blue mussel phenoloxidase.
44
45
46
47
48
49
50
51
52

53 Inhibition of blue mussel phenoloxidase by DBHB, NBHB and 2,3-BMO analyzed in
54 this study is reflected by their ability to complex copper(II) ions (e.g. as shown for
55 DBHB in **Fig. 5**) whereas compound TT failed to do this. Complexation of copper(II)
56 ions was also demonstrated in this study for a naturally occurring mixture of bastadin
57 derivatives as present in a crude extract of the sponge *I. basta*. Comparative X-ray
58
59
60

1
2
3 structure determinations of DBHB and the dinuclear copper(II) complex
4
5 (DBHB(DMF)CuCl₂)₂ · 4CH₂Cl₂ as well as the comparison with structurally related
6
7 complexes characterize DBHB as N,O-bidentate chelating ligand with strong binding
8
9 potential for copper(II).
10
11

12
13
14
15 In conclusion: This study demonstrates for the first time the strong inhibition of blue
16
17 mussel phenoloxidase by DBHB and by structurally related compounds at
18
19 submicromolar to low micromolar concentrations. Additional experimental evidence
20
21 obtained earlier with barnacle larvae (Ortlepp et al., 2007) furthermore indicates that
22
23 inhibition of the target molecule phenoloxidase extends also to the *in vivo* level. We
24
25 provide evidence for the α -oxo-oxime group as the decisive pharmacophore of the
26
27 studied molecules that is responsible for the complexation of copper(II) ions as
28
29 shown by X-ray analysis of a copper(II) complex of DBHB. The data presented finally
30
31 suggest that the documented strong inhibition of blue mussel phenoloxidase is
32
33 mechanistically due to complexation of copper(II) ions that are present within the
34
35 catalytical centre of the enzyme.
36
37
38
39
40
41
42

43 **Acknowledgement:** Financial support of the BMBF to Peter Proksch under the
44
45 project *BiotecMarin* is gratefully acknowledged. We thank the Industrieclub
46
47 Düsseldorf e.V. for financial support of Mirko Bayer and Mrs Eleonore Hammes for
48
49 technical assistance during the X-ray diffraction experiments.
50
51
52
53
54
55
56
57
58
59
60

Literature

Aladaileh S, Rodney P, Nair SV, Raftos DA (2007). Characterization of phenoloxidase activity in Sydney rock oysters (*Saccostrea glomerata*). Comparative Biochemistry and Physiology, Part B. 148: 470-480

Asokan R, Arumugam M, Mullainadhan P (1997). Activation of prophenoloxidase in the plasma and haemocytes of the marine mussel *Perna viridis* Linnaeus. Developmental and Comparative Immunology. 21: 1-12

Blunt J.W., Copp B.R., Hu W.-P.H., Munro M.H.G., Northcote P.T., Prinsep M.R. (2009). Marine natural products. Nat. Prod. Rep. 26: 170 - 244

Chen JS, Wei C, Rolle RS, Otwell WS, Balaban MO, Marshall MR (1991). Inhibitory Effect of Kojic Acid on Some Plant and Crustacean Polyphenol Oxidases. J. Agric. Food Chem. 39: 1396-1401

Cheung PJ, Ruggieri GD, Nigrelli RF (1977). A New Method for Obtaining Barnacle Cement in the Liquid State for Polymerization Studies. Marine Biology. 43: 157-163

Cima, F., Ballarin, L., 2000. Tributyltin induces cytoskeletal alterations in the colonial ascidian *Botryllus schlosseri* phagocytes via interaction with calmodulin. Aquat. Toxicol. 48: 419–429.

Cima, F., Ballarin, L., Bressa, G., Burighel, P., 1998a. Cytoskeleton alterations of tributyltin (TBT), in lunicante phagocytes. Ecotoxicol. Environ. Saf. 40, 160–165.

1
2
3
4
5 Cima, F., Spinazzi, R., Ballarin, L., 1998b. Possible tributyltin–calmodulin interaction
6 in morphofunctional alterations of ascidian phagocytes. *Fresenius Environ. Bull.* 7,
7
8 396–401.
9
10

11
12
13
14
15 Clare AS (1996). Signal transduction in barnacle settlement: calcium re-visited.
16
17 *Biofouling*. 10: 141-159
18

19
20
21
22 Clare AS, Hoeg JT (2008). *Balanus amphitrite* or *Amphibalanus amphitrite*? A note
23
24 on barnacle nomenclature. *Biofouling*. 24: 55-57
25

26
27
28
29 Decker H, Dillinger R, Tuczek F (2000). Wie funktioniert die Tyrosinase? Neue
30
31 Einblicke aus Modellchemie und Strukturbiologie. *Angew. Chem.* 112: 1656-1660
32

33
34
35
36 Hellio C, Bourgougnon N, Le Gal Y (2000). Phenoloxidase (E.C. 1.14.18.1) from
37
38 *Mytilus edulis* byssus gland: purification, partial characterization and application for
39
40 screening products with potential antifouling activities. *Biofouling*. 16: 235-244
41
42

43
44
45
46 Fusetani N (2004). Biofouling and antifouling. *Nat Prod. Rep.* 21: 94-104
47

48
49
50
51 Hellio C, Bado-Nilles A, Gagnaire B, Renault T, Thomas-Guyon H (2007).
52
53 Demonstration of a true phenoloxidase activity and activation of a ProPO cascade in
54
55 Pacific Oyster, *Crassostrea gigas* (Thunberg) in vitro. *Fish & Shellfish Immunology*.
56
57 22: 433-440
58
59
60

1
2
3 Hellio C, Maréchal JP, Da Gama BAP, Pereira R, Clare AS (2009) Natural marine
4 products with antifouling activities In: Advances in marine antifouling coatings and
5 technologies. Woodshead, Cambridge, pp 572–622
6
7
8
9

10
11
12 Hellio C, Yebra DM (2009). Legislation affecting antifouling products. In: Advances. In
13 marine antifouling coatings and technologies. Woodshead, Cambridge, pp 240-260
14
15
16
17

18
19
20
21
22 Kamino K (2001). Novel barnacle underwater adhesive protein is a charged amino
23 acid-rich protein constituted by a Cys-rich repetitive sequence. *Biochem J.* 356: 503-
24
25
26
27 507
28
29
30

31
32 Kapoor P, Pathak A, Kapoor R, Venugopalan P, Corbella M, Rodríguez M, Robles J,
33 Llobet A (2002). Structural, Electronic, and Magnetic Consequences of O-Carbonyl
34 vs O-Alkoxy Ester Coordination in New Dicopper Complexes Containing the $\text{Cu}_2(\mu\text{-Cl})_2$
35 Core. *Inorg. Chem.* 41: 6153-6160
36
37
38
39
40
41
42

43
44 Kapoor P, Pathak A, Prabhjot K, Venugopalan P, Kapoor R, (2004). Steric control of
45 coordination: Unusual coordination mode of dimethylpyridine-2,6-dicarboxylate in a new
46 dinuclear copper(II) complex $[(\text{dmpc})(\text{Cl})(\mu\text{-Cl})\text{Cu}_2(\mu\text{-Cl})(\text{Cl})(\text{dmpc})]$ and reversal of the
47 coordination mode in $[\text{Cu}(\text{dmpc})(\text{H}_2\text{O})_3](\text{ClO}_4)_2$. *Trans. Met. Chem.* 29: 251-258.
48
49
50
51
52
53

54
55 Kim D, Park J, Kim J, Han C, Yoon J, Kim N, Seo J, Lee C (2006). Flavonoids as
56 Mushroom Tyrosinase Inhibitors: A Fluorescence Quenching Study. *J. Agric. Food*
57
58
59
60 Chem. 54: 935-941

1
2
3 Kim YJ, Uyama H (2005). Tyrosinase inhibitors from natural and synthetic sources:
4 structure, inhibition mechanism and perspective for the future. Cell. Mol. Life Sci. 62:
5
6 1707-1723
7
8

9
10
11
12 Kubo I, Kinst-Hori I, Chaudhuri SW, Kubo Y, Sánchez Y, Ogura T (2000). Flavonols
13 from *Heterotheca inuloides*: Tyrosinase Inhibitory Activity and Structural Criteria.
14
15 Bioorganic and Medicinal Chemistry. 8: 1749-1755
16
17

18
19
20
21
22 Lau M (1991). Tributyltin antifouling: a treat to the Hong Kong marine environment.
23
24 Arch Environ Contam Toxicol. 20: 299-304
25
26

27
28
29 Liu D, Maguire R, Lau Y, Pacepavicius G, Okamura H, Aoyama I (1997).
30 Transformation of the new antifouling compound Iragarol 1051 by *Phanerochaete*
31 *chrysosporium*. Water Res. 31: 2363-2369
32
33
34

35
36
37
38 Marechal JP, Hellio C (2009). Challenges for the development of new non-toxic
39 antifouling solutions. Int J Mol Sci. 10: 4623-37
40
41
42

43
44
45
46 Mokrini R, Mesaoud MB, Daoudi M, Hellio C, Maréchal JP, El Hattab M, Ortalo-
47 Magné A, Piovetti L, Culioli G (2008). Meroditerpenoids and Derivatives from the
48
49 Brown Alga *Cystoseira baccata* and Their Antifouling Properties. J Nat Prod. 71:
50
51 1806-1811
52
53
54
55
56
57
58
59
60

1
2
3 Nagasawa T, Utagawa T, Goto J, Kim CJ, Tani Y, Kumagai H, Yamada H (1981).
4
5 Syntheses of L-Tyrosine-Related Amino Acids by Tyrosine Phenol-lyase of
6
7 Citrobacter intermedius. Eur J Biochem. 117: 33-40
8
9

10
11
12
13
14
15 Olivares C, García-Borrón J, Solano F (2002). Identification of Active Site Residues
16
17 Involved in Metal Cofactor Binding and Stereospecific Substrate Recognition in
18
19 Mammalian Tyrosinase. Implications of the Catalytic Cycle. Biochemistry. 41: 679-
20
21 686
22
23

24
25
26
27 Ortlepp S, Sjögren M, Dahlström M, Weber H, Ebel R, Edrada RA, Thoms C, Schupp
28
29 P, Bohlin L, Proksch P (2007). Antifouling Activity of Bromotyrosine-Derived Sponge
30
31 Metabolites and Synthetic Analogues. Mar Biotechnol. 9: 776-785
32
33

34
35
36 Pereira M, Ankjaergaard C (2009). Legislation affecting antifouling products. In:
37
38 Advances. In marine antifouling coatings and technologies. Woodshead, Cambridge,
39
40 pp 240-260
41
42

43
44
45
46
47
48 Rittschof D, Maki J, Mitchell R, Costlow JD (1986). Ion and neuropharmacological
49
50 studies of barnacle settlement. Neth J Sea Res. 20: 269-275
51
52

53
54
55 Sheldrick GM (1990). SHELXS97. Program for the Solution of Crystal Structures,
56
57 University of Göttingen, Germany
58
59
60

1
2
3 Sheldrick GM (1997). SHELXL97. Program for the Refinement of Crystal Structures,
4
5 University of Göttingen, Germany
6
7

8
9
10 Sommerer SO, Westcott BL, Jircitano AJ, Abboud KA (1995). The synthesis and
11
12 structure of two novel metal-di-2-pyridyl ketone oxime dimers. *Inorganica Chimica*
13
14 *Acta*, 238: 149-153
15
16

17
18
19 Suci PA, Geesey GG (2000). Influence of Sodium Periodate and Tyrosinase on
20
21 Binding of Alginate and Adlayers of *Mytilus edulis* Foot Protein 1. *Journal of Colloid*
22
23 *and Interface Science*. 230: 340-348
24
25

26
27
28 Tsoukatou M, Hellio C, Vagias C, Harvala C, Roussis V (2002). Chemical defense
29
30 and antifouling activity of three Mediterranean sponges of the genus *Ircinia*. *Z*
31
32 *Naturforsch* 57:161–171
33
34

35
36
37 Tsoukatou M, Maréchal JP, Hellio C, Novaković I, Tufegdžic S, Sladić D, Gašić MJ,
38
39 Clare AS, Vagias C, Roussis V (2007) Evaluation of the activity of the sponge
40
41 metabolites avarol and avarone and their synthetic derivatives against fouling micro-
42
43 and macroorganisms. *Molecules* 12:1022–1034
44
45
46
47

48
49
50 Tujula N, Radford J, Nair SV, Raftor DA (2001). Effects of tributyltin and other metals
51
52 on the phenoloxidase activation system of the tunicate, *Styela plicata*. *Aquatic*
53
54 *toxicology*. 55: 191-201
55
56

57
58
59 Voulvoulis N, Scrimshaw M, Lester J (1999). Alternative antifouling biocides. *Appl*
60
Organometal Chem. 13: 135-145

1
2
3
4
5
6 Wahl N (1989) Marine epibiosis. 1 Fouling and antifouling - some basic aspects. Mar
7
8 Ecol Prog Ser 58, 175–189
9

10
11
12 Yamamoto H, Satuito CG, Yamazaki M, Natoyama K, Tachibana A, Fusetani N
13
14 (1998). Neurotransmitter blockers as antifoulants against planktonic larvae of the
15
16 barnacle *Balanus amphitrite* and the mussel *Mytilus galloprovincialis*. Biofouling. 13:
17
18 69-82
19
20
21

22
23
24 Yebra DM, Kiil S, Dam-Johansen K (2004) Antifouling technology: past, present and
25
26 future steps towards efficient and environmentally friendly antifouling coatings. Progr
27
28 Org Coating 50:75–104
29
30
31

32
33
34 Zentz F, Hellio C, Valla A, De La Broise D, Bremer G, Labia R (2002). Antifouling
35
36 Activities of N-Substituted Imides: Antimicrobial Activities and Inhibition of *Mytilus*
37
38 *edulis* Phenoloxidase. Mar Biotechnol. 4: 431-430
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Figures




Fig. 1: Inhibition of blue mussel phenoloxidase by 2,3-BMO, NBHB, DBHB and TT as indicated by the IC₅₀ values of the respective compounds obtained from *in vitro* experiments.




Fig. 2: Part of the absorption spectra of DBHB (333 μmol) with various concentrations of CuSO_4 in methanolic solution: (I) 3333 $\mu\text{mol CuSO}_4/\text{ml}$; (II) 1667 $\mu\text{mol CuSO}_4/\text{ml}$; (III) 1111 $\mu\text{mol CuSO}_4/\text{ml}$; (IV) 555 $\mu\text{mol CuSO}_4/\text{ml}$; (V) 167 $\mu\text{mol CuSO}_4/\text{ml}$; (VI) 0 $\mu\text{mol CuSO}_4/\text{ml}$




Fig. 3: Influence of the different molar ratios of DBHB and CuSO_4 on the absorption maxima at 685 nm: (I) 1:10; (II) 1:5; (III) 1:3.3; (IV) 1:1.7; (V) 1:0.5; (VI) only DBHB without Cu(II)SO_4





Fig. 4: Proposed structure of the neutral copper(II)-precipitate of DBHB



(a) DBHB


(b) (DBHB(DMF)CuCl₂)₂ · 4CH₂Cl₂

Fig. 5: Molecular structure of DBHB in the crystal (30 % probability ellipsoids) (a) and structure of the dinuclear complex di- μ -chloro-bis(6,6'-dibromohemibastadin-1)(dichloro)bis(dimethylformamide)dicopper(II) in its crystalline tetrakis(dichloromethane) solvate (25 % probability ellipsoids) (b). The directions of supramolecular association *via* hydrogen bonding are indicated by dashed lines.

Table 1 Selected bond lengths (in Å) and angles (in °) for DBHB (**1**) and (DBHB(DMF)CuCl₂)₂ · 4CH₂Cl₂ (**2**)

	1	2		1	2
Br1–C10	1.908(5)	1.892(6)	N3–C19		1.454(8)
Br2–C8	1.898(5)	1.919(5)	N3–C20		1.448(9)
Br3–C16	1.908(5)	1.903(6)	C1–C2	1.499(7)	1.497(7)
Br4–C14	1.891(5)	1.906(6)	C1–C6	1.517(7)	1.530(7)
Cu1–Cl1		2.2920(15)	C2–C3	1.504(7)	1.511(7)
Cu1–Cl2		2.6420(13)	C4–C5	1.500(8)	1.526(8)
Cu1–Cl1A		2.9384(14)	C5–C12	1.530(8)	1.517(8)
Cu1–O2		2.000(4)	C6–C7	1.397(7)	1.390(7)
Cu1–O5		1.962(4)	C6–C11	1.382(6)	1.388(8)
Cu1–N1		1.999(4)	C7–C8	1.389(6)	1.388(8)
O1–C9	1.350(6)	1.356(7)	C8–C9	1.396(7)	1.367(9)
O2–C3	1.245(5)	1.259(6)	C9–C10	1.390(7)	1.406(8)
O3–N1	1.401(6)	1.370(5)	C10–C11	1.385(7)	1.391(8)
O4–C15	1.359(6)	1.361(7)	C12–C13	1.412(8)	1.394(8)
O5–C18		1.248(7)	C12–C17	1.412(8)	1.389(8)
N1–C2	1.280(6)	1.283(7)	C13–C14	1.379(8)	1.379(8)
N2–C3	1.336(7)	1.307(7)	C14–C15	1.391(7)	1.398(8)
N2–C4	1.459(7)	1.468(7)	C15–C16	1.381(7)	1.412(9)
N3–C18		1.316(8)	C16–C17	1.388(8)	1.368(9)
	2			1	2
Cl1–Cu1–Cl2	96.82(5)		O3–N1–C2	112.3(5)	116.3(4)
Cl1–Cu1–Cl1A	87.14(5)		C6–C1–C2	114.9(4)	112.7(4)
Cl1–Cu1–O2	168.91(11)		N1–C2–C1	125.5(5)	125.4(5)
Cl1–Cu1–O5	97.41(12)		N1–C2–C3	115.0(5)	109.0(5)
Cl1–Cu1–N1	90.79(13)		C1–C2–C3	119.5(4)	125.5(5)
Cl2–Cu1–Cl1A	174.85(5)		O2–C3–N2	121.8(5)	122.9(5)
Cl2–Cu1–O2	87.87(10)		O2–C3–C2	121.6(5)	118.5(5)
Cl2–Cu1–O5	96.82(11)		N2–C3–C2	116.5(4)	118.6(5)
Cl2–Cu1–N1	94.02(12)		C3–N2–C4	124.7(5)	122.3(5)
Cl1A–Cu1–O2	87.65(11)		N2–C4–C5	116.1(3)	110.4(4)
Cl1A–Cu1–O5	85.89(11)		C4–C5–C12	110.2(5)	114.7(5)
Cl1A–Cu1–N1	82.59(12)		O5–C18–N3		124.8(5)
O5–Cu1–O2	91.98(16)		C18–N3–C19		123.0(6)
O5–Cu1–N1	165.52(17)		C18–N3–C20		120.6(6)
O2–Cu1–N1	78.83(16)		C19–N3–C20		116.3(6)
Cu1–O2–C3	113.8(3)				
Cu1–N1–C2	118.9(3)				

Symmetry code: A 1.5–x, 0.5–y, 2–z

Table 2 The hydrogen bond geometries (in Å and °) for DBHB (**1**) and (DBHB(DMF)CuCl₂)₂ · 4CH₂Cl₂ (**2**)

D–H···A	D–H	H···A	angle	D···A
Compound 1				
O1–H1···O2A	0.83(5)	1.92(5)	148(5)	2.663(5)
O3–H3···O2 B	0.80(6)	2.29(6)	154(6)	3.031(6)
O4–H4···Br4	0.82	2.60	120.5	3.099(4)
Compound 2				
N2–H2···Cl2C	0.88	2.42	152.4	3.228(5)
O1–H1···Cl2D	0.84	2.35	146.2	3.080(4)
O3–H3···O5E	0.84	2.58	112.3	3.003(5)
O4–H4···Cl2F	0.84	2.48	137.0	3.151(4)

Symmetry codes: A 1–x, 1–y, 1–z; B 1–x, y–0.5, 0.5–z; C x, 1–y, z–0.5; D 1.5–x, y–0.5, 1.5–z; E 1.5–x, 0.5–y, 2–z; F 1–x, y, 1.5–z

Table 3: Summary of crystal data, details of intensity measurements and structure refinements of DBHB and ((DBHB(DMF)CuCl₂)₂ · 4CH₂Cl₂).

	C ₁₇ H ₁₄ Br ₄ N ₂ O ₄	C ₄₀ H ₄₂ Br ₈ Cl ₄ Cu ₂ N ₆ O ₁₀ · 4CH ₂ Cl ₂
Empirical formula	C ₁₇ H ₁₄ Br ₄ N ₂ O ₄	C ₄₄ H ₅₀ Br ₈ Cl ₁₂ Cu ₂ N ₆ O ₁₀
M _r	629.90	2014.60
Crystal system	monoclinic	monoclinic
Space group	<i>P</i> 2 ₁ / <i>c</i>	<i>C</i> 2/ <i>c</i>
Z	4	4
Temperature [K]	291(2)	173(2)
Unit cell parameters		
<i>a</i> [Å]	8.8333(5)	33.0662(19)
<i>b</i> [Å]	13.9719(10)	14.7836(11)
<i>c</i> [Å]	16.2404(13))	13.9389(7)
β [°]	100.255(8)	100.841(6)
Volume [Å ³]	1999.7(2)	6692.3(7)
D _{calcd.}	2.092	2.000
Absorption coefficient	8.076	5.946
F(000)	1208	3912
Crystal size [mm ³]	0.23 x 0.22 x 0.09	0.28 x 0.25 x 0.03
Crystal colour	colourless	green
Diffractometer type	Stoe-IPDS	Stoe-IPDS
Scan mode	φ	φ
θ range for data collection	1.92-25.00	2.04-25.00
Limiting indices	-10 < <i>h</i> < 10 -16 < <i>k</i> < 16 -19 < <i>l</i> < 19	-39 < <i>h</i> < 39 -17 < <i>k</i> < 17 -16 < <i>l</i> < 15
Reflections collected	18311	43969
Reflections unique	3502	5808
Reflections observed	2391	4235
Criterion for observation	<i>I</i> > 2 σ (<i>I</i>)	<i>I</i> > 2 σ (<i>I</i>)
Completeness	0.994	0.983
Refined Parameters	256	392
R ₁ ^[a] observed	0.035	0.045
wR ₂ ^[b] , all data	0.072	0.112
Goodness-of-Fit, S ^[c]	1.01	1.13
Largest diff. peak/hole	0.63/-0.34	2.29/-1.10
CCDC-identifier	782628	782629

^[a] $R_1 = \frac{\sum ||F_o| - |F_c||}{\sum |F_o|}$; ^[b] $wR_2 = \frac{[\sum w(F_o^2 - F_c^2)^2 / \sum w(F_o^2)^2]^{1/2}}{P}$, where $w = 1/[\sigma^2(F_o^2) + (aP^2)]$ and $P = (F_o^2 + 2F_c^2)/3$; ^[c] $S = [\sum w(F_o^2 - F_c^2)^2 / (n-p)]^{1/2}$.