

Degeneracy between warm and coupled cold dark matter: A clarifying noteHermano Velten,^{1,2,*} Humberto A. Borges,^{3,4,†} and Thiago R. P. Caramês^{4,‡}¹*Departamento de Física, Universidade Federal do Espírito Santo,
Campus Goiabeiras, Vitória 29075-910, Brazil*²*CPT, Aix Marseille Université, UMR 7332, 13288 Marseille, France*³*Instituto de Física, Universidade Federal da Bahia Campus de Ondina, Salvador 40210-340, Brazil*⁴*Institute of Cosmology & Gravitation, University of Portsmouth, Dennis Sciama Building, Burnaby Road,
Portsmouth PO1 3FX, United Kingdom*

(Received 30 November 2015; published 7 March 2016)

Wei *et al.* [Phys. Rev. D **88**, 043510 (2013)] have proposed the existence of a cosmological degeneracy between warm dark matter (WDM), modified gravity and coupled cold dark matter (CDM) cosmologies at both the background expansion and the growth of density perturbation levels; i.e., corresponding cosmological data would not be able to differentiate such scenarios. Here, we will focus on the specific indistinguishability between warm dark matter plus cosmological constant (Λ) and coupled scalar field CDM scenarios. Although the statement of Wei *et al.* is true for very specific conditions we present a more complete discussion on this issue and show in more detail that these models are indeed distinguishable. We show that the degeneracy breaks down since coupled models leave a specific signature in the redshift space distortion data which is absent in the uncoupled warm dark matter cosmologies. Furthermore, we complement our claim by providing the reasons which suggest that even at nonlinear level a breaking of such apparent equivalence is also expected.

DOI: [10.1103/PhysRevD.93.063503](https://doi.org/10.1103/PhysRevD.93.063503)**I. INTRODUCTION**

The actual nature of the dark matter particle remains unknown but the standard cosmological model relies on a candidate belonging to the weakly interacting massive particle (WIMP) category [1]. In this acronym “massive” means such particles are heavy, with masses $O(\text{GeV})$; i.e., they were nonrelativistic at freeze-out. As a consequence, since this ensemble of cold dark matter (CDM) particles presents a very low velocity dispersion the pressure of this component is assumed to be irrelevant for the background expansion, i.e., $p_c = 0$.

Lighter particles $m \sim O(\text{keV})$ would have decoupled still in the nonrelativistic regime and represent the basis of the warm dark matter (WDM) [2]. Consequently, their relativistic free-streaming is able to prevent structure formation below the dwarf galaxy scale and since they have a non-negligible thermal velocity there exists a positive pressure contribution. Such pressure can play a role at both the background expansion level and the perturbative sector since it leads to a effective speed of sound. This contribution is sometimes parametrized by a constant equation of state parameter $w_w = p_w/\rho_w$.¹ Current constraints limit the dark matter equation of state to the range $|w_w| \lesssim 10^{-3}$ values [3].

The prevailing view for the dark matter sector also states that these particles should interact weakly. This means that they have very small cross sections in scattering processes. However, on large scales, dark matter could eventually transfer energy-momentum to the dark energy (DE) sector whose nature is still not known. The simplest DE candidate is a cosmological constant, but barotropic fluids and dynamical fields have been also associated to the accelerated cosmic expansion. The suggestion of a possible energy-momentum transfer between DM and DE gave rise to interacting/coupled cosmological models [4] (see a recent review in [5]). In this scenario, the densities of DM and DE are somehow connected, leading to a possible way out for the cosmic coincidence problem (see [6] and references therein).

Some cosmological observations are capable of probing the expansion rate of the Universe H . A remarkable example is the supernovae type Ia data. From the Friedmann equation however we know that H is determined by the sum of all cosmic components. Then, we are not able to distinguish the nature of each single cosmic fluid or different cosmologies which lead to the same H . Moreover, a similar situation can also happen when studying the evolution of first order matter density perturbations which give us a direct understanding of how structures grow. If some set of observations is indeed unable to distinguish the actual physics behind that observable then we are left with the so-called cosmic degeneracy problem (see for example related discussion in Refs. [7–9]).

Wei *et al.* [10] have proposed a possible equivalence between three distinct cosmological models:

- (A) WDM which does not interact with dark energy in the form of a cosmological constant Λ ;

*velten@pq.cnpq.br

†humberto@ufba.br

‡thiago.carames@port.ac.uk

¹In our notation, the subscript c refers to CDM and the subscript w to WDM.

- (B) CDM coupled to a scalar dark energy field;
- (C) A CDM model driven by a modified gravity theory parametrized in terms of an effective gravitational coupling G_{eff} .

In comparison to the minimal standard Λ CDM cosmology, both models have one additional degree of freedom. Namely, model A has the constant equation of state parameter of WDM w_w , model B has the coupling Q between dark matter and the scalar field dark energy, and model C has the constant effective gravitational coupling G_{eff} . Reference [10] argues that by setting the same background expansion H and the same density contrast δ in such models, one can find an exact mapping between quantities in both models. Therefore, once one satisfies such mapping, cosmological data for the background expansion and the growth history would not be able to differentiate specific signatures of WDM particles, dark matter–dark energy coupling or modified gravity. Such a claim has also motivated further investigations [11].

We do not focus here on the possible degeneracy involving modified gravity scenarios since this discussion also relies on screening mechanisms to recover the success of general relativity in local tests. In any case, we follow recent literature and we support the idea that modified gravity models can be distinguishable from other cosmologies, more specifically, interacting cosmologies [12,13].

The purpose of this work is twofold: (1) to extend the analysis of Ref. [10] and to enrich the discussion on the degeneracy between Λ WDM and coupled CDM scenarios and (2) to identify specific cosmological signatures which are able to break such degeneracy. We anticipate that the proposed degeneracy breaks down at linear and nonlinear levels.

In the next section we review the basic dynamics presented in Ref. [10] as well as their main outcomes. In Sec. III we present in each subsection one argument advocating that the models A and B can be in fact distinguishable. In Sec. IV, we also provide an analysis concerning the nonlinear clustering to reinforce our viewpoint. We conclude in the final section.

II. WARM DARK MATTER AND COUPLED DARK MATTER: DYNAMICS

Let us introduce a more general dynamics associated to model A in which the Universe contains only WDM and dark energy. Hence, the Friedmann equation reads

$$3H^2 = \kappa^2(\rho_w + \rho_{de}), \quad (1)$$

where $\kappa^2 = 8\pi G$. Since there is no interaction between such components

$$\dot{\rho}_w + 3H(\rho_w + p_w) = 0 \quad (2)$$

$$\dot{\rho}_{de} + 3H(\rho_{de} + p_{de}) = 0, \quad (3)$$

where the equation of state parameter of a fluid $w = p/\rho$ can even be a function of time. However, in standard cosmology one has for CDM $w_c = 0$ and for relativistic fluids $w_r = 1/3$. Restricting to model A, w_w will be a free constant and $w_{de} = -1$.

At first order, subhorizon density perturbations $\delta = \delta\rho/\rho$ in the matter fluid obey

$$\begin{aligned} \delta_w'' + \left[2 - 3(2w_w - c_{ad}^2) + \frac{H'}{H} \right] \delta_w' \\ = \frac{3}{2} \Omega_w \delta_w \left(1 - 6c_{ad}^2 + 8w_w - 3w_w^2 + \frac{k^2 c_{\text{eff}}^2}{a^2 H^2} \right). \end{aligned} \quad (4)$$

where the symbol ' (prime) means derivative with respect to $\ln a$. The effective speed of sound has been defined as $c_{\text{eff}}^2 = \delta p / \delta \rho$. For adiabatic fluids this quantity reads $c_{\text{eff}}^2 = c_{ad}^2 = p' / \rho'$. In model A, $c_{\text{eff}}^2 = w_w$. We have also defined the fractionary energy density such as $\Omega_w = \kappa^2 \rho_w / 3H^2$.

Now we describe model B with more details. For the scalar field dynamics ϕ , its energy density and pressure are, respectively,

$$\hat{\rho}_\phi = \frac{\dot{\phi}^2}{2} + V(\phi) \quad \text{and} \quad \hat{p}_\phi = \frac{\dot{\phi}^2}{2} - V(\phi), \quad (5)$$

where the symbol $\hat{}$ (hat) refers to quantities belonging to model B. A specific scalar field dynamics corresponds to a choice of the potential $V(\phi)$. Then, the corresponding Friedmann equations read

$$3\hat{H}^2 = \kappa^2(\hat{\rho}_c + \hat{\rho}_\phi). \quad (6)$$

An interaction between dark matter and dark energy corresponds to a source term in their energy conservation equations. More specifically, and for the purposes of our work, we present here the coupling adopted by Amendola [14] which was used in [10]:

$$\dot{\hat{\rho}}_c + 3H(\hat{\rho}_c + p_c) = -\kappa Q \hat{\rho}_c \dot{\phi}, \quad (7)$$

$$\dot{\hat{\rho}}_\phi + 3H(\hat{\rho}_\phi + p_\phi) = \kappa Q \hat{\rho}_c \dot{\phi}. \quad (8)$$

The new degree of freedom of the model is encoded in the dimensionless arbitrary function $Q \equiv Q(\phi)$.

The dynamics of the coupled scalar field to CDM is therefore determined by

$$\ddot{\phi} + 3\hat{H} \dot{\phi} + \frac{dV}{d\phi} = \kappa Q \hat{\rho}_c. \quad (9)$$

For the subhorizon ($k/H \gg 1$) perturbative dynamics of the coupled CDM component we have [15]

$$\hat{\delta}'_c + \left(2 + \frac{\hat{H}'}{H} - \kappa Q\phi'\right)\hat{\delta}'_c = \frac{3}{2}(1 + 2Q^2)\hat{\Omega}_c\hat{\delta}_c. \quad (10)$$

Let us review the results of Ref. [10] which are related to the equivalence between a Λ WDM scenario and the coupled CDM described in the last section. The Λ WDM model has been assumed to possess the parameters $\Omega_{w0} = 0.28$, $w_w = \text{const}$ and, of course, $w_{de} = -1$.

The main argument of Wei *et al.* is that by demanding the same background expansion and structure growth history, i.e.,

$$H = \hat{H} \quad \text{and} \quad \delta_w = \hat{\delta}_c, \quad (11)$$

there is an exact mapping between the quantities in both models. From adopting that $H = \hat{H}$ they found

$$(\kappa\phi')^2 = -3\hat{\Omega}_c - 2\frac{\hat{H}'}{\hat{H}} \quad (12)$$

$$\kappa Q\phi' = -3 - 2\frac{\hat{H}'}{\hat{H}} - \frac{\hat{\Omega}'_c}{\hat{\Omega}_c}, \quad (13)$$

which combined produce an expression for Q . The function $\hat{\Omega}_c$ is not yet known and should be in general different from the WDM density Ω_w . Then, by assuming $\delta = \hat{\delta}_c$ it is possible to find a first order differential equation for $\hat{\Omega}_c$ [Eq. (32) of [10]]. According to Ref. [10], by comparing (4) and (10), and taking the subhorizon limit ($k/H \gg 1$) of the former, one finds

$$\begin{aligned} & \left[3(2w_w - c_{ad}^2) + 3 + 2\frac{\hat{H}'}{\hat{H}} + \frac{\hat{\Omega}'_c}{\hat{\Omega}_c}\right]\delta'_w \\ &= \frac{3}{2}\delta_w[\hat{\Omega}_c(1 + 2Q^2) - \Omega_w(1 - 6c_{ad}^2 + 8w_w - 3w_w^2)]. \end{aligned} \quad (14)$$

Notice that if w_w and Ω_{w0} are specified we can obtain δ_w from Eq. (4) and therefore $\hat{\Omega}_c$ is found after integration of the above equation.

Here we point out a drawback in the analysis of Ref. [10]. Since WDM has a non-negligible c_{eff}^2 the k^2 -dependence in Eq. (4) cannot be eliminated as it was claimed. Equation (14) is therefore formally wrong because of the missing $c_{\text{eff}}^2 k^2$ term which should be indeed present in the subhorizon limit of (4).

III. DISTINGUISHING COUPLED QUINTESSENCE CDM FROM Λ WDM

We have shown in the last section that the determination of $\hat{\Omega}_c$ according to (14) is inconsistent within the subhorizon limit. However, let us assume that the freedom in choosing w_w allows one to make it as small as necessary in order to make $c_{\text{eff}}^2 k^2 / a^2 H^2 \ll 1$ for any wave number, i.e.,

taking the CDM limit. With this we are temporarily relying on the assumption that $\hat{\Omega}_c$ is well determined from (14).

In the next subsections we will discuss how to distinguish models A and B while keeping this ‘‘misleading’’ assumption.

A. Implications of $\Omega_w \neq \hat{\Omega}_c$

Part of the statistics of the matter field clustering is encoded in the power spectrum $P(k) = \delta_k^2$. One observable which is affected by the evolution of the matter density is the turnover scale k_{eq} in the matter power spectrum. This scale is set by the equality time between matter and radiation, i.e., when $\Omega_r(z_{\text{eq}}) = \Omega_m(z_{\text{eq}})$. For the standard Λ CDM this happens around $z_{\text{eq}} \sim 3200$. Modes which enter the horizon during the radiation epoch cannot grow due to the Hubble drag caused by the radiative background. Only after z_{eq} , when the background is dominated by the matter component, do perturbations grow according to $\delta \sim a$ (this is also known as the Meszaros effect). If galaxy surveys are able to probe very large volumes then k_{eq} can be used to constrain cosmological models. Unfortunately, we have so far only moderate bounds on the turnover scale provided by the WiggleZ survey $k_{\text{eq}} = 0.0160^{+0.0035}_{-0.0041} h \text{ Mpc}^{-1}$ [16].

Therefore, since we have in general $\Omega_w \neq \hat{\Omega}_c$ we expect that the k_{eq} will differ in both models.

In the specific example given in [10] $\hat{\Omega}_c$ is found from integration of Eq. (14) since some arbitrary initial condition for $\hat{\Omega}_c$ is provided. We will also show here that this choice plays a crucial role.

Let us assume two submodels for the coupled scenario, Model AI: $\hat{\Omega}_c(z_i) = 0.995$ at a redshift $z_i = 1000$;

Model AII: $\hat{\Omega}_c(z_i) = 0.28$ at a redshift $z_i = 0$.

Reference [10] adopted model AI while our model AII should be—see (10)—even more similar to the Λ WDM scenario (model B).

In model AI we find that $\hat{\Omega}_{c0} = 0.16$. During this procedure it has also been assumed that $h = 0.7$ and $w_w = 0.003$ for the WDM. We have estimated the corresponding turnover scale in this specific coupled CDM example as $k_{\text{eq}} = 0.0145h \text{ Mpc}^{-1}$ ($z_{\text{eq}} = 3265$). The latter result is the same for AI and AII. On the other hand, the supposed degenerated Λ WDM cosmology presents $k_{\text{eq}} = 0.0148h \text{ Mpc}^{-1}$ ($z_{\text{eq}} = 3180$). Although we have shown that there is indeed a difference between the models, it is unfortunately very tiny in light of current precision in determining the turnover scale.

B. Matter power spectrum on small scales

Since the free-streaming of WDM particles with masses $\sim \text{keV}$ leads to a cutoff in $P(k)$ on small scales one can wonder whether the absence of such feature in CDM coupled models could be used to distinguish models A and B. In practice, this suppression is implemented via the power spectrum definition $P_w(k) = T_{\text{lin}}^2(k)P_c(k)$ where the suppression effects are encoded in the *ad hoc* function

$T_{\text{lin}}^2(k)$ which in fact also depends on the WDM particle mass m_w [17]. Current constraints from Lyman- α observations place a bound $m_{\text{WDM}} \geq 3.3$ keV (2σ) [18]. Moreover, the WDM signature on the HI 21 cm power spectrum which shall be measured in near the future by SKA [19] has been discussed in [20].

Now, let us provide a qualitative comparison between the Λ WDM and coupled CDM cosmologies concerning the $P(k)$ construction. The matter power spectrum in coupled CDM cosmologies has been also widely studied assuming many different coupling functions [21]. One remarkable fact is that one does not observe a sharp suppression on small scales as in the WDM case. This already provides a hint at differentiating models A and B.

Another important remark here is related to the numerical simulation outcomes. In general, the standard Λ CDM model is plagued with the prediction of the large number of satellites and the cusp density profile of halos. In this sense, successful cosmologies should avoid the same CDM clustering patterns. Numerical results for coupled cosmologies usually identify specific signatures which are absent in uncoupled cases [22,23]. But in general, both coupled cosmologies and WDM seem to converge to similar predictions. However, available results for numerical simulations do not allow us to make clear statements on such distinguishability since different research groups have their own numerical suites which differ in aspects like initial conditions and merging assumptions, for instance. As far as we know, a direct comparison between coupled CDM and Λ WDM cosmologies has not yet been performed by the same group. Then, this strategy would lead to a clear picture on the indistinguishability of such models.

C. The redshift space distortions signature of coupled/interacting models

We calculate here in detail the predictions for the redshift space distortions (RSD) in models A and B. In measuring the RSD the key physical quantity is the galaxies' peculiar velocities \vec{v} which quantifies how anisotropic the galactic clustering is. The derived quantity ($f\sigma_8$) which has been widely used for constraining cosmologies captures information from both the expansion growth rate via the definition $f = d \ln \delta / d \ln a$ and the variance of the density field σ_8 .

In the Λ WDM scenario we can project the WDM energy momentum tensor $T_{w;\mu}^{\mu\nu} = 0$ parallel to its four-velocity u_w^μ such that

$$aH\delta'_w + (1+w)\Theta = 0 \quad (15)$$

where $\Theta/aH = \nabla \cdot \vec{v}$. Therefore, we find that

$$-\frac{\Theta}{aH} = \left(\frac{f}{1+w} \right) \delta_w. \quad (16)$$




FIG. 1. Ratio of the RSD predictions between the coupled CDM and the Λ WDM models (20).

For the coupled cosmology, however, we should start from

$$T_{c;\mu}^{\mu\nu} = Q^\nu. \quad (17)$$

The projection of the above equation into u_c^μ leads to

$$a\hat{H}\hat{\delta}'_c + \frac{a\hat{Q}}{\hat{\rho}_c}\delta_c + \hat{\Theta} = 0. \quad (18)$$

The corresponding velocity divergence in coupled CDM models will be then [24]

$$-\frac{\hat{\Theta}}{a\hat{H}} = \left(\hat{f} + \frac{Q}{\hat{\rho}_c\hat{H}} \right) \hat{\delta}_c. \quad (19)$$

It is worth noting that the condition $\delta = \hat{\delta}$ means $f = \hat{f}$ and therefore (19) provides a clearer distinguished prediction for the RSD measurements than for the Λ WDM case. Clearly, we have $\hat{\Theta}/\Theta \neq 1$ and therefore the degeneracy is broken at a linear level.

In order to quantify whether or not the interacting term leads to a sizable effect in the RSD measurements we calculate the ratio between Eqs. (19) and (16),

$$\frac{\hat{\Theta}}{\Theta} = \left(1 - \frac{\kappa Q \phi'}{H^2 f} \right) (1 + w_w), \quad (20)$$

and plot it in Fig. 1. This shows that the configuration adopted by Wei *et al.* (AI) leads indeed to a large discrepancy in comparison to the Λ WDM model. However, the magnitude of this effect depends on the choice of the integration constant for $\hat{\Omega}_c$ since the dashed curve (AII) leads to differences of order $\lesssim 1\%$. Such sensitivity is not reached by RSD surveys and therefore in practice the degeneracy can persist.

IV. DISTINGUISHABILITY AT THE NONLINEAR LEVEL

Differences between WDM and CDM during the final process of clustering where effects like dynamical friction, tidal stripping, and tidal heating takes place are expected [25], but such details should be investigated via numerical techniques.

The consequences at a nonlinear level of possible interactions in the dark sector have received considerable attention from cosmologists in recent years [26,27]. In these works the authors analyze the impacts of the interacting dark energy on small scales, gravitational bias between baryons and cold dark matter, halo mass function and the full nonlinear matter power spectrum obtained from simulations, comparing with a fiducial Λ CDM framework. In this context, recently some authors succeeded at finding new fitting functions that reproduce the nonlinear power spectrum provided by the N-body simulations [27]. Likewise, similar investigations looking for a potential signature of a WDM model on the nonlinear power spectrum evolution and other nonlinearity effects can be found in [17,18].

In order to reinforce our claim we extend our investigation to the nonlinear ambit, by adopting the spherical collapse mechanism. Let us note that our line of reasoning has had so far the Fourier space as standpoint; thus a glance at the nonlinear regime enriches our analysis, allowing us to check the distinguishability of both models also in the real space. For the sake of simplicity we work within a neo-Newtonian fluid description and consider the top-hat profile for the spherical overdensity. In models satisfying this condition the perturbed physical quantities are assumed to be homogeneous throughout the collapse, so that the gradients of both the density and pressure perturbations are neglected. Our aim is to show that also within this framework the equivalence between Λ WDM and coupled ϕ CDM models claimed by Wei *et al.* is not fulfilled. So let us now look at the dynamical equations describing the time evolution of the matter perturbation in these models. For the Λ WDM model the clustering component is the WDM whose equation of state is constant. In this case the spherical collapse predicts the following perturbative equation [28],

$$\begin{aligned} \ddot{\delta}_w + 2H\dot{\delta}_w - \frac{3}{2}\Omega_w(1+3w_w)(1+w_w)\delta_w \\ = \frac{3}{2}\Omega_w(1+3w_w)(1+w_w)\delta_w^2 + \frac{(4+3w_w)}{3(1+w_w)}\frac{\delta_w^2}{(1+\delta_w)}, \end{aligned} \quad (21)$$

whereas for the interacting ϕ CDM the analogue equation comprises [which can be compared with Eq. (17) of Ref. [29]]

$$\begin{aligned} \ddot{\hat{\delta}} + (2H - \kappa Q\dot{\phi})\dot{\hat{\delta}} - \left[2H\kappa Q\dot{\phi} + \frac{d}{dt}(\kappa Q\dot{\phi}) + \frac{3}{2}H^2\hat{\Omega}_c \right] \hat{\delta} \\ = \frac{3}{2}H^2\hat{\Omega}_c\hat{\delta}^2 + \frac{1}{3(1+\hat{\delta})} [4\hat{\delta}^2 - 5\kappa Q\dot{\phi}\hat{\delta}\delta + (\kappa Q\dot{\phi})^2\hat{\delta}^2]. \end{aligned} \quad (22)$$

For the purpose of giving the reader a proper idea about the role played by the nonlinear terms in both models we separate them from the linear ones by displaying them at the right side of each corresponding equation. The breaking of the alleged equivalence claimed by Wei *et al.*, at a linear level, becomes even clearer when we observe the full nonlinear equations above. By using the condition $\delta = \hat{\delta}$ and $H = \hat{H}$ a straightforward comparison between these dynamical equations is impossible. The nonlinear increments appearing in (21) are proportional to $\delta^2(1+\delta)^{-1}$ and δ^2 , while in Eq. (22) the nonlinearity effects contribute with the terms $\hat{\delta}^2(1+\hat{\delta})^{-1}$, $\hat{\delta}^2$, $\hat{\delta}\hat{\delta}(1+\hat{\delta})^{-1}$ and $\hat{\delta}^2(1+\hat{\delta})^{-1}$. It is important to notice that unlike the former two terms the latter two do not have any counterparts in Eq. (21). Furthermore these additional terms arise as direct contributions coming from the interaction assumption at nonlinear level, since they vanish automatically when the interaction is switched off ($\phi = \text{const}$), and the Λ CDM model is recovered. This feature allows us to interpret the terms $\hat{\delta}\hat{\delta}(1+\hat{\delta})^{-1}$ and $\hat{\delta}^2(1+\hat{\delta})^{-1}$ in Eq. (22) as a sign of the existence of the interactions in the dark sector and since it is expected that these contributions become more and more important as the collapse reaches its final stages, we can infer that these terms can leave imprints on the estimate of abundance of collapsed objects which shall be different from the predictions coming from Λ WDM. Therefore this difference seems to be crucial to prevent the direct mapping between both models.

In fact it is possible to confirm the nonequivalence between these two scenarios and verify that the spherical collapse is indeed affected differently in both models. In order to provide an estimation on the magnitude of such difference we have solved numerically Eqs. (21) and (22), adopting model AII for comparison. Using the same initial condition for both overdensities we found that $\delta_w^{NL}(z=0)$ and $\hat{\delta}_c^{NL}(z=0)$ differ at the 22% level. For the sake of comparison, from Fig. 1 we can infer that $\hat{\Theta}$ and Θ calculated at $z=0$ differ only at the 1% level. Therefore, the nonlinear analysis seems to be more effective in differentiating these cosmologies.

As is well known the abundance of collapsed objects in a given cosmological scenario can be appraised using the linear overdensity evaluated at the collapse redshift $\delta_{\text{crit}} \equiv \delta_{\text{linear}}(z = z_{\text{collapse}})$, the so-called *threshold*. The collapse time means the instant at which the nonlinear density enhancement blows up which in general has a strong dependence upon the amplitude of the initial

overdensity: the greater (smaller) the initial density contrast is, the earlier (later) the collapse occurs. Thus, the values of $\delta_w^{NL}(z=0)$ and $\hat{\delta}_c^{NL}(z=0)$ reported above could be arbitrarily amplified by adjusting the initial conditions until the collapse (divergent nonlinear overdensity) is reached. Thus by aiming to verify how the degeneracy breaking of the models influences the corresponding threshold values, we computed such a parameter for each model at a specific time z_c . We pursued the same redshift collapse for each perturbative equation and adjusted it to $z_c \sim 0$ by means of a meticulous choice of the initial conditions. For the Λ WDM model we set $\delta_w(z_i=1000) = 3.4467 \times 10^{-3}$ and $\delta'_w(z_i=1000) = 6.43943 \times 10^{-8}$, whereas for the coupled model we have $\delta_c(z_i=1000) = 3.779 \times 10^{-3}$ and $\delta'_c(z_i=1000) = 10^{-8}$. Both cases provide redshift collapse values at the order $z_c \sim 10^{-5}$, and both values are approximately the same, differing just in the ninth decimal place. We observed that the discrepancy in both threshold values at z_c corresponds to a $\sim 2.2\%$ level. Although quite small, the distinguishability appears again through a difference between the corresponding δ_{crit} for each aforesaid scenario. This tells us that investigating the amount of bound objects over a certain range of mass, following for instance the Press-Schechter formalism or Sheth-Tormen fitting, indicates a possible way to disrupt the supposed degeneracy between both models.

V. FINAL REMARKS

We have discussed the possible indistinguishability at background and linear order between coupled CDM-quintessence and Λ WDM scenarios presented in [10]. Such a situation is indeed quite intriguing since most of the available observational data would be unable to differentiate cosmological models.

Assuming the strategy of [10] such that the scale dependence of Eq. (4) is eliminated then, even so we were able to show that differences between such models persist. In this case the question becomes actually how sizable are such differences and whether or not observations of the structure growth history can distinguish them.

In particular the expression for the redshift space distortions in the coupled cosmology (19) has a clear signature of the interaction Q .

Finally, we have also shown in Sec. IV that the nonlinear density evolution of such models is indeed different.

ACKNOWLEDGMENTS

We acknowledge enlightening conversations with Marco Bruni, Federico Piazza, Valentina Salvatelli and David Wands. We thank CNPq for financial support. H. V. also thanks the A*MIDEX Foundation under Contract No. ANR-11-IDEX-0001-02 and H. B. thanks Fapesb.

-
- [1] G. Bertone, D. Hooper, and J. Silk, *Phys. Rep.* **405**, 279 (2005).
 - [2] P. Bode, J. P. Ostriker, and N. Turok, *Astrophys. J.* **556**, 93 (2001); M. R. Lovell, C. S. Frenk, V. R. Eke, A. Jenkins, L. Gao, and T. Theuns, *Mon. Not. R. Astron. Soc.* **439**, 300 (2014).
 - [3] C. M. Mueller, *Phys. Rev. D* **71**, 047302 (2005); E. Calabrese, M. Migliaccio, L. Pagano, G. De Troia, A. Melchiorri, and P. Natoli, *Phys. Rev. D* **80**, 063539 (2009); L. Xu and Y. Chang, *Phys. Rev. D* **88**, 127301 (2013).
 - [4] L. Amendola, *Phys. Rev. D* **62**, 043511 (2000); W. Zimdahl, D. Pavon, and L. P. Chimento, *Phys. Lett. B* **521**, 133 (2001).
 - [5] Y. L. Bolotin, A. Kostenko, O. A. Lemets, and D. A. Yerokhin, *Int. J. Mod. Phys. D* **24**, 1530007 (2015).
 - [6] H. E. S. Velten, R. F. vom Martens, and W. Zimdahl, *Eur. Phys. J. C* **74**, 3160 (2014).
 - [7] M. Kunz, *Phys. Rev. D* **80**, 123001 (2009).
 - [8] S. Carneiro and H. A. Borges, *J. Cosmol. Astropart. Phys.* **06** (2014) 010.
 - [9] A. Aviles and J. L. Cervantes-Cota, *Phys. Rev. D* **84**, 083515 (2011).
 - [10] H. Wei, J. Liu, Z. C. Chen, and X. P. Yan, *Phys. Rev. D* **88**, 043510 (2013).
 - [11] J. Shim, J. Lee, and M. Baldi, arXiv:1404.3639; J. Lee and F. Hoyle, *Astrophys. J.* **803**, 45 (2015).
 - [12] C. Wetterich, *Lect. Notes Phys.* **892**, 57 (2015).
 - [13] T. Clemson and K. Koyama, *J. Cosmol. Astropart. Phys.* **01** (2010) 010.
 - [14] L. Amendola, *Phys. Rev. D* **69**, 103524 (2004).
 - [15] L. Amendola and D. Tocchini-Valentini, *Phys. Rev. D* **66**, 043528 (2002).
 - [16] G. B. Poole *et al.*, *Mon. Not. R. Astron. Soc.* **429**, 1902 (2013).
 - [17] M. Viel, K. Markovic, M. Baldi, and J. Weller, *Mon. Not. R. Astron. Soc.* **421**, 50 (2012).
 - [18] M. Viel, G. D. Becker, J. S. Bolton, and M. G. Haehnelt, *Phys. Rev. D* **88**, 043502 (2013).
 - [19] D. J. Schwarz *et al.*, *Proc. Sci.*, AASKA14 (2015) 032 [arXiv:1501.03820]; M. J. Jarvis *et al.*, arXiv:1501.03825.
 - [20] I. P. Carucci, F. Villaescusa-Navarro, M. Viel, and A. Lapi, *J. Cosmol. Astropart. Phys.* **07** (2015) 047.
 - [21] G. Olivares, F. Atrio-Barandela, and D. Pavon, *Phys. Rev. D* **74**, 043521 (2006); S. Lee, G.-C. Liu, and K.-W. Ng, *Phys. Rev. D* **73**, 083516 (2006); S. Calogero and H. Velten, *J. Cosmol. Astropart. Phys.* **11** (2013) 025; D. R. Castro, H. Velten, and W. Zimdahl, *J. Cosmol. Astropart. Phys.* **06** (2012) 024; R. F. vom Martens, W. S. Hipólito-Ricardi, and W. Zimdahl, *J. Cosmol. Astropart. Phys.* **08** (2014) 004;

- H. Velten, H. A. Borges, S. Carneiro, R. Fazolo, and S. Gomes, *Mon. Not. R. Astron. Soc.* **452**, 2220 (2015).
- [22] M. Baldi, V. Pettorino, G. Robbers, and V. Springel, *Mon. Not. R. Astron. Soc.* **403**, 1684 (2010).
- [23] E. Carlesi, A. Knebe, G. F. Lewis, and G. Yepes, *Mon. Not. R. Astron. Soc.* **439**, 2958 (2014); P. J. Elahi, G. F. Lewis, C. Power, E. Carlesi, and A. Knebe, *Mon. Not. R. Astron. Soc.* **452**, 1341 (2015).
- [24] Y. Wang, G.-B. Zhao, D. Wands, L. Pogosian, and R. G. Crittenden, *Phys. Rev. D* **92**, 103005 (2015).
- [25] A. R. Pullen, A. J. Benson, and L. A. Moustakas, *Astrophys. J.* **792**, 24 (2014).
- [26] M. Baldi, *Mon. Not. R. Astron. Soc.* **409**, L89 (2010); M. Baldi and F. Simpson, *Mon. Not. R. Astron. Soc.* **449**, 2239 (2015); W. Cui, M. Baldi, and S. Borgani, *Mon. Not. R. Astron. Soc.* **424**, 993 (2012); J. Beyer and C. Wetterich, *Phys. Lett. B* **738**, 418 (2014); C. Penzo, A. V. Macciò, M. Baldi, L. Casarini, and J. Oñorbe, [arXiv:1504.07243](https://arxiv.org/abs/1504.07243).
- [27] S. Casas, L. Amendola, M. Baldi, V. Pettorino, and A. Vollmer, *J. Cosmol. Astropart. Phys.* **01** (2016) 045.
- [28] F. Pace, J.-C. Waizmann, and M. Bartelmann, *Mon. Not. R. Astron. Soc.* **406**, 1865 (2010).
- [29] N. C. Devi, H. A. Borges, S. Carneiro, and J. S. Alcaniz, *Mon. Not. R. Astron. Soc.* **448**, 37 (2015).