

Space-based gravitational wave observatories will be able to use eccentricity to unveil stellar-mass binary black hole formation

Han Wang (王晗)¹, Ian Harry², Alexander Nitz^{3,4} and Yi-Ming Hu (胡一鸣)^{1,*}

¹*MOE Key Laboratory of TianQin Mission, TianQin Research Center for Gravitational Physics and School of Physics and Astronomy, Frontiers Science Center for TianQin, Gravitational Wave Research Center of CNSA, Sun Yat-sen University (Zhuhai Campus), Zhuhai 519082, China*

²*University of Portsmouth, Portsmouth, PO1 3FX, United Kingdom*

³*Department of Physics, Syracuse University, Syracuse, New York 13244, USA*

⁴*Max-Planck-Institut für Gravitationsphysik (Albert-Einstein-Institut), D-30167 Hannover, Germany*



(Received 26 October 2023; accepted 28 February 2024; published 25 March 2024)

The measurement of eccentricity would provide strong constraints on the formation channel of stellar-mass binary black holes. However, current ground-based gravitational wave detectors will, in most cases, not be able to measure eccentricity due to orbital circularization. Space-based observatories, in contrast, can determine binary eccentricity at 0.01 Hz to $e_{0.01} \gtrsim \mathcal{O}(10^{-4})$. Directly observing stellar-mass binary black holes with space-based observatories remains a challenging problem. However, observing such systems with ground-based detectors allows the possibility to identify the same signal in archival data from space-based observatories in the years previous. Since ground-based detectors provide little constraints on eccentricity, including eccentricity in the archival search will increase the required number of filter waveforms for the archival search by 5 orders of magnitudes [from $\sim \mathcal{O}(10^3)$ to $\sim \mathcal{O}(10^8)$], and will correspondingly need $\sim 8 \times 10^5$ core hours (and $\sim 10^5$ GB of memory), even for a mild upper limit on eccentricity of 0.1. In this work, we have constructed the first template bank for an archival search of space-based gravitational wave detectors, including eccentricity. We have demonstrated that, even though the inclusion of eccentricity brings extra computational burden, an archival search including eccentricity will be feasible in the time frame of planned space-based observatories and will provide strong constraints on the eccentricities of stellar-mass binary black holes.

DOI: [10.1103/PhysRevD.109.063029](https://doi.org/10.1103/PhysRevD.109.063029)

I. INTRODUCTION

Stellar-mass black holes (sBBHs) detected before 2015 were mainly observed through x-ray binaries [1,2], with measured masses $\lesssim 20M_{\odot}$ [3]. The first gravitational wave (GW) signal GW150914 observed by Laser Interferometer Gravitational Wave Observatory (LIGO) and Virgo has been identified as the coalescence of sBBHs with component masses $36^{+5}_{-4}M_{\odot}$ and $29^{+4}_{-4}M_{\odot}$ [4]. The observed masses posed a significant challenge to our understanding of the formation mechanism of sBBHs [5]. To date, nearly 100 sBBH mergers have been reported, many of them as heavy as GW150914 [6,7]. With the accumulation of GW observations, numerous models have been proposed to explain the formation of these sBBHs [8].

The eccentricity of a sBBH system is a key probe in unveiling the system's formation mechanism. However, among all GW detections, none was claimed to have measurable eccentricity (eccentricity at 10 Hz $e_{10} \gtrsim 0.1$) [9–11]

until GW190521, which some argue could be eccentric [12–14]. The sensitive frequency band of current ground-based detectors makes them only capable of observing sBBHs seconds before coalescence. Advanced LIGO/Virgo can measure the eccentricity for binaries with $e_{10} \gtrsim 0.05$ [15], but most sBBHs cannot retain eccentricity that high because of orbital circularization due to gravitational wave emission before entering the ground-based frequency band [16]. Therefore, it is challenging for ground-based detectors to distinguish and identify the formation channels of sBBHs [17].

Space-based GW observatories, like TianQin [18] and Laser Interferometer Space Antenna (LISA) [19], offer a promising solution to this question. They have longer baselines than their ground-based counterparts and therefore are sensitive in a lower frequency band and could observe sBBHs for years. This makes space-based observatories capable of precise mass measurements and unveiling the evolution of eccentricity and spin of sBBH sources [20–25]. For example, eccentricity evolves as $e \sim e_i (f/f_i)^{-19/18}$ at leading order [26]. If the GW of a

*huyiming@mail.sysu.edu.cn




FIG. 1. Predicted eccentricity distributions from different evolution models. The black dots and error bars represent the median values and 50% credible intervals, respectively. The vertical solid (dashed) lines indicate the minimum (maximum) detectable eccentricities of different GW observatories.

binary system evolves to the ground-based detector frequency band at $f \gtrsim 1$ Hz with eccentricity equal to 10^{-3} , the system has a significantly larger eccentricity, $e_i \sim 0.1$, at a frequency $f_i \sim 0.01$ Hz, which is a typical sensitive frequency for space-based observatories.

Figure 1 shows eccentricity distributions predicted by different evolution models. sBBHs formed in isolation are likely to have $e_{0.01} \lesssim 10^{-3}$ [27,28]. sBBHs dynamically formed in globular clusters and subsequently ejected into the field have similar distributions, with $e_{0.01} \lesssim 10^{-2}$ [28,29]. However, sBBHs that evolve inside clusters can retain a high eccentricity with $e_{0.01} \gtrsim 10^{-2}$ [30], and eccentricities can reach extreme values ($e_{0.01} \sim 1$) for systems involved in various triplets [31–33] or in active galactic nuclei (AGN) disks [34]. Space-based observatories have the capability to detect eccentricities $e_{0.01} \gtrsim 10^{-3}$ [20,23]. Therefore, sBBH detections with space-based observatories, alongside observations with ground-based facilities, offer a unique opportunity to identify the formation channel of sBBHs.

Considering eccentricity for the sBBHs can bring additional benefits. The inclusion of eccentricity can break parameter degeneracy [35], improve the precision of measuring source distance and sky localization [36,37], and make future tests of general relativity more reliable [38,39].

Matched-filtering methods have been widely used in ground-based GW detection [40]. These searches require a suitable set of waveform filters or “template bank.” Applying this method to TianQin or LISA will be challenging because of the number of waveform templates required. An example search for compact binary mergers in LIGO/Virgo data requires $\lesssim 4 \times 10^5$ templates [41]. In contrast, Moore *et al.* [42] predicts that a bank of order 10^{30} templates would be needed to cover the whole sBBH

parameter space for LISA, far exceeding a reasonable computational cost.

It has been proposed that a search of archival data from space-based observatories, triggered by detection with ground-based facilities, can achieve the multiband detection of sBBHs [25,43–45]. Next-generation ground-based detectors, like Einstein Telescope (ET) [46] and Cosmic Explorer (CE) [47], will be able to detect GW events with signal-to-noise ratios (SNRs) $\mathcal{O}(10^{2-3})$ and will therefore place tight constraints on the source parameters, for example, measuring the chirp mass to one part in 10^6 [48]. Therefore, the parameter space of an archival search of TianQin/LISA data can be greatly reduced and the required template bank size reduced to the level of 10^4 templates [45].

However, the impact of the eccentricity on archival searches has not been explored. In this paper, for the first time, we implement a matched-filtering bank generation process for an archival search in space-based observatories incorporating eccentricity, triggered by an observation using next-generation ground-based detectors. Using GW150914 and GW190521 as examples, we find that, even though the inclusion of eccentricity would enlarge the template bank by a factor of $\sim \mathcal{O}(10^5)$, the task is still tangible. This work provides a practical solution to the realistic multiband GW observation scenario.

II. METHODOLOGY

To detect GWs by matched filtering, we use EccentricFD [26,49], a nonspinning inspiral-only frequency-domain waveform approximant with eccentricity at the initial frequency e_i valid up to 0.4, for constructing the template bank. EccentricFD includes post-Newtonian (PN) corrections up to 3.5PN order and has been included into LALSuite [50]. The eccentricity in EccentricFD is expanded to $\mathcal{O}(e^8)$ and then further expanded in e_i up to $\mathcal{O}(e_i^8)$. The parameter set follows $\lambda^\mu = (\mathcal{M}, \eta, D_L, t_c, \phi_c, \iota, \lambda, \beta, \psi, e_i)$, where $\mathcal{M} \equiv (m_1 m_2)^{3/5} (m_1 + m_2)^{-1/5}$ and $\eta \equiv (m_1 m_2) (m_1 + m_2)^{-2}$ given by the component masses m_1 and m_2 ($m_1 > m_2$) are the chirp mass and symmetric mass ratio, D_L is the luminosity distance, t_c and ϕ_c are the coalescence time and phase, ι is the inclination angle, (λ, β) are ecliptic longitude and ecliptic latitude, ψ is the polarization angle, and e_i is the eccentricity at the initial frequency f_i in the quadrupolar GW mode. For space-based observatories, f_i is determined by the evolution time T from the beginning of observation to the merger. In this work, we assume a fully continuous five-year observation for both TianQin and LISA, and the merger happens at the end of the five-year period. For $M_{\text{tot}} \lesssim 10^5 M_\odot$ and $T \gtrsim 1$ yr, the correction for f_i from the eccentricity can be neglected (see Ref. [26], Appendix E), so we will use the noncentric frequency-time relation at leading PN order in the following calculation: $f_i = (5/256)^{3/8} \pi^{-1} \mathcal{M}^{-5/8} T^{-3/8}$.

The size of the parameter space that would need to be searched in an archival search depends on the parameter estimation precision of the next-generation ground-based detectors. One can use the Fisher information matrix (FIM) Γ_{ij} to estimate the statistical uncertainties in measuring parameters. $\Gamma_{ij} = \left(\frac{\partial h}{\partial \lambda^i} \middle| \frac{\partial h}{\partial \lambda^j}\right)$, where $(h|g) \equiv 4\Re \int_0^{+\infty} \frac{\tilde{g}^*(f)\tilde{h}(f)}{S_n(f)} df$, $S_n(f)$ is the one-sided detector noise power spectral density, $\tilde{h}(f) = \tilde{h}(f, \lambda^\mu)$ is the Fourier transform of the waveform $h(t)$, and λ^μ is the parameter set. The overall FIM of a detector network is the summation of the FIM of each detector. Under the Gaussian stationary assumption, the covariance matrix can be approximated by $\Sigma = \Gamma^{-1}$, and the marginalized parameter uncertainties can be estimated as $\sigma_{\lambda^i} = \sqrt{\Sigma_{ii}}$.

Here we consider a ground-based detector network including ET and two CEs, with their sites randomly chosen. Since GW emission will cause a binary orbit to circularize over time [16], we assume that events are noneccentric in the ground-based observation window. Higher-order modes, however, will be important for ET or CE, especially given the large SNR that events visible to LISA and TianQin will have. We therefore use the non-eccentric IMRPhenomHM [51] waveform to estimate the precision with which next-generation ground detectors can measure source parameters. We choose a low-frequency cutoff of $f_{\text{low}} = 1$ Hz for both CE and ET during the calculation. This is motivated by the result that one would acquire 20% of the whole SNR between 1 and 10 Hz with ET [52]. Our estimation is consistent with previous studies [23,43,45], which show that, for a GW event that retains no eccentricity when entering the ground-based observation window, the only two parameters that space-based observatories can measure more precisely are the chirp mass \mathcal{M} and initial eccentricity e_i . Therefore, we assume that all the parameters except for chirp mass and eccentricity are known exactly when performing an archival search, and the chirp mass range is determined by the uncertainty from the network of the ET and two CEs, i.e., $\mathcal{M} \in [\mathcal{M}_0 - 10\sigma_{\mathcal{M}}, \mathcal{M}_0 + 10\sigma_{\mathcal{M}}]$. In the future, we should directly use the posterior from Bayesian inference in ground-based detectors, but for this study, the uncertainty range generated by the FIM is a reasonable and conservative estimate.

We construct a template bank using SBANK [53–55], a Python package for generating stochastic template banks for compact binaries. When generating template banks stochastically, we need to determine how much those two waveforms overlap with each other. The fitting factor (FF) is used to define the maximum “similarity” between a given waveform and the best matching template in a bank [56],

$$\text{FF}(\lambda^\mu) \equiv \max_{\lambda^{\mu'}} \frac{(h(\lambda^\mu)|h(\lambda^{\mu'}))}{\sqrt{(h(\lambda^\mu)|h(\lambda^\mu))(h(\lambda^{\mu'})|h(\lambda^{\mu'}))}}. \quad (1)$$

Here $\lambda^{\mu'}$ denotes the parameter set for a template in the bank, and λ^μ is the parameter set for the test waveform. For a template bank to be complete (or “valid”), any GW signal in its parameter range should have $\text{FF} \geq M$, where M is the minimal match. Here we set $M = 0.97$, which is a commonly used value [6,7,41].

Ground-based detectors observe the GW signal over a period of only seconds before coalescence, so that the Doppler frequency modulation from the movement at Earth’s orbit can be ignored. However, the long observation time and the orbital motion of space-based observatories make the response time dependent, and one must consider these time-dependent response terms during bank generation. Additionally, unlike ground-based detectors that have fixed arm lengths during operation, the relative spacecraft motion results in unequal arm lengths. The method of time delay interferometry (TDI) has been proposed for canceling out the laser phase noise from different arms. It constructs particular combinations to make virtual equal arm interferometers. This is further complicated when considering eccentric waveforms. Here we use the frequency-domain TDI response [57,58] and combine it with EccentricFD, which contains a set of eccentric harmonics. We follow the arm length and noise budget in Luo *et al.* [18] for TianQin, and $L = 2.5 \times 10^9$ m with noise budget from Babak *et al.* [59] for LISA. We consider the response in the A channel as an example during all the calculations in this work.

Since different eccentric harmonics have different correspondences with the Fourier frequency, we should provide a frequency cutoff during the calculation to avoid the waveform generation exceeding the valid range for a specific GW detector: $\tilde{h}_{\text{det}} = \sum_j \tilde{h}_j \times \Theta(j \cdot f_{\text{high}} - 2f) \Theta(2f - j \cdot f_{\text{low}})$, where $\Theta(x)$ is the Heaviside step function and j denotes the j th eccentric harmonic [26]. For TianQin or LISA, we have $f_{\text{low}} = \max[10^{-4} \text{ Hz}, f_0]$, $f_{\text{high}} = \min[f_{\text{ISCO}}, 1 \text{ Hz}]$, where $f_{\text{ISCO}} = (6^{3/2} \pi (m_1 + m_2))^{-1}$ is the quadrupolar frequency at innermost-stable circular orbit (ISCO).

III. STOCHASTIC TEMPLATE BANK GENERATION

If a signal has small eccentricity, it could be that a circular waveform would be sufficient to recover it. The question is, how small is small enough? We therefore use a noneccentric bank (i.e., banks of \mathcal{M} in Table I) and match it with an eccentric signal. In Fig. 2, we plot the fitting factor between the injected eccentric waveform and the template bank. As expected, the mismatch increases as eccentricity gets larger and we find that the eccentricity is distinguishable for TianQin/LISA when $e_i \gtrsim 5 \times 10^{-4}$. Many models, for example, dynamical interaction mechanisms [21,27,30], predict larger initial eccentricity at ~ 0.01 Hz. We also investigate the bias between the injected and recovered chirp mass when neglecting eccentricity, which increases




FIG. 2. The fitting factor between a noneccentric template bank and a signal with different eccentricities. The blue (green) lines denote the banks of TianQin (LISA); the solid (dashed) lines correspond to the banks of a GW150914-like (GW190521-like) scenario.

from $\lesssim 10^{-6} M_{\odot}$ at $e_i = 0$ to $\gtrsim 10^{-3} M_{\odot}$ at $e_i = 0.1$. Such systematic bias could be even larger in the full parameter space. It is therefore necessary for searches to take eccentricity into account.

In Table I we show the size of the stochastic template banks, with different parameter spaces for both GW150914- and GW190521-like sources. We first assume that all the parameters (including chirp mass) are known exactly, except for eccentricity, and thus generate a one-dimensional bank with $e_i \in [0, 0.1]$. The bank size is as large as $\mathcal{O}(10^5)$ when only searching over eccentricity and requires $\lesssim 80$ core hours (and $\lesssim 100$ GB of memory). Therefore, for TianQin/LISA, we consider $e_i = 5 \times 10^{-4}$ ($e_i = 0.1$) as the smallest distinguishable eccentricity (the upper limit by the current computational cost), which corresponds to the red solid (dashed) line in Fig. 1. Figure 3 shows the distribution of the eccentricity, which follows an e^2 cumulative distribution. It agrees with the theoretical estimate in previous studies [20], subject to Poisson fluctuation as indicated by the shaded region. We then generate a one-dimensional bank covering only a range of chirp mass. Since the range is small, \mathcal{M} appears to be uniformly distributed.

As our current result fits well with a theoretical distribution for both eccentricity and chirp mass, we can give a

TABLE I. Template bank sizes for GW150914- and GW190521-like events with different parameter spaces.

	Parameter space	GW150914-like	GW190521-like
TianQin	$e_i \in [0, 0.1]$	117202	49943
	$\mathcal{M} \in \mathcal{M}_0 \pm 10\sigma_{\mathcal{M}}$	3034	4250
LISA	$e_i \in [0, 0.1]$	100403	44867
	$\mathcal{M} \in \mathcal{M}_0 \pm 10\sigma_{\mathcal{M}}$	2070	3088




FIG. 3. The distribution of the eccentricity in the archival search template bank. The shaded regions represent the 1σ Poisson fluctuation.

reasonable estimation for the full 2D parameter space of the archival search. By assuming the 2D bank follows the same relationship as the 1D eccentric bank as the eccentricity range increases, the full 2D archival search banks are expected to have $N_T \sim \mathcal{O}(10^8)$ templates; if we consider the maximal valid range for EccentricFD, i.e., $e_i \in [0, 0.4]$, N_T will be up to $\mathcal{O}(10^9)$.

To evaluate if we have overestimated the magnitude of 2D bank size due to any degeneracy between the eccentricity and the chirp mass [60–62], we generate a 2D bank for both detectors and for both sources. Restricted by the huge computational burden, we choose to verify the estimate through a bank within a smaller eccentricity range of $e_i \in [0, 0.001]$. All 2D banks have $N_T \sim \mathcal{O}(10^4)$, which is smaller but of the same order as the direct multiplication of bank sizes that are calculated separately in their parameter spaces. Such results do not change our magnitude estimation of the full 2D archival search bank size. This indicates the challenge of computational cost: an example 2D bank with $e_i \in [0, 0.001]$ includes 13372 templates and would need ~ 80 hr for one core (and 18 GB of memory to cache waveforms) to generate. By slicing the full parameter space along eccentricity and generating the 2D bank in parallel, a bank with $N_T \sim \mathcal{O}(10^8)$ needs $\sim 8 \times 10^5$ core hours (and $\sim 10^5$ GB of memory).

To evaluate the performance of our template banks, we perform tests to quantify both the validity and redundancy. First, we randomly generate 10,000 test waveforms with parameter values drawn from within the parameter space of the bank and calculate the fitting factor for each waveform. If the bank is valid, all the test waveforms will have at least one template with which the match is larger than the minimal match threshold ($M = 0.97$). In Fig. 4 we present the histogram of the fitting factor for the 10,000 injected waveforms. The red vertical line represents the threshold $M = 0.97$, and we find that, for almost all cases, the injected waveform has a FF larger than 0.97; only 0.44% of them fall lower than 0.97.

Then we move on to test the redundancy of the generated bank. We calculate the match between every template in the




FIG. 4. Validity and redundancy test of the example 2D template bank. The histogram in purple (cyan) shows the result of the validity test (redundancy test). The vertical red line corresponds to the match criteria $M = 0.97$.

template bank. An ideal bank will have no redundancy, meaning the matches between all pairs of templates should be smaller than the minimal match threshold. In Fig. 4, following the validity test, for each template we present the histogram of the fitting factor, which is calculated on a bank that excludes the template itself. We find that only 6.22% of all templates are redundant. This brings marginal extra computational cost.

IV. SUMMARY AND DISCUSSION

Numerous studies pointed out that the eccentricity of sBBHs will play a significant role in unveiling their origin. In this paper, we demonstrate that the archival search of the sBBHs from space-based observatories is highly sensitive to the eccentricity. Furthermore, for the first time, we successfully implement a GW template bank generation process that includes eccentricity.

We generate one-dimensional template banks for either initial eccentricity or for chirp mass. The upper limit of initial eccentricity at a system five years before merger is 0.1. The range of chirp mass is determined by the estimation with the ground-based network. By extrapolating the one-dimensional bank results, we conclude that a two-dimensional eccentric bank will comprise $N_T \sim \mathcal{O}(10^{8-9})$ templates, which is $\sim \mathcal{O}(10^5)$ larger compared to

the zero eccentricity case, and will require $\sim \mathcal{O}(10^6)$ core hours [and $\sim \mathcal{O}(10^5)$ GB of memory] for the pipeline to generate it [also $\sim \mathcal{O}(10^5)$ larger compared to noncentric case]. This conclusion is verified by a small 2D bank, where the upper limit on the initial eccentricity is 0.001. Constructing and filtering a template bank of 10^{8-9} waveforms will therefore be a challenging task, but it is not outside the scope of the expected computational facilities in the late 2030s and could be further improved with additional optimization of the relevant software techniques. Our work provides a practical solution to the realistic multiband GW observation scenario, with which one can determine the formation mechanism of sBBHs with successful archival searches.

It should be noted that we use a nonspinning eccentric waveform model in the paper. It is already known that spin effects are largely negligible during the inspiral [63] phase. However, in our technique, this is not a concern at all because the spin would already be constrained by ground-based facilities. Our space-based archival search would then just search a range of chirp mass and eccentricity values, using the measured black hole spins. It is important to note though that, for both ground- and space-based detectors, more precise waveform models will be needed in the future to avoid potential systematic errors [58,64–67].

One caveat in the study is the duty cycle. We consider ET + dual CE for the ground-based detectors, whereas in reality the duty cycle cannot reach 100%; so the sky localization from realistic future networks might be worse than our calculation. Space-based observatories will also be limited by duty cycles [18,68]. We leave the detailed calculation to future studies.

ACKNOWLEDGMENTS

We are grateful to Xiang-Yu Lyu, En-Kun Li, Jian-dong Zhang, and Shuai Liu for the helpful discussions. This work has been supported by the Guangdong Major Project of Basic and Applied Basic Research (Grant No. 2019B030302001), the Natural Science Foundation of China (Grant No. 12173104), and the National Key Research and Development Program of China (No. 2020YFC2201400). I. H. acknowledges support from the UK Space Agency through Grant No. ST/X002225/1.

[1] Jerome A. Orosz, Jeffrey E. McClintock, Jason P. Aufdenberg, Ronald A. Remillard, Mark J. Reid, Ramesh Narayan, and Lijun Gou, The mass of the black hole in Cygnus X-1, *Astrophys. J.* **742**, 84 (2011).

[2] Jesus M. Corral-Santana, Jorge Casares, Teo Munoz-Darias, Franz E. Bauer, Ignacio G. Martinez-Pais, and David M. Russell, BlackCAT: A catalogue of stellar-mass black holes in x-ray transients, *Astron. Astrophys.* **587**, A61 (2016).

- [3] Feryal Özel, Dimitrios Psaltis, Ramesh Narayan, and Jeffrey E. McClintock, The black hole mass distribution in the galaxy, *Astrophys. J.* **725**, 1918 (2010).
- [4] B. P. Abbott *et al.*, Observation of gravitational waves from a binary black hole merger, *Phys. Rev. Lett.* **116**, 061102 (2016).
- [5] B. P. Abbott *et al.*, Astrophysical implications of the binary black-hole merger GW150914, *Astrophys. J. Lett.* **818**, L22 (2016).
- [6] R. Abbott *et al.*, GWTC-3: Compact binary coalescences observed by LIGO and Virgo during the second part of the third observing run, *Phys. Rev. X* **13**, 041039 (2023).
- [7] Alexander H. Nitz, Sumit Kumar, Yi-Fan Wang, Shilpa Kastha, Shichao Wu, Marlin Schäfer, Rahul Dhurkunde, and Collin D. Capano, 4-OGC: Catalog of gravitational waves from compact-binary mergers, *Astrophys. J.* **946**, 59 (2023).
- [8] Leor Barack *et al.*, Black holes, gravitational waves and fundamental physics: A roadmap, *Classical Quantum Gravity* **36**, 143001 (2019).
- [9] B. P. Abbott *et al.*, Search for eccentric binary black hole mergers with Advanced LIGO and Advanced Virgo during their first and second observing runs, *Astrophys. J.* **883**, 149 (2019).
- [10] Isobel M. Romero-Shaw, Paul D. Lasky, and Eric Thrane, Searching for eccentricity: Signatures of dynamical formation in the first gravitational-wave transient catalogue of LIGO and Virgo, *Mon. Not. R. Astron. Soc.* **490**, 5210 (2019).
- [11] Shichao Wu, Zhoujian Cao, and Zong-Hong Zhu, Measuring the eccentricity of binary black holes in GWTC-1 by using the inspiral-only waveform, *Mon. Not. R. Astron. Soc.* **495**, 466 (2020).
- [12] R. Abbott *et al.*, Properties and astrophysical implications of the $150M_{\odot}$ binary black hole merger GW190521, *Astrophys. J. Lett.* **900**, L13 (2020).
- [13] Isobel M. Romero-Shaw, Paul D. Lasky, Eric Thrane, and Juan Calderon Bustillo, GW190521: Orbital eccentricity and signatures of dynamical formation in a binary black hole merger signal, *Astrophys. J. Lett.* **903**, L5 (2020).
- [14] V. Gayathri, J. Healy, J. Lange, B. O'Brien, M. Szczepanczyk, Imre Bartos, M. Campanelli, S. Klimentko, C. O. Lousto, and R. O'Shaughnessy, Eccentricity estimate for black hole mergers with numerical relativity simulations, *Nat. Astron.* **6**, 344 (2022).
- [15] Marcus E. Lower, Eric Thrane, Paul D. Lasky, and Rory Smith, Measuring eccentricity in binary black hole inspirals with gravitational waves, *Phys. Rev. D* **98**, 083028 (2018).
- [16] P. C. Peters, Gravitational radiation and the motion of two point masses, *Phys. Rev.* **136**, B1224 (1964).
- [17] Carl L. Rodriguez, Pau Amaro-Seoane, Sourav Chatterjee, Kyle Kremer, Frederic A. Rasio, Johan Samsing, Claire S. Ye, and Michael Zevin, Post-Newtonian dynamics in dense star clusters: Formation, masses, and merger rates of highly-eccentric black hole binaries, *Phys. Rev. D* **98**, 123005 (2018).
- [18] Jun Luo *et al.*, TianQin: A space-borne gravitational wave detector, *Classical Quantum Gravity* **33**, 035010 (2016).
- [19] Pau Amaro-Seoane, Heather Audley, Stanislav Babak, John Baker, Enrico Barausse, Peter Bender, Emanuele Berti, Pierre Binetruy, Michael Born, Daniele Bortoluzzi *et al.*, Laser interferometer space antenna, [arXiv:1702.00786](https://arxiv.org/abs/1702.00786).
- [20] Atsushi Nishizawa, Emanuele Berti, Antoine Klein, and Alberto Sesana, eLISA eccentricity measurements as tracers of binary black hole formation, *Phys. Rev. D* **94**, 064020 (2016).
- [21] Atsushi Nishizawa, Alberto Sesana, Emanuele Berti, and Antoine Klein, Constraining stellar binary black hole formation scenarios with eLISA eccentricity measurements, *Mon. Not. R. Astron. Soc.* **465**, 4375 (2017).
- [22] Xian Chen and Pau Amaro-Seoane, Revealing the formation of stellar-mass black hole binaries: The need for deci-hertz gravitational wave observatories, *Astrophys. J. Lett.* **842**, L2 (2017).
- [23] Shuai Liu, Yi-Ming Hu, Jian-dong Zhang, and Jianwei Mei, Science with the TianQin observatory: Preliminary results on stellar-mass binary black holes, *Phys. Rev. D* **101**, 103027 (2020).
- [24] Riccardo Buscicchio, Antoine Klein, Elinore Roebber, Christopher J. Moore, Davide Gerosa, Eliot Finch, and Alberto Vecchio, Bayesian parameter estimation of stellar-mass black-hole binaries with LISA, *Phys. Rev. D* **104**, 044065 (2021).
- [25] Antoine Klein, Geraint Pratten, Riccardo Buscicchio, Patricia Schmidt, Christopher J. Moore, Eliot Finch, Alice Bonino, Lucy M. Thomas, Natalie Williams, Davide Gerosa, Sean McGee, Matt Nicholl, and Alberto Vecchio, The last three years: Multiband gravitational-wave observations of stellar-mass binary black holes, [arXiv:2204.03423](https://arxiv.org/abs/2204.03423).
- [26] Nicolas Yunes, K. G. Arun, Emanuele Berti, and Clifford M. Will, Post-circular expansion of eccentric binary inspirals: Fourier-domain waveforms in the stationary phase approximation, *Phys. Rev. D* **80**, 084001 (2009).
- [27] I. Kowalska, T. Bulik, K. Belczynski, M. Dominik, and D. Gondek-Rosinska, The eccentricity distribution of compact binaries, *Astron. Astrophys.* **527**, A70 (2011).
- [28] Katelyn Breivik, Carl L. Rodriguez, Shane L. Larson, Vassiliki Kalogera, and Frederic A. Rasio, Distinguishing between formation channels for binary black holes with LISA, *Astrophys. J. Lett.* **830**, L18 (2016).
- [29] Carl L. Rodriguez, Sourav Chatterjee, and Frederic A. Rasio, Binary black hole mergers from globular clusters: Masses, merger rates, and the impact of stellar evolution, *Phys. Rev. D* **93**, 084029 (2016).
- [30] Johan Samsing and Daniel J. D'Orazio, Black hole mergers from globular clusters observable by LISA I: Eccentric sources originating from relativistic N -body dynamics, *Mon. Not. R. Astron. Soc.* **481**, 5445 (2018).
- [31] Fabio Antonini and Hagai B. Perets, Secular evolution of compact binaries near massive black holes: Gravitational wave sources and other exotica, *Astrophys. J.* **757**, 27 (2012).
- [32] Fabio Antonini, Silvia Toonen, and Adrian S. Hamers, Binary black hole mergers from field triples: Properties, rates and the impact of stellar evolution, *Astrophys. J.* **841**, 77 (2017).
- [33] Michael Zevin, Johan Samsing, Carl Rodriguez, Carl-Johan Haster, and Enrico Ramirez-Ruiz, Eccentric black hole mergers in dense star clusters: The role of binary-binary encounters, *Astrophys. J.* **871**, 91 (2019).
- [34] J. Samsing, I. Bartos, D. J. D'Orazio, Z. Haiman, B. Kocsis, N. W. C. Leigh, B. Liu, M. E. Pessah, and H. Tagawa, AGN

- as potential factories for eccentric black hole mergers, *Nature (London)* **603**, 237 (2022).
- [35] Zeyuan Xuan, Smadar Naoz, and Xian Chen, Detecting accelerating eccentric binaries in the LISA band, *Phys. Rev. D* **107**, 043009 (2023).
- [36] Baosan Sun, Zhoujian Cao, Yan Wang, and Hsien-Chi Yeh, Parameter estimation of eccentric inspiraling compact binaries using an enhanced post circular model for ground-based detectors, *Phys. Rev. D* **92**, 044034 (2015).
- [37] Tao Yang, Rong-Gen Cai, Zhoujian Cao, and Hyung Mok Lee, Eccentricity of long inspiraling compact binaries sheds light on dark sirens, *Phys. Rev. Lett.* **129**, 191102 (2022).
- [38] Pankaj Saini, Marc Favata, and K. G. Arun, Systematic bias on parametrized tests of general relativity due to neglect of orbital eccentricity, *Phys. Rev. D* **106**, 084031 (2022).
- [39] Sajad A. Bhat, Pankaj Saini, Marc Favata, and K. G. Arun, Systematic bias on the inspiral-merger-ringdown consistency test due to neglect of orbital eccentricity, *Phys. Rev. D* **107**, 024009 (2023).
- [40] S. Babak *et al.*, Searching for gravitational waves from binary coalescence, *Phys. Rev. D* **87**, 024033 (2013).
- [41] Tito Dal Canton and Ian W. Harry, Designing a template bank to observe compact binary coalescences in Advanced LIGO's second observing run, [arXiv:1705.01845](https://arxiv.org/abs/1705.01845).
- [42] Christopher J. Moore, Davide Gerosa, and Antoine Klein, Are stellar-mass black-hole binaries too quiet for LISA?, *Mon. Not. R. Astron. Soc.* **488**, L94 (2019).
- [43] Alberto Sesana, Prospects for multiband gravitational-wave astronomy after GW150914, *Phys. Rev. Lett.* **116**, 231102 (2016).
- [44] Kaze W.K. Wong, Ely D. Kovetz, Curt Cutler, and Emanuele Berti, Expanding the LISA horizon from the ground, *Phys. Rev. Lett.* **121**, 251102 (2018).
- [45] Becca Ewing, Surabhi Sachdev, Ssohrab Borhanian, and B. S. Sathyaprakash, Archival searches for stellar-mass binary black holes in LISA data, *Phys. Rev. D* **103**, 023025 (2021).
- [46] M. Punturo, M. Abernathy, F. Acernese, B. Allen, Nils Andersson, K. Arun, F. Barone, B. Barr, M. Barsuglia, M. Beker *et al.*, The Einstein Telescope: A third-generation gravitational wave observatory, *Classical Quantum Gravity* **27**, 194002 (2010).
- [47] David Reitze, Rana X. Adhikari, Stefan Ballmer, Barry Barish, Lisa Barsotti, GariLynn Billingsley, Duncan A. Brown, Yanbei Chen, Dennis Coyne, Robert Eisenstein *et al.*, Cosmic Explorer: The U.S. contribution to gravitational-wave astronomy beyond LIGO, *Bull. Am. Astron. Soc.* **51**, 35 (2019), <https://arxiv.org/abs/1907.04833>.
- [48] Hee-Suk Cho, Improvement of the parameter measurement accuracy by the third-generation gravitational wave detector Einstein Telescope, *Classical Quantum Gravity* **39**, 085006 (2022).
- [49] E. A. Huerta, Prayush Kumar, Sean T. McWilliams, Richard O'Shaughnessy, and Nicolás Yunes, Accurate and efficient waveforms for compact binaries on eccentric orbits, *Phys. Rev. D* **90**, 084016 (2014).
- [50] LIGO Scientific Collaboration, LIGO Algorithm Library—LALSuite, free software (GPL) (2018), <https://doi.org/10.7935/GT1W-FZ16>.
- [51] Lionel London, Sebastian Khan, Edward Fauchon-Jones, Cecilio García, Mark Hannam, Sascha Husa, Xisco Jiménez-Forteza, Chinmay Kalaghatgi, Frank Ohme, and Francesco Pannarale, First higher-multipole model of gravitational waves from spinning and coalescing black-hole binaries, *Phys. Rev. Lett.* **120**, 161102 (2018).
- [52] Amber K. Lenon, Duncan A. Brown, and Alexander H. Nitz, Eccentric binary neutron star search prospects for Cosmic Explorer, *Phys. Rev. D* **104**, 063011 (2021).
- [53] <https://github.com/gwastro/sbank>.
- [54] Stanislav Babak, Building a stochastic template bank for detecting massive black hole binaries, *Classical Quantum Gravity* **25**, 195011 (2008).
- [55] Ian W. Harry, Bruce Allen, and B. S. Sathyaprakash, A stochastic template placement algorithm for gravitational wave data analysis, *Phys. Rev. D* **80**, 104014 (2009).
- [56] Jolien D. E. Creighton and Warren G. Anderson, *Gravitational-Wave Physics and Astronomy: An Introduction to Theory, Experiment and Data Analysis* (John Wiley & Sons, New York, 2011).
- [57] Sylvain Marsat, John G. Baker, and Tito Dal Canton, Exploring the Bayesian parameter estimation of binary black holes with LISA, *Phys. Rev. D* **103**, 083011 (2021).
- [58] Xiangyu Lyu, En-Kun Li, and Yi-Ming Hu, Parameter estimation of stellar mass binary black holes under the network of TianQin and LISA, *Phys. Rev. D* **108**, 083023 (2023).
- [59] Stanislav Babak, Antoine Petiteau, and Martin Hewitson, LISA sensitivity and SNR calculations, [arXiv:2108.01167](https://arxiv.org/abs/2108.01167).
- [60] Marc Favata, Chunglee Kim, K. G. Arun, JeongCho Kim, and Hyung Won Lee, Constraining the orbital eccentricity of inspiraling compact binary systems with Advanced LIGO, *Phys. Rev. D* **105**, 023003 (2022).
- [61] Eamonn O'Shea and Prayush Kumar, Correlations in parameter estimation of low-mass eccentric binaries: GW151226 & GW170608, *Phys. Rev. D* **108**, 104018 (2023).
- [62] Amber K. Lenon, Alexander H. Nitz, and Duncan A. Brown, Measuring the eccentricity of GW170817 and GW190425, *Mon. Not. R. Astron. Soc.* **497**, 1966 (2020).
- [63] Alberto Mangiagli, Antoine Klein, Alberto Sesana, Enrico Barausse, and Monica Colpi, Post-Newtonian phase accuracy requirements for stellar black hole binaries with LISA, *Phys. Rev. D* **99**, 064056 (2019).
- [64] Luc Blanchet, Gravitational radiation from post-Newtonian sources and inspiraling compact binaries, *Living Rev. Relativity* **17**, 2 (2014).
- [65] Michael Pürrer and Carl-Johan Haster, Gravitational wave-form accuracy requirements for future ground-based detectors, *Phys. Rev. Res.* **2**, 023151 (2020).
- [66] Isobel M. Romero-Shaw, Paul D. Lasky, and Eric Thrane, Signs of eccentricity in two gravitational-wave signals may indicate a subpopulation of dynamically assembled binary black holes, *Astrophys. J. Lett.* **921**, L31 (2021).
- [67] Isobel M. Romero-Shaw, Davide Gerosa, and Nicholas Loutrel, Eccentricity or spin precession? Distinguishing subdominant effects in gravitational-wave data, *Mon. Not. R. Astron. Soc.* **519**, 5352 (2023).
- [68] Pau Amaro Seoane *et al.*, The effect of mission duration on LISA science objectives, *Gen. Relativ. Gravit.* **54**, 3 (2022).